

Blood-flow restricted exercise following ankle fractures — A feasibility study

Larsen, Peter; Platzer, Oscar Just; Lollesgaard, Lærke; Pedersen, Samuel Krogh; Nielsen, Peter Kruse; Rathleff, Michael S.; Bandholm, Thomas; Jensen, Stefan Teglhuis; Elsoe, Rasmus

Published in:
Foot and Ankle Surgery

DOI (link to publication from Publisher):
[10.1016/j.fas.2021.08.010](https://doi.org/10.1016/j.fas.2021.08.010)

Creative Commons License
CC BY 4.0

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Larsen, P., Platzer, O. J., Lollesgaard, L., Pedersen, S. K., Nielsen, P. K., Rathleff, M. S., Bandholm, T., Jensen, S. T., & Elsoe, R. (2022). Blood-flow restricted exercise following ankle fractures — A feasibility study. *Foot and Ankle Surgery*, 28(6), 726-731. <https://doi.org/10.1016/j.fas.2021.08.010>


General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Blood-flow restricted exercise following ankle fractures — A feasibility study

Peter Larsen^{a,b,*}, Oscar Just Platzer^c, Lærke Lollesgaard^c, Samuel Krogh Pedersen^c, Peter Kruse Nielsen^c, Michael S. Rathleff^{a,c}, Thomas Bandholm^d, Stefan Tegllhus Jensen^b, Rasmus Elsoe^b

^a Department of Occupational Therapy and Physiotherapy, Aalborg University Hospital, Aalborg, Denmark

^b Department of Orthopaedic Surgery, Aalborg University Hospital, Aalborg, Denmark

^c Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

^d Physical Medicine & Rehabilitation Research – Copenhagen (PMR-C), Department of Physio- and Occupational Therapy, Department of Orthopaedic Surgery, Department of Clinical Research, Copenhagen University Hospital, Amager and Hvidovre, Copenhagen, Denmark

ARTICLE INFO

Article history:

Received 9 June 2021

Received in revised form 17 August 2021

Accepted 31 August 2021

Keywords:

Blood flow restricted exercise

BFRE

Ankle fracture

Rehabilitation

ABSTRACT

Background: The objective was to investigate the feasibility of blood flow restricted exercise (BFRE) as a rehabilitation modality in patients with a unilateral ankle fracture.

Methods: Feasibility study with a prospective cohort design. Inclusion criteria were above 18 years of age and unilateral ankle fractures. Exclusion criteria: history of cardiac or embolic diseases, cancer, diabetes, hypertension and family history of cardio or vascular diseases. The predefined feasibility outcome was based on three criteria regarding patients experience with participating in the BFRE protocol and the absence of any serious adverse events.

Results: Eight patients were included. Median age was 33 years (range: 23–60). All eight patients reported maximum satisfaction on the two questions regarding patient's perception of the overall experience with BFRE training and the feasibility to introduce BFRE as an intervention.

Conclusion: Early use of BFRE in patients with unilateral ankle fractures seems feasible in patients without comorbidity.

© 2021 The Author(s). Published by Elsevier Ltd on behalf of European Foot and Ankle Society. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Fractures of the ankle are very common and have an overall incidence of 169.7/100,000/year [1] - representing almost 10% of all bone fractures [2]. The primary aim when treating ankle fractures is to restore normal function of the ankle joint. The acute management of ankle fractures can either be surgical or non-surgical, depending on fracture classification and concomitant injuries to bone and ligaments [3]. Following the initial treatment, the majority of patients have a prolonged period of relative immobilization with partly or no weight bearing to allow for accurate bone union. Such a prolonged period of immobilization often results in skeletal muscle atrophy and loss of range of joint motion [4]. Even a short period of three weeks knee

immobilization in cast and no weight bearing cause a 47% decrease in muscle strength in healthy college students [5]. These deficits can negatively impact function and daily activities and the main aim of rehabilitation is to help the patient regain strength and function to the presurgical level.

Resistance training aims to build muscle mass, muscle strength and improve physical function [6]. The basic principle of progressive strength training is to progressively overload the muscle using external weight loads, which is known to increase muscle growth and strength [7]. However, in patients recovering from an ankle fracture, external weights and progressive overload may put too much stress on healing tissues and delay bone union. Therefore, “classic” progressive strength training of the lower extremity with increasing external weight loads might be contraindicated [7].

A new modality, blood flow restricted exercise (BFRE), has recently been tested in other conditions where patients do not tolerate high external loads [8]. It is characterized by muscle strength training with low external weight loads (20–30% of one-repetition maximum (1RM)) combined with a pneumatic cuff

* Corresponding author at: Department of Occupational Therapy and Physiotherapy/Orthopaedic Surgery, Aalborg University Hospital, Aalborg University, 18-22 Hobrovej, DK-9000 Aalborg, Denmark.

E-mail address: peter.larsen@rn.dk (P. Larsen).

inflation that partly reduces arterial blood flow and limits venous return, thus, elevating metabolic stimulus in the working muscles [9–16]. Due to the low external weight needed, BFRE seems useful in the rehabilitation of patients with ankle fractures, and may reduce the negative effects of immobilization.

The objective of the present pilot study was to investigate the feasibility of BFRE when used as a rehabilitation modality in patients with a unilateral ankle fracture.

2. Patients and methods

2.1. Study design

This study used a prospective cohort design to investigate the feasibility of BFRE in patients with unilateral ankle fractures, using pre-defined criteria for feasibility. Patients were included from the Department of Orthopedic Surgery, Aalborg University Hospital, Denmark and assessed for eligibility between September 15, 2020 and December 17, 2020. Outcomes were collected unblinded at baseline, after each BFRE session and after 3 weeks of BFRE.

The Danish Data Protection Agency and the Committee for Science Ethics of Northern Denmark (journal number N-20200052) approved the study, which was performed according to the principles of the Helsinki Declaration. Oral and written information about the study were provided and all patients completed a written informed consent. The reporting of the study complies with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement [17] and intervention description complies with the Consensus on Exercise Reporting Template (CERT) [18].

2.2. Participants

The inclusion criteria were above 18 years of age, patients with unilateral ankle fractures (AO classification 44). The exclusion criteria were not able to understand written and spoken Danish and patients with previous history of periphery vascular diseases, cancer, deep vein thrombosis or pulmonary embolus, arrhythmia or coronary diseases, diabetic, hypertension 160/95 mmHg and family history of cardio or vascular diseases.

2.3. Intervention

All patients were prescribed an individualized BFRE/exercise program which was supervised by a physiotherapist. The exercise program started about 14 days postoperatively and included 3 weeks of intervention with the BFRE program two times per week. Each exercise session comprises a single unilateral tension band strength training exercise; knee-extension, which was performed at about 30% of 1RM. (Please find detailed CERT in appendix 1).

The blood flow occlusion pressure was established with an air cuff (FitCuffs® V3 Performance Lower) placed on the inguinal fold with a moderate limb occlusion pressure (LOP) at 60% sustained throughout the training session.

The load was individualized and each patient exercised with a resistance of an elastic band (TheraBand® dark blue) corresponding to a relative load of about 30% 1 RM. While the patient sat on a chair, a 150 cm double elastic band was used as external resistance (Fig. 1). The knee-extension exercise was performed in 4 sets with 30, 15, 15, 15 repetitions in each set, with a tempo of 1 s concentric and 1 s eccentric contraction phase. A 30-sec rest interval was allowed between sets throughout the training protocol. When a given resistance in the elastic exercise band became too low (i.e., more than 30,15,15,15 repetitions per set could be performed), the patient was instructed to adjust the resistance in the elastic



Fig. 1. Tension band strength training exercise.

exercise band (moving the chair 30 cm. further away from the door to achieve a new resistance).

2.4. Descriptives and outcomes

Baseline characteristics including age, gender, height, weight, BMI and AO classification were obtained at the time of inclusion/baseline. Feasibility data from individual patients were collected after each BFRE session.

2.4.1. Feasibility outcome questions and criteria

1. Based on your current knowledge and experience, how likely is it that you will choose BFRE training if you experience an ankle fracture tomorrow?

This question intends to evaluate the patients overall combined experience with participating in the BFRE training following an ankle fracture and the patients' reflections regarding discomfort and insecurity during the training combined with patients perceived and expected outcome.

2. How likely are you, based on your current knowledge and experience, to recommend BFRE training to friends and family?

This question intends to evaluate the patients' perception of to what degree BFRE training can be introduced in the general population of ankle fractures.

A feasible outcome was predefined based on at least 75% of patients reporting 4 point or above on a 5-point Likert scale on both questions regarding patients experience with participating in the BFRE protocol. Feasibility outcome was evaluated after the last BFRE session. The scale for answers were as follow: much likely, likely, neither/nor, not likely, not at all likely.

Furthermore, feasibility will be claimed based on the absence of any serious adverse events (SAE). SAE was predefined as events resulting in life-threatening conditions, death, permanent

Table 1
Baseline characteristics.

Baseline characteristics of the 8 patients	
Age at follow-up, median (range)	33 (23–60)
Gender, n	
Male	3
Female	5
Fracture classification, n	
AO-44A	3
AO-44B	3
AO-44C	2
Treatment, n	
Conservative	3
Operative	5
LOP	197 (160–220)

N = number, LOP = Limb occlusion pressure.

disability, or damage. Information regarding SAE were obtained at each BFRE session.

2.4.2. Secondary outcome measurements

Acceptance of pain/discomfort during treatment was evaluated on a 5-point Likert scale by a single question after each BFRE session (very much, much, niter/nor, not, not at all):

How acceptable was it for you to have pain/discomfort during the BFRE session?

Patients evaluated perception of security of BFRE on a 5-point Likert scale after each BFRE session by a single question (very much, much, niter/nor, not, not at all):

To what degree did you experience security in today's BFRE session?

2.4.3. Muscle function outcomes

Maximum isometric knee-extension strength was tested bilateral with the patient sitting on an examination table with 60 degrees knee flexion by a strap-mounted dynamometer attached to the wall (Mecmesin AFG2500, Mecmesin Ltd, West Sussex, UK) [19]. Maximum isometric knee-extension strength was obtained at baseline and after the last BFRE session.

Adverse events/harms: An AE was defined as any undesirable experience during follow-up. If an AE result in hospitalization, prolonged inpatient hospital care, result in re-surgery, or if an AE is life-threatening, result in death, permanent disability or damage, they will be categorized as serious adverse events (SAEs). AE and SAE were obtained at each BFRE session.

2.5. Statistics

Approaches to sample size justification for pilot and feasibility trials vary. We aimed for a target sample size of 8 participants due to the descriptive design and as no efficacy testing was planned [20]. Data was assumed missing at random and no imputations were made.

3. Results

During the study period 21 patients were admitted at Aalborg University Hospital with a unilateral ankle fracture and all were assessed for eligibility. Eight patients were included in the study.

Eleven patients were excluded due to the exclusion criteria and two patients did not want to participate.

The median age was 33 years with a range of 23–60. Baseline characteristics are presented in Table 1.

3.1. Outcomes related to intervention feasibility

Based on the predefined three criteria of feasibility of BFRE as early intervention the unilateral ankle fractures the degree of feasibility was high.

All eight patients reported maximum on the two questions regarding patient's perception of the overall experience with BFRE training and then possibility to introduce BFRE training as an intervention on unilateral ankle fracture patients. No serious adverse event (SAE) was observed during the study period.

3.2. Other outcomes

The individual patient's acceptance of pain/discomfort during each BFRE training together with patient's confident status are reported in Table 2. Results indicate very high degree of patients acceptance of pain and discomfort and patients confident status with BFRE training.

The development in maximum isometric knee-extension strength for individual patients are shown in Fig. 2. Results indicate that sitting test of maximum isometric knee-extension strength are feasible both at baseline and after 6–8 weeks following the fracture. Results show that muscle strength increase for both the injured and non-injured leg during the BFRE treatment. However, the difference after 6–8 weeks was median 18% decrease in muscle strength for the injured leg compared with the non-injured leg.

3.3. Adverse events

Seven of the eight patients reported delayed onset muscle soreness (DOMS). One patient reported pain from the cuff after one session. One patient reported temporary peripheral paresthesia to the front of the knee and the anterior portion of the crura after a single session. All AE disappeared within 24 h.

4. Discussion

This study indicates that early use of BFRE in patients with ankle fractures is feasible and have the potential to be used in future studies to evaluate the effect of BFRE on the development in rehabilitation following ankle fractures. Patients' overall satisfaction with the BFRE intervention were very high, and no serious adverse events were reported. Pilot study data in new treatment modalities is essential prior to large scale RCT trials evaluating efficacy, to guard against inability to recruit, feasibility of intervention, sample size calculation and SAE [21,22]. Data from the present study indicate that a large-scale, confirmatory trial is feasible.


4.1. The potential of enhanced and accelerated rehabilitation following ankle fractures

Ankle fractures are very common in the emergency departments world-wide [1]. During the past decades, the number of ankle fractures has increased steadily, which is thought to be caused by an increase in the number of people participating in sports and a shift in demographics towards an elderly population [2,23,24]. Following ankle fractures, patients commonly report a period of sick leave, pain, restrictions in range of joint motion, muscle weakness and difficulties with weight-bearing tasks such as sport, walking and climbing stairs [25–28]. Furthermore, long-term outcomes following ankle fractures are reported with increased risk of ankle pain and posttraumatic osteoarthritis [29]. Although optimal treatment, including surgery and rehabilitation are offered, the socioeconomic cost associated with

Table 2

Individual patient's acceptance of pain/discomfort during each BFRE training and patient's confident status.

Intervention		1	2	3	4	5	6
How acceptable was the level of pain/discomfort during the BFRE session?	Very much	7	7	7	7	7	7
	Much	1	1	1	1	1	1
	Neither/nor						
	Not at all						
To what degree did you experience security in today's BFRE session?	Very much	8	8	8	8	8	7
	Much						1
	Neither/nor						
	Not at all						

**Fig. 2.** Development in maximum isometric knee-extension strength. Legend: N = newton.

treatment of ankle fractures are considerable with an estimated cost between \$8688 and \$20,414 (2016 USD) per patient [30]. As a consequence, time effective, evidence based and safe rapid rehabilitation returning patients to previous function are of high interest for both patients and society. To achieve successful outcome following an ankle fracture restoration of muscle mass and muscle strength may be critical. BFRE training may represent the potential to facilitate a rapid rehabilitation following an ankle fracture.

4.2. Feasibility of BFRE in patients with ankle fractures

From the first introduction of BFRE to increase muscle strength and hypertrophy in healthy athletes [31,32] the indications have expanded to medical rehabilitation of different musculoskeletal diagnosis [33]. This study shows that early use of BFRE in patients with ankle fractures is feasible and associated with a high degree of patient satisfaction. Results from the present study are supported by Cancio et al. [34] reporting that BFRE are well-tolerated for patients with distal radius fractures. However, the current literature included limited information regarding the feasibility of BFRE in patients with fractures of weight bearing bones in the lower extremities [35].

4.3. Intervention

The use of tension band for exercise may limit the accuracy of the load for exercise. This may influence the predefined goal of 30% of 1RM in the present study and thereby the effect of the intervention. However, other studies have used elastic exercise band exercise together with BFRE for the lower extremities in older adults with excellent results [36]. The advantages for elastic tension band versus knee extension machine primary includes the ability for home exercise which may increase the ability for the use BFRE as an early intervention following ankle fractures. However, the present study did not include feasibility of home base BFRE.

4.4. Safety of BFRE in patients with ankle fractures

In general BFRE is considered safe for healthy active adults [11]. The safety of BFRE as intervention for patients with musculoskeletal diseases has been discussed in several papers. [16,33]. However, few studies associate BFRE with increased risk of serious cardiovascular adverse events, especially in patients with cardiovascular diseases [37,38]. Furthermore, rhabdomyolysis, numbness, bruising, subcutaneous haemorrhage, delayed onset muscle soreness, and pain are reported as potential side effects to BFRE [37,38]. This study did not observe any serious adverse events. One patient reported temporary peripheral paresthesia to the front of the knee and the anterior portion of the crura after a single session which disappeared within 24 h and almost all patients reported DOMS. To the authors knowledge, no studies have been reported on the safety of BFRE in patients with fractures of the lower extremities and therefore, little is known regarding safety. From the literature it is well known that surgery increases the risk of venous thromboembolism [39]. In the present study all patients with a previous history of cardiovascular diseases were excluded and the feasibility of BFRE may be interpreted with this in conclusion. Due to the lower number of participants even unexpected common SAE are not likely to have been captured in the present study population. As a consequence, data regarding safety should be interpreted with caution and are an important question in future research.

4.5. Limitations

Main limitations to this study is the selected group of patients included represented by almost young and healthy patients and a short follow-up period. Our group of patients do not represent the full spectrum of patient with ankle fractures which include a bimodal distribution with younger men and older women [1].

Another limitation is the use of use of tension band for exercise which may limit the accuracy of the load for exercise. Furthermore, the sample size are limited due to the pilot and feasibility study design, and no conclusion regarding potential effect of the intervention can be made.

5. Conclusion

Early use of BFRE in patients with unilateral ankle fractures seems feasible in patients without severe comorbidities. Patients' overall satisfaction with the BFRE intervention was very high. No serious adverse events were reported, although the study was not powered for these outcomes. BFRE training may represent the potential to facilitate an enhanced and accelerated rehabilitation following unilateral ankle fractures. Data from the present study will help inform a large-scale, confirmatory trial.

Conflict of interest

Peter Larsen, Oscar Just Platzer, Lærke Lollesgaard, Samuel Krogh Pedersen, Peter Kruse Nielsen, Michael S. Rathleff, Stefan Tegllhus Jensen and Rasmus Elsoe have no conflicts of interest to report.

Professor Thomas Bandholm report the following conflicts of interest:

Financial interests: I have received speaker's honoraria for talks or expert testimony on the efficacy of exercise therapy to enhance recovery after surgery at meetings or symposia held by biomedical companies (Zimmer Biomet and Novartis).

I have received fees for writing textbook chapters (Munksgaard) and for organising post-graduate education, such as post-graduate courses in clinical exercise physiology (Danish Physical Therapy Organization) or PhD courses on clinical research methodology (University of Copenhagen). Non-financial interests: I am an editorial board member with Br J Sports Med and PLOS ONE. I am a steering group member for "Clinical guidance catalogues" commissioned by the Danish Society of Sports Physical Therapy. I am a board member for the PhD-program: Basic and Clinical Research in Musculoskeletal Sciences (MUSKOS), University of Copenhagen. I am an exercise physiologist and physical therapist and may have a cognitive exercise bias.

The authors did not receive benefits or grants in any form from a commercial party related directly or indirectly to the subject of this article.

References

- [1] Elsoe R, Ostgaard SE, Larsen P. Population-based epidemiology of 9767 ankle fractures. *Foot Ankle Surg* 2018;24(February (1)):34–9, doi:<http://dx.doi.org/10.1016/j.fas.2016.11.002>.
- [2] Court-Brown CM. Rockwood and Green's fractures in adults, Vol. 8th. USA: Walters Kluwer; 2015.
- [3] Larsen P, Rathleff MSMS, Elsoe R. Surgical versus conservative treatment for ankle fractures in adults – a systematic review and meta-analysis of the benefits and harms. *Foot Ankle Surg* 2019;25(August (4)):409–17, doi:<http://dx.doi.org/10.1016/j.fas.2018.02.009>.
- [4] Appell HJ. Muscular atrophy following immobilisation. A review. *Sports Med* 1990;10:42–58.
- [5] Hortobágyi T, Dempsey L, Fraser D, Zheng D, Hamilton G, Lambert J, et al. Changes in muscle strength, muscle fibre size and myofibrillar gene expression after immobilization and retraining in humans. *J Physiol* 2000;524(Pt. 1):293–304, doi:<http://dx.doi.org/10.1111/j.1469-7793.2000.00293.x>.
- [6] Suetta C, Aagaard P, Rosted A, Jakobsen AK, Duus B, Kjaer M, et al. Training-induced changes in muscle CSA, muscle strength, EMG, and rate of force development in elderly subjects after long-term unilateral disuse. *J Appl Physiol* 2004;97:1954–61, doi:<http://dx.doi.org/10.1152/japplphysiol.01307.2003>.
- [7] Delorme TL, Watkins AL. Techniques of progressive resistance exercise. *Arch Phys Med Rehabil* 1948;29:263–73.
- [8] Husted RS, Troelsen A, Thorborg K, Rathleff MS, Husted H, Bandholm T. Efficacy of pre-operative quadriceps strength training on knee-extensor strength before and shortly following total knee arthroplasty: protocol for a randomized, dose-response trial (The QUADX-1 trial). *Trials* 2018;19:47, doi:<http://dx.doi.org/10.1186/s13063-017-2366-9>.
- [9] Loenneke JP, Abe T, Wilson JM, Thiebaud RS, Fahs CA, Rossow LM, et al. Blood flow restriction: an evidence based progressive model (Review). *Acta Physiol Hung* 2012;99:235–50, doi:<http://dx.doi.org/10.1556/APhysiol.99.2012.3.1>.
- [10] Clarkson MJ, May AK, Warmington SA. Chronic blood flow restriction exercise improves objective physical function: a systematic review. *Front Physiol* 2019;10:1058, doi:<http://dx.doi.org/10.3389/fphys.2019.01058>.
- [11] Patterson SD, Hughes L, Warmington S, Burr J, Scott BR, Owens J, et al. Blood flow restriction exercise position stand: considerations of methodology, application, and safety. *Front Physiol* 2019;10:533, doi:<http://dx.doi.org/10.3389/fphys.2019.00533>.
- [12] Nascimento D da C, Petriz B, Oliveira S da C, Vieira DCL, Funghetto SS, Silva AO, et al. Effects of blood flow restriction exercise on hemostasis: a systematic review of randomized and non-randomized trials. *Int J Gen Med* 2019;12:91–100, doi:<http://dx.doi.org/10.2147/IJGM.S194883>.
- [13] Bond CW, Hackney KJ, Brown SL, Noonan BC. Blood flow restriction resistance exercise as a rehabilitation modality following orthopaedic surgery: a review of venous thromboembolism risk. *J Orthop Sports Phys Ther* 2019;49:17–27, doi:<http://dx.doi.org/10.2519/jospt.2019.8375>.
- [14] Domingos E, Polito MD. Blood pressure response between resistance exercise with and without blood flow restriction: a systematic review and meta-analysis. *Life Sci* 2018;209:122–31, doi:<http://dx.doi.org/10.1016/j.lfs.2018.08.006>.
- [15] Lixandrão ME, Ugrinowitsch C, Berton R, Vechin FC, Conceição MS, Damas F, et al. Magnitude of muscle strength and mass adaptations between high-load resistance training versus low-load resistance training associated with blood-flow restriction: a systematic review and meta-analysis. *Sports Med* 2018;48:361–78, doi:<http://dx.doi.org/10.1007/s40279-017-0795-y>.
- [16] Hughes L, Paton B, Rosenblatt B, Gissane C, Patterson SD. Blood flow restriction training in clinical musculoskeletal rehabilitation: a systematic review and meta-analysis. *Br J Sports Med* 2017;51:1003–11, doi:<http://dx.doi.org/10.1136/bjsports-2016-097071>.
- [17] von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *Int J Surg* 2014;12:1495–9, doi:<http://dx.doi.org/10.1016/j.ijsu.2014.07.013>.
- [18] Slade SC, Dionne CE, Underwood M, Buchbinder R, Beck B, Bennell K, et al. Consensus on exercise reporting template (CERT): modified Delphi study. *Phys Ther* 2016;96:1514–24, doi:<http://dx.doi.org/10.2522/ptj.20150668>.
- [19] Rathleff CR, Baird WN, Olesen JL, Roos EM, Rasmussen S, Rathleff MS. Hip and knee strength is not affected in 12–16 year old adolescents with patellofemoral pain—a cross-sectional population-based study. *PLoS One* 2013;8:e79153, doi:<http://dx.doi.org/10.1371/journal.pone.0079153>.
- [20] Arain M, Campbell MJ, Cooper CL, Lancaster GA. What is a pilot or feasibility study? A review of current practice and editorial policy. *BMC Med Res Methodol* 2010;10:67, doi:<http://dx.doi.org/10.1186/1471-2288-10-67>.
- [21] Blümlé A, Schandelmaier S, Oeller P, Kasenda B, Briel M, von Elm E, et al. Premature discontinuation of prospective clinical studies approved by a research ethics committee – a comparison of randomised and non-randomised studies. *PLoS One* 2016;11:e0165605, doi:<http://dx.doi.org/10.1371/journal.pone.0165605>.
- [22] Leon AC, Davis LL, Kraemer HC. The role and interpretation of pilot studies in clinical research. *J Psychiatr Res* 2011;45:626–9, doi:<http://dx.doi.org/10.1016/j.jpsychires.2010.10.008>.
- [23] Bengtner U, Johnell O, Redlund-Johnell I. Epidemiology of ankle fracture 1950 and 1980. Increasing incidence in elderly women. *Acta Orthop Scand* 1986;57:35–7.
- [24] Court-Brown CM, McBirnie J, Wilson G. Adult ankle fractures—an increasing problem? *Acta Orthop Scand* 1998;69:43–7.
- [25] C-WCW Lin, Moseley AM, Herbert RD, Refshauge KM, C-W. CL, A.M. M, et al. Pain and dorsiflexion range of motion predict short-and medium-term activity limitation in people receiving physiotherapy intervention after ankle fracture: an observational study. *Aust J Physiother* 2009;55:31–7.
- [26] Stevens JE, Walter GA, Okereke E, Scarborough MT, Esterhai JL, George SZ, et al. Muscle adaptations with immobilization and rehabilitation after ankle fracture. *Med Sci Sports Exerc* 2004;36:1695–701.
- [27] Psatha M, Wu Z, Gammie FM, Ratkevicius A, Wackerhage H, Lee JH, et al. A longitudinal MRI study of muscle atrophy during lower leg immobilization following ankle fracture. *J Magn Reson Imaging* 2012;35:686–95, doi:<http://dx.doi.org/10.1002/jmri.22864>.
- [28] M.J. H, R.D. H. Prediction of outcome after ankle fracture. *J Orthop Sports Phys Ther* 2005;35:786–92.
- [29] Lübbecke A, Salvo D, Stern R, Hoffmeyer P, Holzer N, Assal M. Risk factors for post-traumatic osteoarthritis of the ankle: an eighteen year follow-up study. *Int Orthop* 2012;36:1403–10, doi:<http://dx.doi.org/10.1007/s00264-011-1472-7>.
- [30] Bielska IA, Wang X, Lee R, Johnson AP. The health economics of ankle and foot sprains and fractures: a systematic review of English-language published papers. Part 2: the direct and indirect costs of injury. *Foot (Edinb)* 2019;39:115–21, doi:<http://dx.doi.org/10.1016/j.foot.2017.07.003>.
- [31] Cook SB, Clark BC, Ploutz-Snyder LL. Effects of exercise load and blood-flow restriction on skeletal muscle function. *Med Sci Sports Exerc* 2007;39:1708–13, doi:<http://dx.doi.org/10.1249/mss.0b013e31812383d6>.
- [32] Scott BR, Loenneke JP, Slattery KM, Dascombe BJ. Blood flow restricted exercise for athletes: a review of available evidence. *J Sci Med Sport* 2016;19:360–7, doi:<http://dx.doi.org/10.1016/j.jsams.2015.04.014>.

- [33] Minniti MC, Statkevich AP, Kelly RL, Rigsby VP, Exline MM, Rhon DI, et al. The safety of blood flow restriction training as a therapeutic intervention for patients with musculoskeletal disorders: a systematic review. *Am J Sports Med* 2020;48:1773–85, doi:<http://dx.doi.org/10.1177/0363546519882652>.
- [34] Cancio JM, Sgromolo NM, Rhee PC. Blood flow restriction therapy after closed treatment of distal radius fractures. *J Wrist Surg* 2019;8:288–94, doi:<http://dx.doi.org/10.1055/s-0039-1685455>.
- [35] Hylden C, Burns T, Stinner D, Owens J. Blood flow restriction rehabilitation for extremity weakness: a case series. *J Spec Oper Med* 2015;15:50–6.
- [36] Yasuda T, Fukumura K, Tomaru T, Nakajima T. Thigh muscle size and vascular function after blood flow-restricted elastic band training in older women. *Oncotarget* 2016;7:33595–607, doi:<http://dx.doi.org/10.18632/oncotarget.9564>.
- [37] Nakajima T, Kurano M, Iida H, Takano H, Oonuma H, Morita T, et al. Use and safety of KAATSU training: results of a national survey. *Int J KAATSU Train Res* 2006;2:5–13, doi:<http://dx.doi.org/10.3806/ijktr.2.5>.
- [38] Patterson SD, Brandner CR. The role of blood flow restriction training for applied practitioners: a questionnaire-based survey. *J Sports Sci* 2018;36:123–30, doi:<http://dx.doi.org/10.1080/02640414.2017.1284341>.
- [39] Sweetland S, Green J, Liu B, Berrington de González A, Canonico M, Reeves G, et al. Duration and magnitude of the postoperative risk of venous thromboembolism in middle aged women: prospective cohort study. *BMJ* 2009;339:b4583, doi:<http://dx.doi.org/10.1136/bmj.b4583>.