

## Opening the black box of the relationship between neighborhood socioeconomic status and mental health

Neighborhood social-interactive characteristics as contextual mechanisms

Jakobsen, Andreas Lindegaard; Jørgensen, Anja; Tølbøll, Lene; Johnsen, Sisse Buch

*Published in:*  
Health & Place

*DOI (link to publication from Publisher):*  
[10.1016/j.healthplace.2022.102905](https://doi.org/10.1016/j.healthplace.2022.102905)

*Creative Commons License*  
CC BY 4.0

*Publication date:*  
2022

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

### *Citation for published version (APA):*

Jakobsen, A. L., Jørgensen, A., Tølbøll, L., & Johnsen, S. B. (2022). Opening the black box of the relationship between neighborhood socioeconomic status and mental health: Neighborhood social-interactive characteristics as contextual mechanisms . *Health & Place*, 77, Article 102905.  
<https://doi.org/10.1016/j.healthplace.2022.102905>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.





# Opening the black box of the relationship between neighborhood socioeconomic status and mental health: Neighborhood social-interactive characteristics as contextual mechanisms

Andreas Lindegaard Jakobsen<sup>a,\*</sup>, Anja Jørgensen<sup>a</sup>, Lene Tølbøll<sup>a</sup>, Sisse Buch Johnsen<sup>b</sup>

<sup>a</sup> Department of Sociology and Social Work, Aalborg University, Fibigerstræde 13, 9220, Aalborg, Denmark

<sup>b</sup> Department of Business Intelligence and Analysis, North Denmark Region, Niels Bohrs Vej 30, 9220, Aalborg, Denmark

## ARTICLE INFO

### Keywords:

Mental health  
Neighborhood socioeconomic status  
Trust  
Mediation  
Social capital

## ABSTRACT

Previous studies have linked low neighborhood socioeconomic status (NSES) to mental health problems. However, few studies have investigated the mechanisms underlying this association and most focused on the association with negative indicators of mental health, such as symptoms of depression or anxiety. This paper investigated whether neighborhood social characteristics (social interaction, trust, safety, organization participation, and attachment) mediate the association between NSES and mental health. We combined Danish register data with survey data from the North Denmark Region Health Survey 2017. Mental health was assessed with the Rand 12-item Short-form Survey (SF-12). The sample consisted of 14,969 individuals nested in 1047 neighborhoods created with an automated redistricting algorithm. We fitted multilevel structural equation mediation models and used a Monte Carlo simulation method to estimate confidence intervals for the indirect effects. NSES was positively associated with mental health. Neighborhood trust significantly mediated this relationship, accounting for 34% of the association after controlling for other mediators. These results indicate that higher levels of mental health in more affluent neighborhoods are partially explained by higher levels of trust. Improving neighborhood trust could mitigate sociogeographic inequalities in mental health.

## 1. Introduction

In recent decades, the number of studies linking neighborhood factors to health has grown exponentially, indicating a shift from focusing solely on individual health determinants to recognition of the importance of the surrounding neighborhood environment (Diez Roux, 2016). Although neighborhood factors have been suggested to affect the mental and physical health of individuals, only a few studies have focused on the challenge of opening the ‘black box’ of neighborhood effects (Galster, 2013; Jivraj et al., 2019; Rose et al., 2021; van Ham et al., 2012). Instead, previous studies have often identified correlations between individual outcomes and neighborhood characteristics without explicitly identifying specific causal mechanisms (van Ham et al., 2012).

One type of mechanisms that possibly explain the link between neighborhood context and mental health are social-interactive mechanisms (Galster, 2013), referring to social processes in neighborhoods such as social cohesion and collective efficacy (Sampson et al., 1997). Exposure to different types of social-interactive factors may ‘get under

the skin’ of residents and influence mental health in various ways. For example, neighborhood disorder including assault and trouble caused by young people or street gangs, may lead to chronic stress responses in residents that cause adverse physiological changes; in the long run, these changes may impair health (Van Deurzen et al., 2016). In contrast, social-interactive characteristics such as social cohesion might promote mental health due to greater positive affect and by buffering the effects of daily stressors (Robinetto et al., 2013) and negative life events such as economic crises (Loureiro et al., 2019).

Many previous studies have focused on structural factors such as neighborhood socioeconomic status (NSES) with several review studies concluding that socioeconomically deprived neighborhoods were associated with symptoms of depression (Julien et al., 2012; Mair et al., 2008; Richardson et al., 2015) as well as other various mental health and well-being outcomes (Silva et al., 2016; Visser et al., 2021). In addition to structural factors, a more holistic perspective is needed, to integrate multiple contextual factors, to capture in more detail different characteristics of various neighborhoods (Loureiro et al., 2019, 2022).

\* Corresponding author. Fibigerstræde 13, 9220, Aalborg, Denmark.

E-mail address: [andreaslj@socsci.aau.dk](mailto:andreaslj@socsci.aau.dk) (A.L. Jakobsen).

<https://doi.org/10.1016/j.healthplace.2022.102905>

Received 21 March 2022; Received in revised form 19 August 2022; Accepted 23 August 2022

Available online 9 September 2022

1353-8292/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Nevertheless, according to multiple reviews, the question of how NSES may affect the mental health status of residents through social-interactive mechanisms often remains unanswered (Aneshensel, 2010; Blair et al., 2014; Kim, 2008; Mair et al., 2008; Paczkowski and Galea, 2010; Visser et al., 2021). A few studies have examined this question with perceived neighborhood disorder (Greene et al., 2020; Kim, 2010; Ross, 2000; Schmidt et al., 2020), neighborhood social cohesion (Bassett and Moore, 2013; Erdem et al., 2015; Rios et al., 2012), network social capital (Haines et al., 2011), neighborhood social contacts (Drukker and van Os, 2003), worry about crime (Jonsson et al., 2020) and neighborhood violence (Joshi et al., 2017) as mediating factors. Most of these studies investigated how disadvantaged and socioeconomically deprived neighborhoods might influence symptoms of depression or anxiety. However, the use of such measures as indicators of overall population mental health has drawbacks, as mental health is more than just the absence of mental illness (Keyes, 2007). To our knowledge, only three of the abovementioned studies use mental health instruments that cover positive and negative as well as hedonic and eudemonic aspects of mental health (Drukker and van Os, 2003; Greene et al., 2020; Jonsson et al., 2020). However, it is important to study mental health from this perspective as well, as the mechanisms that influence ill-being may not be the same as the mechanisms of well-being (Huppert, 2009).

Another shortcoming associated with most studies in this area is the use of predefined administrative/census-based areas to delineate neighborhoods (Mair et al., 2008; Richardson et al., 2015; Truong and Ma, 2006; Visser et al., 2021). Such areas rarely correspond to theoretical concepts or residents' understandings of neighborhoods; thus, the use of such areas is potentially problematic for the study of social processes (Sampson et al., 2002). Instead, the areas used should correspond to the hypothesized mechanisms under study (Visser et al., 2021). This may lead to different results when the use of administrative areas is compared to alternative theoretically based neighborhoods due to the so-called modifiable areal unit problem (Chaix et al., 2006; Flowerdew et al., 2008; Franzini and Spears, 2003; Jakobsen, 2021; Messer et al., 2012; Parenteau and Sawada, 2011).

In an effort to further this area of research, the aim of this study was to analyze various possible contextual social-interactive mechanisms explaining the association between NSES and mental health. Instead of relying on administrative units as neighborhoods, we used neighborhoods created with an automated redistricting algorithm with the use of physical boundaries such as larger roads as neighborhood dividers. The use of physical barriers as dividers stems from findings that such barriers can also function as social dividers and thereby promote or hinder social interaction (Feld, 1981; Grannis, 1998) and result in high within-group sociodemographic homogeneity (Foster and Aaron Hipp, 2011; Lund, 2018). Furthermore, residents may use such major physical barriers to help identify their own neighborhood from distinct areas (Campbell et al., 2009; Grannis, 2009).

Thus, we investigated the indirect effects of multiple contextual mechanisms related to the social-interactive environments of neighborhoods. We hypothesized that (i) higher levels of NSES are positively associated with mental health and (ii) that one or more neighborhood social-interactive characteristics, in the form of neighborhood social interaction, trust, organization participation, attachment, and safety, mediate the relationship between NSES and mental health.

## 2. Theory of neighborhood social-interactive mechanisms

Social-interactive mechanisms include complex multifaceted concepts, such as social capital (e.g., resources from social networks including trust and norms of reciprocity) (Putnam, 2000), social cohesion (e.g., strength of social relationships, sense of belonging, shared values, common identity, trust, and the existence of equal opportunities versus social exclusion within a community) (Berger-Schmitt, 2002), and the related concept of collective efficacy (social cohesion, trust and

informal social control) (Sampson et al., 1997). Despite the differences between these concepts, they share key similarities. In addition, social capital can be separated into structural and cognitive components characterized as what people 'do' and what people 'feel', respectively (Harpham et al., 2002). We focused on social interactions as well as neighborhood organization participation (NOP) as examples of specific structural dimensions, and trust, attachment, and safety as examples of cognitive dimensions.

Social ties and interaction have been well established as generally beneficial for mental health (Kawachi and Berkman, 2001; Santini et al., 2015). In addition, living in disadvantaged neighborhoods has been linked to smaller social networks (York Cornwell and Behler, 2015) and loneliness (Algren et al., 2020). However, structural characteristics of a neighborhood may not always be a strong determinant of social ties and interactions within the neighborhood itself (Berg and Timmermans, 2015; Jørgensen, 2010).

Another structural component of social capital is the level of civic engagement, in the form of NOP, which can help build relationships of trust and reciprocity with others (Collins et al., 2014). Participation in organizations has been linked to higher quality of life and fewer depressive symptoms (Anderson et al., 2014; Santini et al., 2020; Takagi et al., 2013); furthermore, residents in less affluent areas are more likely to have low levels of social participation (Bowling and Stafford, 2007).

When focusing on the quality of the social interactions taking place in neighborhoods, one of the key cognitive aspects is the particularized interpersonal trust between people in the neighborhood (Garoon et al., 2016). Drawing on the work by Simmel and Luhmann, Lewis and Weigert define trust as "a functional alternative to rational prediction for the reduction of complexity" that "allows social interactions to proceed on a simple and confident basis" (Lewis and Weigert, 1985). Previous studies have linked lower levels of trust to neighborhood deprivation (Corcoran et al., 2018; Franzini et al., 2005; Nettle et al., 2014) depression (Fahmi et al., 2019; Fujiwara and Kawachi, 2008) and well-being (Helliwell and Putnam, 2004).

Place attachment, defined as a positive affective bond or association between individuals and their residential environment (Shumaker and Taylor, 1983), has also been linked to better mental health (Scannell and Gifford, 2017; Young et al., 2004). Attachment to places can be considered a fundamental human need based on the emotional significance that geographic spaces evoke in humans, thereby transforming into meaningful "places" (Giuliani, 2003). Moreover, place attachment can occur at both the individual and group levels and thereby be shared by group members such as neighbors (Scannell and Gifford, 2010). Feelings of place attachment may be lower in deprived neighborhoods due to circumstances such as higher levels of crime and lower safety (Bailey et al., 2012).

From a biological perspective, the feeling of being in a safe environment is linked to an optimal bodily state and social engagement behaviors, while feelings of unsafety are linked to defensive stress responses (Porges, 2007). Consistent with these findings, living in less safe neighborhoods has been associated with psychological distress and depression (Booth et al., 2012; Choi and Matz-Costa, 2018; Cromley et al., 2012; Wilson-Genderson and Pruchno, 2013). Moreover, deprived neighborhoods have been linked to lower perceived safety (Franzini et al., 2005; Mouratidis, 2020) and higher levels of crime (Benson et al., 2003; Sampson et al., 1997; Zimmerman and Messner, 2010).

To summarize, previous research has shown that these abovementioned neighborhood characteristics are related to various aspects of mental health but are unequally distributed across neighborhoods with different NSESs, which makes these characteristics theoretically plausible mediators in the pathway between NSES and mental health.

### 3. Methods

#### 3.1. Study area

The study area includes the North Denmark Region which is one of five Regions in Denmark. These regions are administrative entities above the municipality level, and below the central government. The North Denmark Region covers an area of 7,933 km<sup>2</sup> with a population of 587,335 and a population density (inhabitants per km<sup>2</sup>) of 74.5 in 2017 (Statistics Denmark, 2017).

#### 3.2. Data sources

This study used three different types of data: 1) survey data from the North Denmark Region Health Survey 2017 to measure mental health and neighborhood contextual mechanisms; 2) register data for the population from age 16 living in the North Denmark Region in 2017 derived from various registers by Statistics Denmark (Baadsgaard and Quitzau, 2011; Jensen and Rasmussen, 2011; Pedersen, 2011) to measure individual-level sociodemographic characteristics and NSES; and 3) georeferenced micro-areas developed by Lund (2018) to divide individuals into different neighborhoods.

The North Denmark Region Health Survey is a population-based survey designed to be representative of the adult population aged 16 and above in the North Denmark Region. The questionnaire includes a broad range of questions related to morbidity and health, focusing on aspects not available in registers, e.g., questions about neighborhoods. Individuals were randomly sampled from the Danish Civil Registration System and invited by postal or electronic mail. The survey was conducted by the North Denmark Region from February to May 2017, and reminders were sent to nonrespondents. A total of 22,583 individuals completed the questionnaire (60.1%), providing an overall margin of error of <1% at a 95% confidence level. The sample was restricted to respondents from neighborhoods with a minimum of five observations due to discretionary criteria and with no missing data on all variables used in the analyses, resulting in a final sample of  $N = 14,969$  individuals nested in 1047 neighborhoods.

Missing data were omitted from the analysis as the final sample showed no major differences from the full sample on key demographic variables (gender, age and marital status) as well as on mental health status and NSES.

#### 3.3. Neighborhoods

Neighborhoods were measured using micro-areas created by Lund (2018), which have been used in previous studies to investigate neighborhood characteristics in a Danish context (Jakobsen, 2021; Kristiansen and Lund, 2022; Lund, 2018, 2020; Lund et al., 2019). The areas were constructed with an automated redistricting algorithm that formed the smallest areas possible that contained at least 100 inhabitants and were separated by large physical barriers. To run the algorithm, the National Square Grid that assigned addresses in Denmark to 'hectare cells' (100 m x 100 m) in 2000, 2005, 2010, and 2015 was used as georeferenced data. The algorithm works in two overall steps. First, initial areas were delineated with the use of physical barriers such as highways, roads broader than 6 m, rivers and streams broader than 3 m, railways, lakes, forests, coastlines, and intakes. In this step, the square grid was applied with the grids dissolved into areas where the largest part of the square was located with borders formed after the grid was applied. Thus, the straight borders were replaced with the borders of the squares in each area. After this step, the areas were further clustered to ensure that there were at least 100 inhabitants in each area according to discretionary criteria of Statistics Denmark. Further criteria were set to ensure the least possible number of merges and the smallest possible number of inhabitants. These criteria ensure that the algorithm consistently generates the same areas if the process is repeated (Lund, 2018). For a more

detailed description of the micro-areas, see Lund (2019, 2018).

#### 3.4. Mental health

Mental health was measured using the mental component score (MCS-12) of the Rand 12-item Short Form questionnaire (SF-12) (Ware John et al., 1996). The SF-12 is a multipurpose generic assessment of health status that covers eight health concepts: physical function, limitations due to physical health and emotional problems, bodily pain, general health, vitality, social function and psychological distress and well-being over the last 4 weeks. From these items, a physical component score (PCS-12) and a mental component score (MCS-12) can be constructed (Ware et al., 1998). All 12 items are used to construct the MCS-12, but the scale places added emphasis on items covering the concepts of emotional problems, vitality, social function and psychological distress and well-being. The MCS-12 is considered a valid measure of mental health in health research (Cheak-Zamora et al., 2009) and has previously been used to measure mental health in various countries, including Denmark (Christensen et al., 2014, 2020; Fong et al., 2010; Kontodimopoulos et al., 2007; Peterson et al., 2009).

In this study, the 12 items were summed and converted into the MCS-12. After the subscale scores were calculated, they were weighted according to population norms. Weights from the North Denmark Region population were calculated and used as population norms instead of the originally proposed population norms from the scoring algorithms (Bjørner et al., 1997). Using this weighting emphasizes population and time adequacy over the possibility of comparing scale scorings to other studies, which corresponds to the aim of the present study (Tucker et al., 2016). Furthermore, we used an exploratory factor analysis with oblique rotation instead of a principal component analysis with orthogonal rotation, since the latter can cause inconsistencies between the subscales and summarized scales and assumes zero correlation between the mental and physical health factors (Blanchard et al., 2004; Fleishman et al., 2010; Jakobsson et al., 2012; Tucker et al., 2016).

#### 3.5. Neighborhood socioeconomic status

A composite index to measure NSES was created based on the full Danish population above age 16 in 2017 (Jakobsen, 2021). Based on previous contextual socioeconomic deprivation scores, the index was created with the following three indicators: the proportion of the population between 30 and 64 years of age that was unemployed for at least half of the year, including recipients of sickness benefits, persons on leave and recipients of cash benefits (Bender et al., 2015; Juhász et al., 2010; Meijer et al., 2013); proportion of the population between 30 and 64 years of age with a disposable income in the lowest quartile (Bender et al., 2015; Meijer et al., 2013); and proportion of the population between 30 and 64 years of age with basic education (UNESCO Institute for Statistics, 2012) as the highest attained educational level (Bender et al., 2015; Juhász et al., 2010; Lund, 2020).

All three indicators were standardized to z-scores and used to create an index, with principal component analysis (PCA). The index was reversed and standardized to a z-score, with higher scores indicating higher NSES.

#### 3.6. Neighborhood social characteristics

*Neighborhood social interaction* was measured with the question "How often are you in contact with friends, acquaintances and family that you do not live with (specifically neighbors or residents in your local area)?" with the response categories (Daily or almost daily, 1 or 2 times a week, 1 or 2 times a month, Less than once a month, or Never) coded from 5 to 1. *Neighborhood trust* was measured with the question "To what extent do you trust people from your settlement/neighborhood/local area?" with response options (Trust them completely, Trust them a lot, Do not trust them very much, or Do not trust them at all). *NOP* was measured with



the question “Do you participate in associations or volunteer work in one or more of the following areas?” and providing a wide range of options ranging from housing and community organizations to youth work such as scouts or youth clubs with response options (Yes, daily; Yes, weekly, but not every day; Yes, monthly, but not every week; Yes, but rarely participates/is a member; or No, never participates/is not a member). This measure was summed into a composite index, with higher scores indicating a higher degree of participation in various organizations. *Neighborhood attachment* was measured with the question “Do you feel attached to your local area?” with the response categories (Yes, strongly attached; Yes, partly attached; No, not particularly attached; or No, not at all attached). *Neighborhood safety* was measured with the question “How safe do you feel walking alone in your local area after dark?” with response options (Very safe, Fairly safe, Fairly unsafe, or Very unsafe).

To capture the mechanisms at a collective neighborhood level and further reduce same-source bias, the individual scores were used to construct ecological average scores at the neighborhood level. Due to variation in observations across neighborhoods, which ranged from 5 to 217, we used empirical Bayes estimates (EBEs). These estimates borrow strength across neighborhoods and shrink estimates for neighborhoods with few observations toward the overall mean (Mujahid et al., 2007) with a shrinkage factor calculated as follows:

$$\text{Shrinkage factor} = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2/n_j}$$

where  $\sigma_u^2$  is the level-2 variance and  $\sigma_e^2$  is the level-1 variance, and  $n_j$  is the number of observations in neighborhood  $j$  in the sample (Hox et al., 2018). The less precise the group-specific estimate and the less variability observed across groups, the greater the shift toward the overall group mean. The strength of this approach is that it uses information from other neighborhoods to improve estimates for unreliable neighborhoods (Diez Roux, 2002; Mujahid et al., 2008). Finally, for interpretational purposes, the scores were standardized to z-scores, with higher scores indicating a higher degree of neighborhood social interaction, trust, organization participation, attachment, and safety.

### 3.7. Individual-level variables

Gender, age, marital status, education, personal income, employment status and ethnicity were included as individual-level variables. Age was grand-mean centered. Marital status was measured as married, cohabiting or living alone. Education was measured as the highest attained education according to the ISCED collapsed into three categories: ‘Basic education’ (levels 0–2), ‘Medium education’ (levels 3–5) and ‘High education’ (levels 6–8) (UNESCO Institute for Statistics, 2012). Income was measured as the total annual personal income (except for any rental income from one’s own accommodation and before deducting labor-market contributions and pension contributions) and categorized into quartiles. Employment status was collapsed into four categories: employed, unemployed, student and pensioner. Ethnicity was dichotomized to Danish or other.

### 3.8. Statistical analysis

To open the black box of the mechanisms by which NSES influences mental health, we used a structural equation multilevel mediation (MSEM) approach with a 2-2-1 design (as illustrated in Fig. 1), given that individuals were nested in neighborhoods; NSES and neighborhood social characteristics were measured at level 2 and individual mental health was measured at level 1 (Preacher et al., 2010; Zhang et al., 2009).

The basic idea of simple mediation is when the relationship between two variables is fully or partially accounted for by a third variable that conceptually lies on the causal pathway between the exposure and the outcome (Baron and Kenny, 1986). In mediation terminology, the




Fig. 1. Illustration of a simple 2-2-1 multilevel mediation model.

relations between the variables can be divided into distinct paths: the path between the predictor and mediator (path  $a$ ), the path between the mediator and outcome (path  $b$ ) and the path between the predictor and the outcome, also called the total effect (path  $c$ ). Finally, the path between the predictor and outcome, once intervening mediated relations have been accounted for is  $c-ab$  (path  $c'$ ), also called the direct path. The indirect effect of the predictor through the mediator that measures the amount of mediation can then be quantified as the product of  $a$  and  $b$  (i.e.,  $ab$ ). (Preacher and Hayes, 2008).

Due to the skewness and nonnormality of the sampling distribution of the indirect effect, common methods for estimating the confidence interval of the indirect effect, such as the *delta method* (Sobel, 1982), can result in a conservative, underpowered test that fails to detect a true indirect effect (MacKinnon et al., 2004; Preacher and Selig, 2012). To solve this problem, we used the Monte Carlo simulation method to construct confidence intervals for the indirect effects (MacKinnon et al., 2004). The Monte Carlo simulation method has been shown to produce results comparable to those of bootstrap methods; furthermore, this method is much faster and less computationally demanding to implement in a multilevel context (Preacher and Selig, 2012). To construct the confidence interval for the indirect effect, we first estimated the effects of  $\hat{a}$  and  $\hat{b}$  and their standard errors of  $\hat{\sigma}_{\hat{a}}$  and of  $\hat{\sigma}_{\hat{b}}$  from the sample. Next, we generated a simulated sampling distribution of the indirect effect  $ab$  by generating a distribution of 100,000 random samples with population values equal to the sample values of  $\hat{a}$ ,  $\hat{b}$ ,  $\hat{\sigma}_{\hat{a}}$  and  $\hat{\sigma}_{\hat{b}}$ . Finally, percentiles of the sampling distribution were used as limits for a 95% confidence interval of the sample  $\hat{a}\hat{b}$ .

To estimate the overall variance in mental health between neighborhoods, also known as the general contextual effect (GCE) (Merlo et al., 2018) we first modeled an “empty” multilevel model that only included a random intercept with no predictors. From this model the intraclass correlation coefficient (ICC) was calculated as follows:

$$\text{ICC} = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2}$$

where  $\sigma_u^2$  is the level-2 variance and  $\sigma_e^2$  is the level-1 variance. The ICC is the percentage of the total variance attributable to the area-level variance (Snijders and Bosker, 1999). Second, we estimated the crude associations between mental health and NSES, social interaction, trust, NOP, attachment and safety as contextual mechanisms, with each predictor in a separate model. Third, we included the individual-level covariates to estimate the adjusted associations between mental health and each predictor separately, with the estimate of NSES corresponding to the total effect of NSES. Finally, we estimated a model that included all predictors concurrently. Indirect effects were estimated with the use of both single and multiple mediation models. The single mediation models allowed us to investigate if the various potential mechanisms acted as mediators, and the multiple mediation model allowed us to study the possible mediating effect of the potential mechanisms when controlling for all other mediators and thereby the mechanism’s unique

ability to mediate the relationship between NSES and mental health (Preacher and Hayes, 2008). Based on the recommendations of Preacher and Hayes (2008), we allowed the residuals associated with the mediators to covary. To quantify the strength of mediation, we calculated the ratio of the indirect effect to the total effect as  $\hat{P}_M = (\hat{a} \times \hat{b}) / \hat{c}$  (Shrout and Bolger, 2002).

Before conducting the multilevel models, we estimated variance inflation factors (VIFs) to identify possible multicollinearity due to positive zero-order correlations between the neighborhood characteristics. The VIFs for the neighborhood characteristics and mean VIF were below 2.5, indicating no serious problems with multicollinearity.

To account for potential differences in selection probabilities and response rates, we used calibrated weights constructed by Statistics Denmark for the analyses.

#### 4. Results

Fig. 2 shows the micro-areas mapped with the EBES for mean mental health scores divided into four classes using Jenks Natural Breaks (Jenks, 1967) with darker colors indicating higher levels of mental health, and with no clear geographic or urban-rural pattern identified. The zero-order correlations at the ecological level with NSES and EBES for mental health and neighborhood characteristics are presented in Table 1. Results indicated that neighborhoods with higher levels of NSES, social interaction, trust, NOP, attachment, and safety were associated with higher ecological levels of mental health; NSES had the strongest correlation with mental health ( $r = 0.36$ ), followed by trust ( $r = 0.35$ ) and safety ( $r = 0.34$ ). Furthermore, neighborhoods with higher levels of social interaction, trust, NOP, attachment, and safety were associated with higher levels of NSES; trust ( $r = 0.32$ ) and safety ( $r = 0.35$ ) had the strongest associations with NSES. Significant correlations were observed between all neighborhood social characteristics. Neighborhoods with higher levels of attachment were particularly strongly associated with higher levels of social interaction ( $r = 0.58$ ) and trust ( $r = 0.49$ ). In addition, neighborhoods with higher levels of trust were moderately associated with higher levels of social interaction ( $r = 0.48$ ). For descriptive statistics of all study variables, see the Supplementary Material.

The ICC for mental health was 2.5% (SE = 0.004). This is comparable

to other similar studies using the SF-12 or SF-36 (Fone and Dunstan, 2006; Peterson et al., 2009) and shows a relatively small GCE of neighborhoods.

Table 2 shows the model results for the association between the neighborhood characteristics and individual mental health. For the crude estimates, the models showed significantly higher mental health of individuals in neighborhoods with higher levels of all neighborhood characteristics, with NSES being the strongest predictor of mental health. After adjusting for individual characteristics, the associations were largely attenuated, but all neighborhood characteristics were still significant predictors of mental health. In the fully adjusted models (adjusted for both individual-level characteristics and all neighborhood characteristics), only neighborhood trust remained a significant predictor of mental health (Coef: 0.25, 95% CI: 0.05, 0.45).

Before conducting the mediation analysis, we tested for exposure-mediator interactions for all neighborhood characteristics (Valeri and VanderWeele, 2013). We identified a significant interaction between NSES and NOP (see Supplementary Material), showing that the relationship between NSES and mental health was less pronounced in neighborhoods with high NOP than in neighborhoods with low NOP. In the full model that controlled for all other neighborhood characteristics, the interaction was no longer significant and did not improve model fit; therefore, it was not included in the mediation models. We further tested for interactions between all neighborhood mediators but found no significant interactions.

Table 3 shows the indirect effects of the neighborhood characteristics from the separate single mediation models and the indirect effects from a multiple mediation model with all neighborhood mediators entered concurrently. In the single mediation models, social interaction, trust, attachment, and safety were significant mediators. Neighborhood trust showed the largest mediation effect, accounting for 43% of the association between NSES and mental health, followed by neighborhood safety, which accounted for 31%. In the multiple mediation model, only neighborhood trust was a significant mediator of this relationship (Coef: 0.13, 95% CI: 0.03, 0.24) accounting for 34% of the association between NSES and mental health when controlling for the other neighborhood mediators.




Fig. 2. Map of North Jutland, Denmark mapped with neighborhood EBES of mean mental health scores.

**Table 1**

Ecological correlation matrix with EBEs for mental health, neighborhood characteristics and NSES.

	1.	2.	3.	4.	5.	6.	7.
1. Mental health	1.00						
2. NSES	0.36***	1.00					
3. Social interaction	0.22***	0.13***	1.00				
4. Trust	0.35***	0.32***	0.48***	1.00			
5. NOP	0.24***	0.16***	0.28***	0.22***	1.00		
6. Attachment	0.22***	0.16***	0.58***	0.49***	0.24***	1.00	
7. Safety	0.34***	0.35***	0.24***	0.36***	0.13***	0.29***	1.00

(N = 1047 neighborhoods) \*\*\*p &lt; 0.001, \*\*p &lt; 0.01, \*p &lt; 0.05.

**Table 2**

Model results predicting mental health.

	Crude estimates <sup>a</sup>		Adjusted estimates <sup>b</sup>		Full model estimates <sup>c</sup>	
	Coef	(95% CI)	Coef	(95% CI)	Coef	(95% CI)
NSES	0.88***	(0.72, 1.05)	0.37***	(0.22, 0.52)	0.17	(-0.01, 0.35)
Social interaction	0.46***	(0.28, 0.63)	0.26***	(0.11, 0.42)	0.01	(-0.20, 0.22)
Trust	0.77***	(0.61, 0.93)	0.41***	(0.27, 0.55)	0.25*	(0.05, 0.45)
NOP	0.44***	(0.24, 0.64)	0.24**	(0.09, 0.39)	0.09	(-0.06, 0.24)
Attachment	0.50***	(0.31, 0.68)	0.29***	(0.13, 0.45)	0.01	(-0.20, 0.21)
Safety	0.76***	(0.58, 0.93)	0.36***	(0.21, 0.51)	0.14	(-0.04, 0.31)

Coef = path coefficient. <sup>a</sup> Crude estimates from separate models. <sup>b</sup> Adjusted estimates from separate models. <sup>c</sup> Adjusted estimates from the full model containing all predictors concurrently. Adjusted = adjusted for individual-level gender, age, marital status, education, personal income, employment status and ethnicity. (N = 14,969) \*\*\*p < 0.001, \*\*p < 0.01, \*p < 0.05.

**Table 3**

Monte Carlo indirect effect estimates.

	Single mediation <sup>a</sup>			Multiple mediation <sup>b</sup>		
	Coef	(95% CI)	$\hat{P}_M$	Coef	(95% CI)	$\hat{P}_M$
Social interaction	0.05*	(0.01, 0.09)	0.13	0.01	(-0.05, 0.06)	–
Trust	0.17***	(0.09, 0.25)	0.43	0.13*	(0.03, 0.24)	0.34
Attachment	0.06*	(0.01, 0.12)	0.16	0.00	(-0.06, 0.07)	–
Safety	0.12**	(0.04, 0.20)	0.31	0.07	(-0.02, 0.16)	–

Coef = path coefficient. <sup>a</sup> Estimates from separate single mediation models. <sup>b</sup> Estimates from full multiple mediation model.

$\hat{P}_M$  = indirect effect/total effect. All models adjusted for individual-level gender, age, marital status, education, personal income, employment status and ethnicity. (N = 14,969) \*\*\*p < 0.001, \*\*p < 0.01, \*p < 0.05.

## 5. Discussion

In this study, we aimed to open the “black box” of neighborhood effects concerning the relationship between NSES and mental health by empirically testing various potential contextual mechanisms related to the social-interactive environment of neighborhoods. We found that higher levels of NSES, social interaction, trust, NOP, attachment, and safety were significantly associated with higher levels of mental health, even when controlling for individual-level factors. Furthermore, we found an interaction between NSES and NOP, indicating that a high degree of participation in organizations in neighborhoods may buffer the impact of NSES on mental health. However, this interaction was no longer significant after controlling for other neighborhood characteristics. In the single mediation models, neighborhood social interaction, trust, attachment, and safety significantly mediated the relationship between NSES and mental health. In a multiple mediation model, only neighborhood trust was a significant mediator of the relationship between NSES and mental health and accounted for 34% of the association. These results indicate that higher levels of mental health associated with living in more affluent neighborhoods is partially explained by higher levels of mutual trust among neighbors when taking other social-interactive neighborhood characteristics into account.

Our finding that neighborhood social characteristics can mediate the

association between NSES and mental health is in line with previous studies (Bassett and Moore, 2013; Drukker and van Os, 2003; Erdem et al., 2015; Haines et al., 2011; Rios et al., 2012). Rios et al. (2012) reported that neighborhood social cohesion accounted for 10% of the association between NSES and individual psychological distress. In that study, neighborhood social cohesion was measured using the social cohesion dimension of the Collective Efficacy Scale (Sampson et al., 1997), in which mutual trust was one of the core indicators (Rios et al., 2012). In addition, other studies show that neighborhood social cohesion mediates the association between NSES and depressive symptoms (Bassett and Moore, 2013) as well as psychological distress (Erdem et al., 2015). In the present study, we did not investigate composite measures of neighborhood social cohesion, but our finding that neighborhood trust was the strongest and only significant mediator and predictor of mental health in the full models is not surprising, since trust is normally understood to be a key indicator of social cohesion and social capital and is sometimes used as a single indicator (Delhey and Newton, 2003). A study from the UK found generalized trust to be the only social capital indicator showing a positive and significant association with psychological health overtime, where social participation in local groups and the frequency of talking to neighbors showed no such effect (Giordano and Lindström, 2011). A review by Ehsan and De Silva (2015) concluded that both individual and ecological measures of cognitive social capital,



e.g., mutual trust, were associated with a reduced risk of common mental disorders, while there was no overall association with structural social capital at either the individual or ecological level. As one possible explanation for the importance of trust, a trusting neighborhood environment might be characterized by positive and supportive personal relationships beneficial for mental health, while this may not necessarily be the case in neighborhoods with high levels of attachment, safety, or social interaction. In addition, attachment, safety, and social interaction might lead to higher levels of trust, with trust acting as a mediator. However, further studies are needed to clarify how these factors may be causally related.

Even though social capital has been positively linked to mental health, there is a lack of controlled trials determining a causal effect (Flores et al., 2018). Nevertheless, a few intervention studies have shown promising results. For example, a quasi-experimental study that investigated an ecological intervention with the objectives of promoting community participation in urban renewal and engaging residents in the construction of attractive urban places showed improvements in mental health, sense of community and social capital (Semenza et al., 2007).

In our study, neighborhood trust was measured at the ecological level. However, there is disagreement on whether neighborhood social characteristics should be measured individually or collectively (Ehsan et al., 2019; Sampson, 2011; Sampson et al., 2002). Both Sampson (2011) and Kawachi and Berkman (2015) argue that collective efficacy and social capital should be seen as social and collective phenomena; therefore, they simply cannot be captured through individual responses alone. We believe that this is also the case with trust: trusting social environments should be seen as the result of mutual trust among residents and not just as a single resident's degree of trust in his neighbors. When neighborhood trust was measured at the individual level, sensitivity analyses showed that it was a stronger predictor of individual mental health than the ecological measure but with a weaker mediating effect, accounting for 27% of the association between NSES and mental health. This indicates that ecological measures of neighborhood social characteristics, in addition to individual responses, are important to consider when investigating the social mechanisms of place. One explanation of the biosocial link between NSES and mental health was proposed by Prior et al. (2018), who found that allostatic load, a biological response to chronic stress, acted as a biological mediator between neighborhood deprivation and both physical and mental health. The complete pathway between NSES and mental health is likely complex and potentially includes individual, social, structural, cognitive, and biological factors. Future studies should investigate more complex pathways and different types of mediators to provide important insights that allow for a better understanding of the link between neighborhoods and mental health.

### 5.1. Study strengths and limitations

One key strength of this study lies in our ability to use micro-areas designed by an automated redistricting algorithm that accounts for physical barriers in the landscape to function as social dividers (Lund, 2018), instead of relying on administrative delineations as neighborhood units. Sensitivity analyses showed that when using parishes, the smallest administrative area in Denmark, we found no mediation effect of any of the neighborhood characteristics. Furthermore, the ICC for mental health was lower when using parishes ( $ICC = 1.5\%$ ,  $SE = 0.003$ ). Together with previous research, this indicates that micro-areas are better at capturing relevant neighborhood effects when focusing on mental health-related outcomes (Jakobsen, 2021). In addition, we included a variable at the municipality level with four categories — 1) urban, 2) semi urban, 3) rural and 4) outer municipality (The Danish Ministry of Food, Agriculture and Fisheries and The Danish Ministry of the Interior and Health, 2011) - to account for the potential contextual confounder of living in an urban or rural area. However, this did not change the results substantially.

As previously mentioned, physical barriers can serve as barriers to social interaction (Feld, 1981; Grannis, 1998) and may reflect individuals' own perceptions of neighborhoods (Campbell et al., 2009; Grannis, 2009; Lynch, 1971). If neighborhood trust matters for mental health, it is clear why the use of larger administrative boundaries may lead to different results, as the average level of trust in these areas can be only marginally expected to reflect the socially interactive environment that individuals experience and participate in on a daily basis. Physical barriers may however not always function as barriers between neighborhoods. It is possible that people who live close to various attractive blue and green space barriers may form social clusters separated from those who live further away from these barriers (Schüle et al., 2019).

One major limitation of this study was the cross-sectional design, which prevents a determination of temporal order in the association between NSES and mental health. However, previous studies using experimental and cohort designs have found evidence for a temporal causal link between NSES and subsequent mental health conditions (Crump et al., 2011; Galea et al., 2007; Leventhal and Brooks-Gunn, 2003; Ludwig et al., 2013; White et al., 2017).

Regarding the association between NSES and the self-reported neighborhood characteristics, we consider reverse causation more unlikely, as this would mean that NSES is affected by characteristics such as perceived safety and trust. However, the path between the different neighborhood characteristics and mental health is more likely to be reversed, as it is highly possible that a person's mental health may affect their perception of trust, etc. in a neighborhood (Echeverria et al., 2004). Thus, one strength of this study was the use of ecological measures instead of individual responses as measurements of neighborhood characteristics; this method also reduced same-source bias (Mujahid et al., 2007). In addition, we included several individual characteristics to control for various potential exposure-outcome, mediator-outcome and exposure-mediator confounders (VanderWeele, 2016). Another limitation of this study concerns our inability to include both individual and micro-area-level weights, as the available weights designed for this study by Statistics Denmark were combined weights of the individual and municipality levels, which may have biased our results. However, sensitivity analyses showed that when running the mediation models without weights, neighborhood trust was still the only significant mediator in the multiple mediation model.

### 5.2. Conclusion

This study adds to the existing literature as we used a new method of automatically generated neighborhoods, which accounts for physical barriers in the landscape, to study whether social characteristics of neighborhoods (social interaction, trust, safety, NOP, and attachment) mediate the association between NSES and mental health. We found evidence that neighborhood trust is a mediator of the association between NSES and mental health. Thus, the higher mental health levels of residents in more affluent neighborhoods can be partially explained by higher levels of neighborhood trust. Improvements in trust between neighborhood residents could potentially mitigate geographic and socioeconomic inequalities in mental health. However, further studies are needed to test the effect of specific public health interventions targeted at the neighborhood level on mental health.

### Data availability

The authors do not have permission to share data.

### Acknowledgments

We thank members of the SocMap research group at the Department of Sociology and Social Work, Aalborg University, and Pernille Tanggaard Andersen, University of Southern Denmark for providing useful comments on an earlier draft. The North Denmark Region Health Survey

was conducted and funded by The North Denmark Region.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.healthplace.2022.102905>.

## References

- Algren, M.H., Ekholm, O., Nielsen, L., Ersbøll, A.K., Bak, C.K., Andersen, P.T., 2020. Social isolation, loneliness, socioeconomic status, and health-risk behaviour in deprived neighbourhoods in Denmark: a cross-sectional study. *SSM - Popul. Heal.* 10, 100546 <https://doi.org/10.1016/j.ssmph.2020.100546>.
- Anderson, N.D., Damianakis, T., Kröger, E., Wagner, L.M., Dawson, D.R., Binns, M.A., Bernstein, S., Caspi, E., Cook, S.L., 2014. The benefits associated with volunteering among seniors: a critical review and recommendations for future research. *Psychol. Bull.* 140 <https://doi.org/10.1037/a0037610>.
- Aneshensel, C.S., 2010. In: Avison, W.R., Aneshensel, C.S., Schieman, S., Wheaton, B. (Eds.), *Neighborhood as a Social Context of the Stress Process* BT - Advances in the Conceptualization of the Stress Process: Essays in Honor of Leonard I. Pearlin. Springer New York, New York, NY, pp. 35–52. [https://doi.org/10.1007/978-1-4419-1021-9\\_3](https://doi.org/10.1007/978-1-4419-1021-9_3).
- Baadsgaard, M., Quitau, J., 2011. Danish registers on personal income and transfer payments. *Scand. J. Publ. Health* 39, 103–105. <https://doi.org/10.1177/1403494811405098>.
- Bailey, N., Kearns, A., Livingston, M., 2012. Place attachment in deprived neighbourhoods: the impacts of population turnover and social mix. *Hous. Stud.* 27 <https://doi.org/10.1080/02673037.2012.632620>.
- Baron, R.M., Kenny, D.A., 1986. The moderator-mediator variable distinction in social psychological research. Conceptual, strategic, and statistical considerations. *J. Pers. Soc. Psychol.* 51, 1173–1182. <https://doi.org/10.1037/0022-3514.51.6.1173>.
- Bassett, E., Moore, S., 2013. Gender differences in the social pathways linking neighborhood disadvantage to depressive symptoms in adults. *PLoS One* 8. <https://doi.org/10.1371/journal.pone.0076554>.
- Bender, A.M., Kawachi, I., Jørgensen, T., Pisinger, C., 2015. Neighborhood deprivation is strongly associated with participation in a population-based health check. *PLoS One* 10, e0129819.
- Benson, M.L., Fox, G.L., DeMaris, A., Van Wyk, J., 2003. Neighborhood disadvantage, individual economic distress and violence against women in intimate relationships. *J. Quant. Criminol.* 19, 207–235. <https://doi.org/10.1023/A:1024930208331>.
- Berg, P. van den, Timmermans, H., 2015. A multilevel path analysis of social networks and social interaction in the neighbourhood. *REGION 2*. <https://doi.org/10.18335/region.v2i1.32>.
- Berger-Schmitt, R., 2002. Considering social cohesion in quality of life assessments: concept and measurement. *Soc. Indic. Res.* 58, 403–428. <https://doi.org/10.1023/A:1015752320935>.
- Bjørner, J.B., Damsgaard, M.T., Watt, T., Bech, P., Rasmussen, N.K., Kristensen, T.S., Modvig, J., Thunedborg, K., 1997. Dansk manual til SF-36 : et spørgeskema om helbredsstatus (Danish manual for SF-36: A questionnaire on health status). Lif, S.I.
- Blair, A., Ross, N.A., Garipey, G., Schmitz, N., 2014. How do neighborhoods affect depression outcomes? A realist review and a call for the examination of causal pathways. *Soc. Psychiatr. Psychiatr. Epidemiol.* <https://doi.org/10.1007/s00127-013-0810-z>.
- Blanchard, C.M., Côté, I., Feeny, D., 2004. Comparing short form and RAND physical and mental health summary scores: results from total hip arthroplasty and high-risk primary-care patients. *Int. J. Technol. Assess. Health Care* 20. <https://doi.org/10.1017/S0266462304001011>.
- Booth, J., Ayers, S.L., Marsiglia, F.F., 2012. Perceived neighborhood safety and psychological distress: exploring protective factors. *J. Sociol. Soc. Welfare* 39.
- Bowling, A., Stafford, M., 2007. How do objective and subjective assessments of neighbourhood influence social and physical functioning in older age? Findings from a British survey of ageing. *Soc. Sci. Med.* 64, 2533–2549. <https://doi.org/10.1016/j.socscimed.2007.03.009>.
- Campbell, E., Henly, J.R., Elliott, D.S., Irwin, K., 2009. Subjective constructions of neighborhood boundaries: lessons from a qualitative study of four neighborhoods. *J. Urban Aff.* 31, 461–490. <https://doi.org/10.1111/j.1467-9906.2009.00450.x>.
- Chaix, B., Leyland, A.H., Sabel, C.E., Chauvin, P., Råstam, L., Kristersson, H., Merlo, J., 2006. Spatial clustering of mental disorders and associated characteristics of the neighbourhood context in Malmö, Sweden, in 2001. *J. Epidemiol. Community Health* 60. <https://doi.org/10.1136/jech.2005.040360>.
- Cheak-Zamora, N.C., Wyrwich, K.W., McBride, T.D., 2009. Reliability and validity of the SF-12v2 in the medical expenditure panel survey. *Qual. Life Res.* 18, 727–735. <https://doi.org/10.1007/s11136-009-9483-1>.
- Choi, Y.J., Matz-Costa, C., 2018. Perceived neighborhood safety, social cohesion, and psychological health of older adults. *Gerontol.* 58 <https://doi.org/10.1093/geront/gnw187>.
- Christensen, A.I., Davidsen, M., Kjoller, M., Juel, K., 2014. What characterizes persons with poor mental health? A nationwide study in Denmark. *Scand. J. Publ. Health* 42, 446–455. <https://doi.org/10.1177/1403494814532877>.
- Christensen, A.I., Davidsen, M., Koushede, V., Juel, K., 2020. Mental health and the risk of negative social life events: a prospective cohort study among the adult Danish population. *Scand. J. Publ. Health*, 1403494820944718. <https://doi.org/10.1177/1403494820944718>.
- Collins, C.R., Neal, J.W., Neal, Z.P., 2014. Transforming individual civic engagement into community collective efficacy: the role of bonding social capital. *Am. J. Community Psychol.* 54, 328–336. <https://doi.org/10.1007/s10464-014-9675-x>.
- Corcoran, R., Mansfield, R., de Bezenac, C., Anderson, E., Overbury, K., Marshall, G., 2018. Perceived neighbourhood affluence, mental health and wellbeing influence judgements of threat and trust on our streets: an urban walking study. *PLoS One* 13, e0202412.
- Cromley, E.K., Wilson-Genderson, M., Pruchno, R.A., 2012. Neighborhood characteristics and depressive symptoms of older people: local spatial analyses. *Soc. Sci. Med.* 75 <https://doi.org/10.1016/j.socscimed.2012.08.033>.
- Crump, C., Sundquist, K., Sundquist, J., Winkleby, M.A., 2011. Neighborhood deprivation and psychiatric medication prescription: a Swedish national multilevel study. *Ann. Epidemiol.* 21, 231–237. <https://doi.org/10.1016/j.annepidem.2011.01.005>.
- Delhey, J., Newton, K., 2003. Who trusts?: the origins of social trust in seven societies. *Eur. Soc. S.* 5 <https://doi.org/10.1080/1461669032000072256>.
- Diez Roux, A.V., 2016. Neighborhoods and health: what do we know? what should we do? *Am. J. Publ. Health* 106, 430–431. <https://doi.org/10.2105/AJPH.2016.303064>.
- Diez Roux, A.V., 2002. A glossary for multilevel analysis. *J. Epidemiol. Community Health*. <https://doi.org/10.1136/jech.56.8.588>.
- Drukker, M., van Os, J., 2003. Mediators of neighbourhood socioeconomic deprivation and quality of life. *Soc. Psychiatr. Psychiatr. Epidemiol.* 38, 698–706. <https://doi.org/10.1007/s00127-003-0690-8>.
- Echeverria, S.E., Diez-Roux, A.V., Link, B.G., 2004. Reliability of self-reported neighborhood characteristics. *J. Urban Health* 81, 682–701. <https://doi.org/10.1093/jurban/jth151>.
- Ehsan, A., Klaas, H.S., Bastianen, A., Spini, D., 2019. Social capital and health: a systematic review of systematic reviews. *SSM - Popul. Heal.* 8, 100425 <https://doi.org/10.1016/j.ssmph.2019.100425>.
- Ehsan, A.M., De Silva, M.J., 2015. Social capital and common mental disorder: a systematic review. *J. Epidemiol. Community Health*. <https://doi.org/10.1136/jech-2015-205868>.
- Erdem, Ö., Prins, R.G., Voorham, T.A.J.J., Van Lenthe, F.J., Burdorf, A., 2015. Structural neighbourhood conditions, social cohesion and psychological distress in The Netherlands. *Eur. J. Publ. Health* 25. <https://doi.org/10.1093/eurpub/ckv120>.
- Fahmi, M., Panjaitan, N.A., Habibie, I., Siregar, A.Y.M., Amarullah, G., Rahma, Sunjaya, D.K., 2019. Does your neighborhood protect you from being depressed? A study on social trust and depression in Indonesia. *BMC Publ. Health* 19, 1371. <https://doi.org/10.1186/s12889-019-7657-5>.
- Feld, S.L., 1981. The focused organization of social ties. *Am. J. Sociol.* 86, 1015–1035. <https://doi.org/10.1086/227352>.
- Fleishman, J.A., Selim, A.J., Kazis, L.E., 2010. Deriving SF-12v2 physical and mental health summary scores: a comparison of different scoring algorithms. *Qual. Life Res.* 19 <https://doi.org/10.1007/s11136-009-9582-z>.
- Flores, E.C., Fuhr, D.C., Bayer, A.M., Lescano, A.G., Thorogood, N., Simms, V., 2018. Mental health impact of social capital interventions: a systematic review. *Soc. Psychiatr. Psychiatr. Epidemiol.* 53 <https://doi.org/10.1007/s00127-017-1469-7>.
- Flowerdew, R., Manley, D.J., Sabel, C.E., 2008. Neighbourhood effects on health: does it matter where you draw the boundaries? *Soc. Sci. Med.* 66, 1241–1255. <https://doi.org/10.1016/j.socscimed.2007.11.042>.
- Fone, D.L., Dunstan, F., 2006. Mental health, places and people: a multilevel analysis of economic inactivity and social deprivation. *Health Place* 12, 332–344. <https://doi.org/10.1016/j.healthplace.2005.02.002>.
- Fong, D.Y.T., Lam, C.L.K., Mak, K.K., Lo, W.S., Lai, Y.K., Ho, S.Y., Lam, T.H., 2010. The Short Form-12 Health Survey was a valid instrument in Chinese adolescents. *J. Clin. Epidemiol.* 63, 1020–1029. <https://doi.org/10.1016/j.jclinepi.2009.11.011>.
- Foster, K.A., Aaron Hipp, J., 2011. Defining neighborhood boundaries for social measurement: advancing social work research. *Soc. Work. Res.* 35, 25–35. <https://doi.org/10.1093/swr/35.1.25>.
- Franzini, L., Caughy, M., Spears, W., Fernandez Esquer, M.E., 2005. Neighborhood economic conditions, social processes, and self-rated health in low-income neighborhoods in Texas: a multilevel latent variables model. *Soc. Sci. Med.* 61 <https://doi.org/10.1016/j.socscimed.2005.02.010>.
- Franzini, L., Spears, W., 2003. Contributions of social context to inequalities in years of life lost to heart disease in Texas, USA. *Soc. Sci. Med.* 57, 1847–1861. [https://doi.org/10.1016/S0277-9536\(03\)00018-2](https://doi.org/10.1016/S0277-9536(03)00018-2).
- Fujiwara, T., Kawachi, I., 2008. A prospective study of individual-level social capital and major depression in the United States. *J. Epidemiol. Community Health* 62, 627. <https://doi.org/10.1136/jech.2007.064261>. LP – 633.
- Galea, S., Ahern, J., Nandi, A., Tracy, M., Beard, J., Vlahov, D., 2007. Urban neighborhood poverty and the incidence of depression in a population-based cohort study. *Ann. Epidemiol.* 17, 171–179. <https://doi.org/10.1016/j.annepidem.2006.07.008>.
- Galster, G.C., 2013. The Mechanism(s) of Neighbourhood Effects: Theory, Evidence, and Policy Implications. Springer Netherlands. <https://doi.org/10.1007/978-94-007-2309-2-2>.
- Garon, J., Engelman, M., Gitlin, L., Szanton, S., 2016. Where does the neighborhood go? Trust, social engagement, and health among older adults in Baltimore City. *Health Place* 41, 58–66. <https://doi.org/10.1016/j.healthplace.2016.07.002>.
- Giordano, G.N., Lindström, M., 2011. Social capital and change in psychological health over time. *Soc. Sci. Med.* 72, 1219–1227. <https://doi.org/10.1016/j.socscimed.2011.02.029>.
- Giuliani, M.V., 2003. Theory of attachment and place attachment. In: Bonnes, M., Lee, T., Bonaiuto, M. (Eds.), *Psychological Theories for Environmental Issues. Psychological Theories for Environmental Issues*.

- Grannis, R., 2009. From the Ground up: Translating Geography into Community through Neighbor Networks. Princeton University Press. <https://doi.org/10.1177/0094306110367909x>.
- Grannis, R., 1998. The importance of trivial streets : residential streets and residential segregation. *Am. J. Sociol.* 103, 1530–1564. <https://doi.org/10.1086/231400>.
- Greene, G., Fone, D., Farewell, D., Rodgers, S., Paranjothy, S., Carter, B., White, J., 2020. Improving mental health through neighbourhood regeneration: the role of cohesion, belonging, quality and disorder. *Eur. J. Publ. Health* 30, 964–966. <https://doi.org/10.1093/eurpub/ckz221>.
- Haines, V.A., Beggs, J.J., Hurlbert, J.S., 2011. Neighborhood disadvantage, network social capital, and depressive symptoms. *J. Health Soc. Behav.* 52 <https://doi.org/10.1177/0022146510394951>.
- Harpam, T., Grant, E., Thomas, E., 2002. Measuring social capital within health surveys: key issues. *Health Pol. Plann.* 17 <https://doi.org/10.1093/heapol/17.1.106>.
- Helliwell, J.F., Putnam, R.D., 2004. The Social Context of Well-Being. *Philosophical Transactions of the Royal Society B: Biological Sciences*, pp. 1435–1446. <https://doi.org/10.1098/rstb.2004.1522>.
- Hox, J.J., Moerbeek, M., van de Schoot, R., 2018. Multilevel analysis - techniques and applications. In: *Multilevel Analysis, Quantitative Methodology Series*, third ed. Routledge. <https://doi.org/10.4324/9781315650982>.
- Huppert, F.A., 2009. Psychological well-being: evidence regarding its causes and consequences. *Appl. Psychol. Heal. Well-Being* 1, 137–164. <https://doi.org/10.1111/j.1758-0854.2009.01008.x>.
- Jakobsen, A., 2021. Neighborhood socioeconomic deprivation and psychiatric medication purchases. Different neighborhood delineations, different results? A nationwide register-based multilevel study. *Health & Place* 72. <https://doi.org/10.1016/j.healthplace.2021.102675>.
- Jakobsson, U., Westergren, A., Lindskov, S., Hagell, P., 2012. Construct validity of the SF-12 in three different samples. *J. Eval. Clin. Pract.* <https://doi.org/10.1111/j.1365-2753.2010.01623.x>.
- Jenks, G.F., 1967. The data model concept in statistical mapping. *Int. Yearb. Cartogr.* 7.
- Jensen, V.M., Rasmussen, A.W., 2011. Danish education registers. *Scand. J. Publ. Health* 39, 91–94. <https://doi.org/10.1177/1403494810394715>.
- Jivraj, S., Murray, E.T., Norman, P., Nicholas, O., 2019. The impact of life course exposures to neighbourhood deprivation on health and well-being: a review of the long-term neighbourhood effects literature. *Eur. J. Publ. Health*. <https://doi.org/10.1093/eurpub/ckz153>.
- Jonsson, K.R., Busfield, J., Södergren, M., Karen, M., Adjei, N.K., 2020. Social capital, deprivation and psychological well-being among young adolescents: a multilevel study from England and Wales. *Int. J. Environ. Res. Publ. Health* 17. <https://doi.org/10.3390/ijerph17103420>.
- Jørgensen, A., 2010. The Sense of Belonging in New Urban Zones of Transition. *Current Sociology* 58 (1). <https://doi.org/10.1177/0011392109348542>.
- Joshi, S., Mooney, S.J., Rundle, A.G., Quinn, J.W., Beard, J.R., Cerdá, M., 2017. Pathways from neighborhood poverty to depression among older adults. *Health Place* 43, 138–143. <https://doi.org/10.1016/j.healthplace.2016.12.003>.
- Juhász, A., Nagy, C., Páldy, A., Beale, L., 2010. Development of a Deprivation Index and its relation to premature mortality due to diseases of the circulatory system in Hungary, 1998–2004. *Soc. Sci. Med.* 70, 1342–1349. <https://doi.org/10.1016/j.socscimed.2010.01.024>.
- Julien, D., Richard, L., Gauvin, L., Kestens, Y., 2012. Neighborhood characteristics and depressive mood among older adults: an integrative review. *Int. psychogeriatrics* 24, 1207–1225. <https://doi.org/10.1017/S1041610211002894>.
- Kawachi, I., Berkman, L.F., 2015. Social capital, social cohesion, and health. In: *Social Epidemiology*, pp. 290–319. <https://doi.org/10.1093/med/9780195377903.003.0008>.
- Kawachi, I., Berkman, L.F., 2001. Social ties and mental health. *J. Urban Health* 78, 458–467. <https://doi.org/10.1093/jurban/78.3.458>.
- Keyes, C.L.M., 2007. Promoting and protecting mental health as flourishing: a complementary strategy for improving national mental health. *Am. Psychol.* 62 <https://doi.org/10.1037/0003-066X.62.2.95>.
- Kim, D., 2008. Blues from the neighborhood? Neighborhood characteristics and depression. *Epidemiol. Rev.* <https://doi.org/10.1093/epirev/mxn009>.
- Kim, J., 2010. Neighborhood disadvantage and mental health: the role of neighborhood disorder and social relationships. *Soc. Sci. Res.* 39, 260–271. <https://doi.org/10.1016/j.ssresearch.2009.08.007>.
- Kontodimopoulos, N., Pappa, E., Niakas, D., Tountas, Y., 2007. Validity of SF-12 summary scores in a Greek general population. *Health Qual. Life Outcomes* 5, 55. <https://doi.org/10.1186/1477-7525-5-55>.
- Kristiansen, S., Lund, R.L., 2022. The geography of gambling: a socio-spatial analysis of gambling machine location and area-level socio-economic status. *J. Gambl. Issues* 49, 44–67. <https://doi.org/10.4309/jgi.2022.49.2>.
- Leventhal, T., Brooks-Gunn, J., 2003. Moving to opportunity: an experimental study of neighborhood effects on mental health. *Am. J. Publ. Health* 93, 1576–1582. <https://doi.org/10.2105/ajph.93.9.1576>.
- Lewis, J.D., Weigert, A., 1985. Trust as a social reality. *Soc. Forces* 63. <https://doi.org/10.1093/sf/63.4.967>.
- Loureiro, A., Partidário, M.D.R., Santana, P., 2022. Strategic assessment of neighbourhood environmental impacts on mental health in the lisbon region (Portugal): a strategic focus and assessment framework at the local level. *Sustain. Times* 14. <https://doi.org/10.3390/su14031547>.
- Loureiro, A., Santana, P., Nunes, C., Almendra, R., 2019. The role of individual and neighborhood characteristics on mental health after a period of economic crisis in the lisbon region (Portugal): a multilevel analysis. *Int. J. Environ. Res. Publ. Health* 16. <https://doi.org/10.3390/ijerph16152647>.
- Ludwig, J., Duncan, G.J., Genetian, L.A., Katz, L.F., Kessler, R.C., Kling, J.R., Sanbonmatsu, L., 2013. Long-Term neighborhood effects on low-income families: evidence from moving to opportunity. *Am. Econ. Rev.* 103, 226–231.
- Lund, R.L., 2020. Moving to prosperity? The effect of prolonged exposure to neighborhood deprivation. *Scand. J. Educ. Res.* 64, 471–487. <https://doi.org/10.1080/00313831.2019.1577753>.
- Lund, R.L., 2019. Dissecting the Local. Aalborg Univ. Det Samf. Fak. Ph.D.-Serien.
- Lund, R.L., 2018. From the dark end of the street to the bright side of the road: automated redistricting of areas using physical barriers as dividers of social space. *Methodol. Innov.* 11 <https://doi.org/10.1177/2059799118814386>.
- Lund, R.L., Jørgensen, A., Riis, O.P., 2019. Social geographical patterns in membership of the established church in Denmark. *Nord. J. Relig. Soc.* 32, 55–70. <https://doi.org/10.18261/issn.1890-7008-2019-01-04>.
- Lynch, K., 1971. *The Image of the City*. MIT Press, Cambridge.
- MacKinnon, D.P., Lockwood, C.M., Williams, J., 2004. Confidence limits for the indirect effect: distribution of the product and resampling methods. *Multivariate Behav. Res.* 39 [https://doi.org/10.1207/s15327906mbr3901\\_4](https://doi.org/10.1207/s15327906mbr3901_4).
- Mair, C., Roux, A.V.D., Galea, S., 2008. Are neighbourhood characteristics associated with depressive symptoms? A review of evidence. *J. Epidemiol. Community Health* 62, 940. <https://doi.org/10.1136/jech.2007.066605>. LP – 946.
- Meijer, M., Engholm, G., Grittner, U., Bloomfield, K., 2013. A socioeconomic deprivation index for small areas in Denmark. *Scand. J. Publ. Health* 41, 560–569. <https://doi.org/10.1177/1403494813483937>.
- Merlo, J., Wagner, P., Austin, P.C., Subramanian, S.V., Leckie, G., 2018. General and specific contextual effects in multilevel regression analyses and their paradoxical relationship: a conceptual tutorial. *SSM - Popul. Heal.* 5 <https://doi.org/10.1016/j.ssmph.2018.05.006>.
- Messer, L.C., Vinikoor-Imler, L.C., Laraia, B.A., 2012. Conceptualizing neighborhood space: consistency and variation of associations for neighborhood factors and pregnancy health across multiple neighborhood units. *Health Place* 18, 805–813. <https://doi.org/10.1016/j.healthplace.2012.03.012>.
- Mouratidis, K., 2020. Neighborhood characteristics, neighborhood satisfaction, and well-being: the links with neighborhood deprivation. *Land Use Pol.* 99, 104886 <https://doi.org/10.1016/j.landusepol.2020.104886>.
- Mujahid, M.S., Diez Roux, A.V., Morenoff, J.D., Raghunathan, T., 2007. Assessing the measurement properties of neighborhood scales from psychometrics to econometrics. *Am. J. Epidemiol.* 165 <https://doi.org/10.1093/aje/kwm040>.
- Mujahid, M.S., Roux, V.D., Morenoff, J.D., Raghunathan, T.E., Cooper, R.S., Ni, H., Shea, S., 2008. Neighborhood characteristics and hypertension. *Epidemiology* 19, 590–598. <https://doi.org/10.1097/EDE.0b013e3181772cb2>.
- Nettle, D., Pepper, G.V., Jobling, R., Schroeder, K.B., 2014. Being there: a brief visit to a neighbourhood induces the social attitudes of that neighbourhood. *PeerJ* 2, e236. <https://doi.org/10.7717/peerj.236>.
- Paczkowski, M.M., Galea, S., 2010. Sociodemographic characteristics of the neighborhood and depressive symptoms. *Curr. Opin. Psychiatr.* <https://doi.org/10.1097/YCO.0b013e32833ad70b>.
- Parenteau, M.-P., Sawada, M.C., 2011. The modifiable areal unit problem (MAUP) in the relationship between exposure to NO2 and respiratory health. *Int. J. Health Geogr.* 10, 58. <https://doi.org/10.1186/1476-072X-10-58>.
- Pedersen, C.B., 2011. The Danish Civil registration system. *Scand. J. Publ. Health* 39, 22–25. <https://doi.org/10.1177/1403494810387965>.
- Peterson, L.E., Tsai, A.C., Petterson, S., Litaker, D.G., 2009. Rural–urban comparison of contextual associations with self-reported mental health status. *Health Place* 15, 125–132. <https://doi.org/10.1016/j.healthplace.2008.03.001>.
- Porges, S.W., 2007. The polyvagal perspective. *Biol. Psychol.* 74, 116–143. <https://doi.org/10.1016/j.biopsycho.2006.06.009>.
- Preacher, K.J., Hayes, A.F., 2008. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav. Res. Methods* 40, 879–891. <https://doi.org/10.3758/BRM.40.3.879>.
- Preacher, K.J., Selig, J.P., 2012. Advantages of Monte Carlo confidence intervals for indirect effects. *Commun. Methods Meas.* 6 <https://doi.org/10.1080/19312458.2012.679848>.
- Preacher, K.J., Zyphur, M.J., Zhang, Z., 2010. A general multilevel SEM framework for assessing multilevel mediation. *Psychol. Methods* 15, 209–233. <https://doi.org/10.1037/a0020141>.
- Prior, L., Manley, D., Jones, K., 2018. Stressed out? An investigation of whether allostatic load mediates associations between neighbourhood deprivation and health. *Health Place* 52. <https://doi.org/10.1016/j.healthplace.2018.05.003>.
- Putnam, R.D., 2000. *Bowling Alone : the Collapse and Revival of American Community*. Simon & Schuster, New York, New York.
- Richardson, R., Westley, T., Gariépy, G., Austin, N., Nandi, A., 2015. Neighborhood socioeconomic conditions and depression: a systematic review and meta-analysis. *Soc. Psychiatr. Psychiatr. Epidemiol.* 50, 1641–1656. <https://doi.org/10.1007/s00127-015-1092-4>.
- Rios, R., Aiken, L.S., Zautra, A.J., 2012. Neighborhood contexts and the mediating role of neighborhood social cohesion on health and psychological distress among hispanic and non-hispanic residents. *Ann. Behav. Med.* 43, 50–61. <https://doi.org/10.1007/s12160-011-9306-9>.
- Robinette, J.W., Charles, S.T., Mogle, J.A., Almeida, D.M., 2013. Neighborhood cohesion and daily well-being: results from a diary study. *Soc. Sci. Med.* 96 <https://doi.org/10.1016/j.socscimed.2013.07.027>.
- Rose, N., Birk, R., Manning, N., 2021. Towards neuroecosociality: mental health in adversity. *Theor. Cult. Soc.*, 0263276420981614 <https://doi.org/10.1177/0263276420981614>.
- Ross, C.E., 2000. Neighborhood disadvantage and adult depression. *J. Health Soc. Behav.* 41, 177–187. <https://doi.org/10.2307/2676304>.



- Sampson, R.J., 2011. Neighborhood effects, causal mechanisms and the social structure of the city. *Anal. Sociol. Soc. Mech.* 227–249. <https://doi.org/10.1017/CBO9780511921315.012>.
- Sampson, R.J., Morenoff, J.D., Gannon-Rowley, T., 2002. Assessing “neighborhood effects”: social processes and new directions in research. *Annu. Rev. Sociol.* 28, 443–478. <https://doi.org/10.1146/annurev.soc.28.110601.141114>.
- Sampson, R.J., Raudenbush, S.W., Earls, F., 1997. Neighborhoods and violent crime: a multilevel study of collective efficacy. *Science* 277 (80). <https://doi.org/10.1126/science.277.5328.918>.
- Santini, Z.I., Jose, P.E., Koyanagi, A., Meilstrup, C., Nielsen, L., Madsen, K.R., Koushede, V., 2020. Formal social participation protects physical health through enhanced mental health: a longitudinal mediation analysis using three consecutive waves of the Survey of Health, Ageing and Retirement in Europe (SHARE). *Soc. Sci. Med.* 251 <https://doi.org/10.1016/j.socscimed.2020.112906>.
- Santini, Z.I., Koyanagi, A., Tyrovolas, S., Mason, C., Haro, J.M., 2015. The association between social relationships and depression: a systematic review. *J. Affect. Disord.* 175, 53–65. <https://doi.org/10.1016/j.jad.2014.12.049>.
- Scannell, L., Gifford, R., 2017. The experienced psychological benefits of place attachment. *J. Environ. Psychol.* 51, 256–269. <https://doi.org/10.1016/j.jenvp.2017.04.001>.
- Scannell, L., Gifford, R., 2010. Defining place attachment: a tripartite organizing framework. *J. Environ. Psychol.* 30 <https://doi.org/10.1016/j.jenvp.2009.09.006>.
- Schmidt, N.M., Nguyen, Q.C., Kehm, R., Osypuk, T.L., 2020. Do changes in neighborhood social context mediate the effects of the moving to opportunity experiment on adolescent mental health? *Heal. Place* 63. <https://doi.org/10.1016/j.healthplace.2020.102331>.
- Schüle, S.A., Hilz, L.K., Dreger, S., Bolte, G., 2019. Social inequalities in environmental resources of green and blue spaces: a review of evidence in the WHO European region. *Int. J. Environ. Res. Publ. Health* 16, 1216. <https://doi.org/10.3390/ijerph16071216>.
- Semenza, J.C., March, T.L., Bontempo, B.D., 2007. Community-initiated urban development: an ecological intervention. *J. Urban Health* 84. <https://doi.org/10.1007/s11524-006-9124-8>.
- Shrout, P.E., Bolger, N., 2002. Mediation in experimental and nonexperimental studies: new procedures and recommendations. *Psychol. Methods* 7. <https://doi.org/10.1037/1082-989X.7.4.422>.
- Shumaker, S.A., Taylor, R.B., 1983. Toward a clarification of people-place relationships: a model of attachment to place. In: *Environmental Psychology: Directions and Perspectives*.
- Silva, M., Loureiro, A., Cardoso, G., 2016. Social determinants of mental health: a review of the evidence. *Eur. J. Psychiatr.* 30, 259–292.
- Snijders, T., Bosker, R., 1999. *Multilevel Analysis: an Introduction to Basic and Advanced Multilevel Analysis*. Sage, Thousand Oaks CA.
- Sobel, M.E., 1982. Asymptotic confidence intervals for indirect effects in structural equation models. *Socio. Methodol.* 13 <https://doi.org/10.2307/270723>.
- Statistics Denmark, 2017. *Statistical Yearbook 2017*.
- Takagi, D., Kondo, K., Kawachi, I., 2013. Social participation and mental health: moderating effects of gender, social role and rurality. *BMC Publ. Health* 13. <https://doi.org/10.1186/1471-2458-13-701>.
- The Danish Ministry of Food, 2011. *Agriculture and Fisheries, the Danish Ministry of the interior and health*. In: *Landdistriktsrederegørelse. Regeringens Redegørelse Til Folketinget*.
- Truong, K.D., Ma, S., 2006. A systematic review of relations between neighborhoods and mental health. *J. Ment. Health Pol. Econ.* 9, 137–154.
- Tucker, G., Adams, R., Wilson, D., 2016. The case for using country-specific scoring coefficients for scoring the SF-12, with scoring implications for the SF-36. *Qual. Life Res.* 25 <https://doi.org/10.1007/s11136-015-1083-7>.
- UNESCO Institute for Statistics, 2012. *International Standard Classification of Education: ISCED 2011*. UNESCO Institute for Statistics Montreal.
- Valeri, L., VanderWeele, T.J., 2013. Mediation analysis allowing for exposure-mediator interactions and causal interpretation: theoretical assumptions and implementation with SAS and SPSS macros. *Psychol. Methods* 18. <https://doi.org/10.1037/a0031034>.
- Van Deurzen, I., Rod, N.H., Christensen, U., Hansen, Å.M., Lund, R., Dich, N., 2016. Neighborhood perceptions and allostatic load: evidence from Denmark. *Health Place* 40, 1–8. <https://doi.org/10.1016/j.healthplace.2016.04.010>.
- van Ham, M., Manley, D., Ham, M. Van, Manley, D., 2012. Neighbourhood effects research at a crossroads. Ten challenges for future research Introduction. *Environ. Plann.* 44, 2787–2793. <https://doi.org/10.1068/a45439>.
- VanderWeele, T.J., 2016. Mediation analysis: a practitioner’s guide. *Annu. Rev. Publ. Health.* <https://doi.org/10.1146/annurev-publhealth-032315-021402>.
- Visser, K., Bolt, G., Finkenauer, C., Jonker, M., Weinberg, D., Stevens, G.W.J.M., 2021. Neighbourhood deprivation effects on young people’s mental health and well-being: a systematic review of the literature. *Soc. Sci. Med.* <https://doi.org/10.1016/j.socscimed.2020.113542>.
- Ware John, E., Kosinski Mark, D., Keller Susan, D., Ware, J., Mark, K., Keller Susan, D., 1996. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med. Care* 34, 220–233. <https://doi.org/10.1097/00005650-199603000-00003>.
- Ware, J., Kosinski, M., Keller, S., 1998. *SF-12: How to Score the SF-12 Physical and Mental Health Summary Scales*.
- White, J., Greene, G., Farewell, D., Dunstan, F., Rodgers, S., Lyons, R.A., Humphreys, I., John, A., Webster, C., Phillips, C.J., Fone, D., 2017. Improving mental health through the regeneration of deprived neighborhoods: a natural experiment. *Am. J. Epidemiol.* 186, 473–480. <https://doi.org/10.1093/aje/kwx086>.
- Wilson-Genderson, M., Pruchno, R., 2013. Effects of neighborhood violence and perceptions of neighborhood safety on depressive symptoms of older adults. *Soc. Sci. Med.* 85 <https://doi.org/10.1016/j.socscimed.2013.02.028>.
- York Cornwell, E., Behler, R.L., 2015. Urbanism, neighborhood context, and social networks. *City Community* 14, 311–335. <https://doi.org/10.1111/cico.12124>.
- Young, A.F., Russell, A., Powers, J.R., 2004. The sense of belonging to a neighbourhood: can it be measured and is it related to health and well being in older women? *Soc. Sci. Med.* 59, 2627–2637. <https://doi.org/10.1016/j.socscimed.2004.05.001>.
- Zhang, Z., Zyphur, M.J., Preacher, K.J., 2009. Testing multilevel mediation using hierarchical linear models: problems and solutions. *Organ. Res. Methods* 12, 695–719. <https://doi.org/10.1177/1094428108327450>.
- Zimmerman, G.M., Messner, S.F., 2010. Neighborhood context and the gender gap in adolescent violent crime. *Am. Socio. Rev.* 75, 958–980. <https://doi.org/10.1177/0003122410386688>.