

## Running on empty

*a longitudinal global study of psychological well-being among runners during the COVID-19 pandemic*

Vistisen, Helene Tilma; Sønderskov, Kim Mannemar; Dinesen, Peter Thisted; Brund, René Børge Korsgaard; Nielsen, Rasmus Østergaard; Østergaard, Søren Dinesen

*Published in:*  
BMJ Open

*DOI (link to publication from Publisher):*  
[10.1136/bmjopen-2022-063455](https://doi.org/10.1136/bmjopen-2022-063455)

*Creative Commons License*  
CC BY-NC 4.0

*Publication date:*  
2022

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Vistisen, H. T., Sønderskov, K. M., Dinesen, P. T., Brund, R. B. K., Nielsen, R. Ø., & Østergaard, S. D. (2022). Running on empty: a longitudinal global study of psychological well-being among runners during the COVID-19 pandemic. *BMJ Open*, 12(9), Article e063455. <https://doi.org/10.1136/bmjopen-2022-063455>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# BMJ Open Running on empty: a longitudinal global study of psychological well-being among runners during the COVID-19 pandemic

Helene Tilma Vistisen <sup>1,2</sup> Kim Mannemar Sønderskov <sup>3,4</sup>  
 Peter Thisted Dinesen <sup>5,6</sup> René Børge Korsgaard Brund <sup>7</sup>  
 Rasmus Østergaard Nielsen <sup>8,9</sup> Søren Dinesen Østergaard <sup>1,2</sup>

**To cite:** Vistisen HT, Sønderskov KM, Dinesen PT, *et al.* Running on empty: a longitudinal global study of psychological well-being among runners during the COVID-19 pandemic. *BMJ Open* 2022;**12**:e063455. doi:10.1136/bmjopen-2022-063455

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-063455>).

Received 01 April 2022  
 Accepted 10 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Søren Dinesen Østergaard; [soeoes@rm.dk](mailto:soeoes@rm.dk)

## ABSTRACT

**Objectives** There are indications that the COVID-19 pandemic has had a profound negative effect on psychological well-being. Here, we investigated this hypothesis using longitudinal data from a large global cohort of runners, providing unprecedented leverage for understanding how the temporal development in the pandemic pressure relates to well-being across countries.

**Design** Prospective cohort study.

**Setting** Global.

**Participants** We used data from the worldwide Garmin-RUNSAFE cohort that recruited runners with a Garmin Connect account, which is used for storing running activities tracked by a Garmin device. A total of 7808 Garmin Connect users from 86 countries participated.

**Primary and secondary outcome measures** From 1 August 2019 (prepandemic) to 31 December 2020, participants completed surveys every second week that included the five-item WHO Well-Being Index (WHO-5). Pandemic pressure was proxied by the number of COVID-19-related deaths per country, retrieved from the Coronavirus Resource Centre at Johns Hopkins University. Panel data regression including individual- and time-fixed effects was used to study the association between country-level COVID-19-related deaths over the past 14 days and individual-level self-reported well-being over the past 14 days.

**Results** The 7808 participants completed a total of 125 409 WHO-5 records over the study period. We found a statistically significant inverse relationship between the number of COVID-19-related deaths and the level of psychological well-being—independent of running activity and running injuries (a reduction of 1.42 WHO-5 points per COVID-19-related death per 10 000 individuals,  $p < 0.001$ ).

**Conclusions** This study suggests that the COVID-19 pandemic has had a negative effect on the psychological well-being of the affected populations, which is concerning from a global mental health perspective.

## INTRODUCTION


Beyond its obvious negative health consequences for those directly infected with coronavirus, the COVID-19 pandemic—and the ensuing public health measures implemented

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Psychological well-being was tracked every second week over several months prior to and during the COVID-19 pandemic.
- ⇒ The study was based on data from 7808 participants representing 86 countries.
- ⇒ The participants were self-enrolled runners, who are likely more psychologically robust than the general population.
- ⇒ Data on nationwide and regional lockdowns from the 86 countries were not available.

to prevent its spreading (eg, lockdowns and restrictions on social gatherings)—is likely to have had adverse effects on psychological well-being more broadly due to, *inter alia*, the uncertainty, the disruption of everyday routines, and the social disconnectedness it has induced.<sup>1 2</sup>

Previous longitudinal studies, tracking the development in psychological well-being over time by means of surveys, have provided initial evidence documenting the negative consequences of the COVID-19 pandemic. While informative, these studies generally suffer from one or more significant drawbacks. First, only a subset of these studies has a prepandemic baseline measurement that is necessary to enable any inferences about the consequences of the pandemic.<sup>3–12</sup> Further, even if prepandemic benchmarks are available, they are typically few and dating back a longer period of time (often years) before the onset of the pandemic.<sup>4–14</sup> This compromises the value of the prepandemic measure, and, by implication, the credibility of any observed change in well-being after the onset of the pandemic. Several pre-pandemic measurements taken over a period leading directly up to the pandemic, would strengthen the case



**Figure 1** Flow chart of the study population and WHO-5 observations. WHO-5, five-item WHO Well-Being Index.

further for the pandemic causing an observed decline in psychological well-being. Second, beyond the consequences of the COVID-19 pandemic in toto, previous studies—including our own<sup>13 15–17</sup>—have produced limited knowledge about how psychological well-being covaries with pandemic pressure (ie, the severity of the COVID-19 pandemic) given the absence of systematic post-pandemic measurements of well-being. If psychological well-being changes in tandem with the ebb and tide of the pandemic waves, it strengthens the claim of the pandemic influencing well-being. Third, the existing results are typically from single-country studies.<sup>3 9 18 19</sup> While this is a natural starting point, this means that any (inverse) correspondence between pandemic pressure and psychological well-being could be due to other temporal changes that causes changes in well-being (eg, seasonal changes in daylight or weather).<sup>16 20</sup> Using data from several countries with variation in pandemic pressure and seasonal conditions can alleviate this concern, and would therefore lend further credibility to the robustness of the negative effect of the pandemic pressure on psychological well-being.

Against the backdrop of previous studies and their shortcomings, the aim of this study was to investigate the dose–response relationship between pandemic pressure (proxied by number of COVID-19-related deaths) and psychological well-being using shortly-spaced individual-level panel survey data from more than 80 countries with extensive measurement points both before and after the inception of the pandemic. The data stems from a large global cohort of runners (the Garmin-RUNSAFE Running Health Study<sup>21</sup>), and therefore, to fortify our results against idiosyncratic features of this sample, we used auxiliary data on the participants' running-related characteristics (activities and injuries), to establish that the relationship between the pandemic pressure and psychological well-being is independent of changes in

these characteristics and hence likely generalises more broadly.

## METHODS

### Data source

We used data from the international worldwide Garmin-RUNSAFE Running Health Study that recruited English-speaking runners aged 18+ with a Garmin Connect account. Garmin connect is a tool for storing and sharing running activities from a Garmin device.<sup>21</sup> Enrolment was open from 1 August 2019 (prepandemic) to 31 December 2020. For further details on the recruitment, see Nielsen *et al.*<sup>21</sup>

### Data collection

At enrolment, the participants in the Garmin-RUNSAFE Running Health Study provided information on country of residence and date of birth. Furthermore, they gave access to daily information on running distance (in metres) during follow-up (from enrolment to 31 December 2020) from their Garmin Connect account. From the time of enrolment to 31 December 2020, the RUNSAFE participants were asked to complete surveys every 2 weeks (sent via email) that included the five-item WHO Well-Being Index (WHO-5)<sup>22</sup>—a psychometrically valid and widely used measure of psychological well-being experienced over the past 2 weeks. The five WHO-5 items are: 'I have felt cheerful and in good spirit', 'I have felt calm and relaxed', 'I have felt active and vigorous', 'I woke up feeling fresh and rested' and 'My daily life has been filled with things that interest me'. Each item is scored from 0 (none of the time) to 5 (all the time). The WHO-5 total score is calculated by adding the individual item scores and multiplying by four (ranges from 0 (complete lack of well-being) to 100 (maximum well-being)). The participants also provided weekly information on

**Table 1** Characteristics of the 7808 participants at enrolment

	No of participants (unit)
Sex	
Women, n (%)	1753 (22.5)
Men, n (%)	5935 (76.0)
Missing, n (%)	120 (1.5)
Age, mean years (SD)	47.3 (10.6)
18–24, n (%)	105 (1.3)
25–34, n (%)	788 (10.1)
35–44, n (%)	2227 (28.5)
45–54, n (%)	2841 (36.4)
55–64, n (%)	1372 (17.6)
65–74, n (%)	420 (5.4)
75+, n (%)	42 (0.5)
Missing, n (%)	13 (0.2)
Continent	
Asia*, n (%)	55 (0.7)
Africa†, n (%)	145 (1.9)
North America‡, n (%)	3118 (39.9)
USA, n (%)	2727 (34.9)
Canada, n (%)	370 (4.7)
South America§, n (%)	38 (0.5)
Europe¶, n (%)	4436 (56.8)
UK, n (%)	956 (12.2)
Germany, n (%)	409 (5.2)
Italy, n (%)	382 (4.9)
Denmark, n (%)	376 (4.8)
France, n (%)	334 (4.3)
Netherlands, n (%)	291 (3.7)
Spain, n (%)	282 (3.6)
Sweden, n (%)	282 (3.6)
Norway, n (%)	192 (2.5)
Belgium, n (%)	135 (1.7)
Oceania**, n (%)	16 (0.2)

\*Countries participating in Asia: Taiwan, Qatar, Saudi Arabia, Cambodia, Malaysia, Cyprus, UAE, Turkey, Thailand, Singapore, India, Japan, Israel, Brunei, Lebanon, Indonesia, Hong Kong, China.

†Countries participating in Africa: Sudan, Eswatini, Namibia, Algeria, Egypt, South Africa, Mauritius, Morocco, Uganda, Zimbabwe, Kenya, Reunion.

‡USA and Canada accounts for 99% of the participants from North America. Other participating countries in North America: Panama, Costa Rica, Honduras, British Virgin Islands, Mexico, Dominican Republic, Greenland, Barbados, Guatemala.

§Countries participating in South America: Venezuela, Bolivia, Ecuador, Argentina, Peru, Chile, Falkland Islands, Brazil, Colombia, French Guiana.

¶The 10 countries in Europe with the highest number of participants. These 10 countries accounts for 82% of the participants from Europe. Other participating countries in Europe: Luxembourg, Slovenia, Portugal, Romania, Austria, Croatia, Switzerland, Ireland, Bosnia and Herzegovina, Iceland, Russia, Ukraine, Finland, Faroe Islands, Lithuania, Slovakia, Montenegro, Malta, Greece, Czechia, Serbia, Poland.

\*\*Countries participating in Oceania: French Polynesia, New Zealand, Australia.

running-related injuries/problems. Specifically, they were asked to indicate which day in the past week a running-related injury/problem interfered with their running activity and/or affected their activities of daily living.

### Patient or public involvement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

### Study population

For this study, we used data from all participants in the Garmin-RUNSAFE Running Health Study with information on country of residence and with ≥1 completed WHO-5 questionnaire on psychological well-being.

### Data on COVID-19-related deaths

The daily number of COVID-19-related deaths per country was retrieved from the Coronavirus Resource Centre at John Hopkins University.<sup>23</sup> The few instances (0.19%) of negative daily deaths (due to changing definitions) were replaced by the mean number of deaths from the two neighbouring dates. We opted for using country-specific death rates because it, unlike other measures, presumably is highly comparable within countries over time. Other measures like incidence rates of COVID-19 and transmissibility depends heavily on test rates, which varied substantially within countries over time due to variation in availability of tests, pandemic pressure etc.


### Statistical analysis

The data described above were organised in person-week units. Specifically, for each week in the follow-up period (1 August 2019–31 December 2020), we computed participant-level WHO-5 total scores (ie, their well-being the past 14 days), running distance over the past 14 days (in metres), running-related injuries/problems (days affected of the past 14 days), as well as the number of COVID-19-related deaths per 10 000 inhabitants (in the country of the participant) for the past 14 days. The rationale behind the weekly and not 2 weekly organisation was that even though the WHO-5 questionnaires were sent out every second week, responses were returned throughout the subsequent 14-day deadline period. If a participant filled in the WHO-5 twice within the same week, the last WHO-5 total score was used.

The following analyses were carried out: First, the cohort was characterised using descriptive statistics. Subsequently, the relationship between country-level COVID-19-related deaths over the past 14 days and the level of psychological well-being over the past 14 days (WHO-5 total score) was assessed via a linear regression model including individual- and time-fixed effects, which reduces the risk of confounding from stable individual- and country-level characteristics as well as general trends in well-being during the study period:

$$WHO5_{it} = \beta_0 + \beta_1 Deaths_{it} + \beta_2 RunningActivity_{it} + \beta_3 Injury_{it} + a_i + u_t + \epsilon_{it}$$






**Figure 2** Number of participants (orange bars), COVID-19 deaths (grey bars) and mean WHO-5 total score (red line) over the course of the study. The line representing the mean WHO-5 total score is generated using a lowess smoother. The orange bars represent the number of participants having completed the WHO-5 at least once in the specific month. WHO-5, five-item WHO Well-Being Index.

Here,  $WHO-5_{it}$  is the WHO-5 total score for individual  $i$  for the time period  $t$  (past 14 days),  $Deaths_{it}$  is the number of deaths per 10 000 inhabitants in  $i$ 's country of residence over the time period  $t$ ,  $RunningActivity_{it}$  is  $i$ 's running activity (total metres) over time period  $t$ , and  $injury_{it}$  is the number of days over time period  $t$  where  $i$ 's activity was affected by a running-related injury/problem. The three remaining terms represent unobserved factors affecting the WHO-5 total score:  $a_i$  is time-invariant and individual-specific,  $u_t$  is individual-invariant and time-specific, and  $\epsilon_{it}$  represents unobserved determinants of the WHO-5 total score that vary across both individual and time. To remove  $a_i$  we included a full set of individual-level fixed effects, and to remove  $u_t$  we included time-fixed effects. Subsequently, we ran the same analysis for each of the five WHO-5 items (replacing  $WHO-5_{it}$  in the

equation shown above). The rationale behind this analytical model is illustrated in the directed acyclic graph shown in online supplemental figure 1. To check the robustness of the model, we conducted leave-one-out analysis excluding one country from the model at the time. As secondary analyses, to explore potential non-linear effects of the number of COVID-19-related deaths, square root, natural logarithmic and quadratic terms were employed (see online supplemental methods for further description).

Finally, to test whether the RUNSAFE participants had higher psychological well-being than the general population (a priori hypothesis), we compared the WHO-5 total scores of the Danish RUNSAFE participants with the WHO-5 total scores from the first three waves of the COVID-19 Consequences Denmark Panel Survey



**Figure 3** The association between COVID-19-related deaths per 10 000 and psychological well-being (WHO-5 total score). WHO-5, five-item WHO Well-Being Index.

**Table 2** Individual fixed-effects linear-regression analyses with time fixed effects (crude\* and adjusted† model)

	Regression coefficient ( $\beta_1 Deaths_{it}$ ) (95% CI)	P value
Crude model*		
WHO-5 total score	-1.48 (-2.47 to -0.49)	0.004
Individual WHO-5 item scores (0–20)		
Interest	-0.40 (-0.63 to -0.17)	<0.001
Fresh	-0.20 (-0.35 to -0.05)	0.011
Vigorous	-0.25 (-0.52 to 0.01)	0.061
Relaxed	-0.25 (-0.39 to -0.11)	<0.001
Cheerful	-0.38 (-0.63 to -0.13)	0.003
Adjusted model†		
WHO-5 total score	-1.42 (-2.16 to -0.67)	<0.001
Individual WHO-5 item scores (0–20)		
Interest	-0.40 (-0.60 to -0.20)	<0.001
Fresh	-0.20 (-0.30 to -0.10)	<0.001
Vigorous	-0.20 (-0.39 to 0.02)	0.032
Relaxed	-0.27 (-0.40 to -0.15)	<0.001
Cheerful	-0.34 (-0.55 to -0.13)	0.002

\*Observations: 125 409. Individuals: 7808. Model:

$$WHO5_{it} = \beta_0 + \beta_1 Deaths_{it} + a_i + u_t + \epsilon_{it}$$

†Observations: 84 679. Individuals: 6222. Model:

$$WHO5_{it} = \beta_0 + \beta_1 Deaths_{it} + \beta_2 RunningActivity_{it} + \beta_3 Injury_{it} + a_i + u_t + \epsilon_{it}$$

where  $Deaths_{it}$  is a numerical discrete variable measuring the number of deaths per 10 000 inhabitants (cf. table 1) in  $i$ 's country of residence at time period  $t$  ( $t$  represents periods of 14 days),  $RunningActivity_{it}$  is a continuous variable measuring  $i$ 's running activity (total meters) at time period  $t$ ,  $Injury_{it}$  measures the number of days where  $i$ 's activity was affected by a running injury or problem at time period  $t$ . The three remaining terms represent unobserved factors affecting  $WHO5_{it}$ :  $a_i$  is time-invariant and individual-specific;  $u_t$  is unit-invariant and time-specific; and  $\epsilon_{it}$  represents unobserved determinants of  $WHO5_{it}$  that vary across both individual and time. To remove  $a_i$ , we included a full set of individual-level fixed effects, and to remove  $u_t$  we included time-fixed effects.

WHO-5, five-item WHO Well-Being Index.

2020.<sup>13 15 16</sup> The WHO-5 total scores from the COVID-19 Consequences Denmark Panel Survey 2020 respondents were weighted on gender, age, education, region and political party choice in the last election in order to render them representative of the Danish population. Only WHO-5 data from overlapping periods of data collection in the two surveys were included, namely 31 March 2020–6 April 2020; 22 April 2020–30 April 2020 and 20 November 2020–8 December 2020.<sup>13 15 16</sup>

All analyses were carried out using Stata V.17.0 (StataCorp) with 0.05 as the threshold for statistical significance.

## RESULTS

In the period from 1 August 2019 to 31 December 2020, a total of 7808 RUNSAFE-participants completed the WHO-5 questionnaire at least once. Data from these 7808

participants were included in the analyses (see figure 1). The characteristics of the participants are listed in table 1.

The participants covered 86 different countries, the age range was 18–88 years, mean age was 47.3 years (SD=10.61) and 76% were men. The maximum follow-up was 17 months including 39 biweekly WHO-5 questionnaires, and 75 weekly injury questionnaires. The total number of completed WHO-5 questionnaires was 125 409 and the median number of completed WHO-5 questionnaires among the 7808 participants was 12 (IQR: 3–31). A total of 980 (12.6%) of the participants had completed the WHO-5 only once and thereby only contributed to the estimation of the country- and the time fixed effects. For an illustration of the distribution of participants and completed WHO-5 questionnaires across countries, see online supplemental figure 2.

Among the 7808 respondents, 7175 (91.9%) had tracked their running activity through Garmin Connect at least once (with a total of 230 169 weeks with information on running activity), and 7759 (99.4%) had filled out the weekly questionnaire about running-related injuries at least once (with a total of 257 171 weeks with information on injuries). For an illustration of the tracking of running activity and completed injury questionnaires over the course of the study, see online supplemental figure 3.

The range in number of COVID-19-related deaths per 10 000 (within a country) during a fourteen-day period was 0 to 3.65 with a median of 0.02 (IQR: 0.00–0.35) in the study period, and a median of 0.31 (IQR: 0.04–0.59) in the period from March 2020 to December 2020. For an illustration of the number of COVID-19-related deaths, the number of study participants, and the level of psychological well-being of these participants over the study period (see figure 2).

The linear association between the number of COVID-19-related deaths per 10 000 and psychological well-being (WHO-5 total score) is illustrated in figure 3 and reported in table 2.

The results show a statistically significant inverse relationship (regression coefficient of -1.42, 95% CI -2.16 to -0.67), which remained when excluding running activity and running related injuries/problems from the model (table 2) and when leaving specific countries out of the analysis one at the time (online supplemental table 1). The number of COVID-19-related deaths was also inversely associated with the five individual WHO-5 items (table 2). The results of the three non-linear analyses were also consistent with an inverse relationship between the number of COVID-19-related deaths per 10 000 and psychological well-being (table 2 and online supplemental table 2). Specifically, all analyses showed that the strength of the inverse relationship decreased at higher levels of COVID-19-related deaths (see online supplemental figure 4). The results of the quadratic model indicated that the relationship could be positive at very high levels of COVID-19 related deaths (approximately  $\geq 2.0$  COVID-19-related deaths per 10 000 inhabitants). This specific

finding is, however, uncertain, because of few observations with very high levels of COVID-19-related deaths (out of the 125 409 person-week observations, only 1974 (1.6%) had a rate  $\geq 2.0$  deaths per 10 000 inhabitants).

Finally, and as expected, the psychological well-being of the participants in the Garmin-RUNSAFE Running Health Study (mean WHO-5 total score of 71.6, 95% CI 70.0 to 73.2) was substantially higher than that of the participants from the COVID-19 Consequences Denmark Panel Survey 2020 (mean WHO-5 total score of 63.2, 95% CI 62.7 to 63.7), when compared across the same time periods.

## DISCUSSION

In this longitudinal study of 7808 runners from 86 countries, we found a statistically significant inverse relationship between the number of COVID-19-related deaths and the level of psychological well-being, which was independent of running activity and running injuries. These results were generally robust across models and sensitivity (leave-one-out) analyses.

To our knowledge, this study is the first to have tracked the psychological well-being of individuals from >80 countries over several months prior to and during the COVID-19 pandemic. The results bolsters and furthers findings from studies using less fine-grained data and less rigorous designs in showing that there is a dynamic inverse relationship between the pandemic pressure and the level of psychological well-being.<sup>3-14</sup> They are also in line with studies having focused on the opposite of psychological well-being during the COVID-19 pandemic, namely symptoms of anxiety and depression, where a positive relationship with the pandemic pressure has been the most consistent finding.<sup>24-28</sup> Irrespective of the definition of outcome, this body of literature clearly suggests that the COVID-19 pandemic is not only a global crisis from a physical health perspective, but also from a mental health/psychological perspective, as acknowledged by the WHO.<sup>29</sup>

Although this study has strengths, in particular due to the availability of fine-grained prepandemic and in-pandemic data on psychological well-being from many countries across continents, there are also important limitations to take into account. First, participants in the survey are self-enrolled and the sample is therefore probably not representative of runners from the included countries, and—given the heterogeneous participation patterns across countries (table 1; online supplemental figure 2)—certainly not representative of the global population of runners. Second, participation varies over time and there are clear signs of panel attrition over the study period, which also raises questions about generalisability. The inclusion of individual fixed effects, and by implication, country fixed effects, alleviates some of this concern, as it removes the influence of individual-level and country-level variables. Nevertheless, generalisability of the results beyond the specific participants is uncertain. Third, and relatedly, the fact that all participants are runners is also

suboptimal with regard to the generalisability of the results. We also notice that the sample is predominantly male (76%), which is likely due to the recruitment method via Garmin Connect—a platform that may be more appealing to male than female runners. Runners are known to be healthier than the general population—both physically and psychologically<sup>30-33</sup>—as also demonstrated by the comparison of psychological well-being between the participants in the Garmin-RUNSAFE Running Health Study and the participants from the COVID-19 Consequences Denmark Panel Survey 2020. However, while runners are not representative of the general population, the fact that they are considered to be quite robust from a psychological perspective, implies that the inverse relationship is likely to be stronger in the general population, thereby rendering our estimate a conservative one. Fourth, with regard to the exposure, namely the number of COVID-19-related deaths, there are inter-country differences in the reporting/operationalisation.<sup>34,35</sup> This does not constitute a major problem, because country differences are removed with the individual-fixed effects. Nevertheless, identical reporting practices would have been preferable. Fifth and relatedly, data on nationwide and regional lockdowns from the 86 countries were not available to us. We were therefore unable to investigate whether the observed negative relationship between COVID-19-related deaths and psychological well-being is driven by the lockdowns—a downstream consequence of pandemic pressure—as has been suggested by some, but not all, other studies.<sup>36,37</sup> Sixth, although the results of this study do not suggest that running activity and running related injuries/problems have marked effects on the impact of the COVID-19 pandemic pressure on psychological well-being, controlled intervention studies are required to clarify the question of causality. Such studies are, however, also associated with challenges—in particular due to the difficulties with regard to blinding, which is virtually impossible. Seventh, our data does not cover the period from 1 January 2021 and onwards, but based on other studies covering this period, it seems that the psychological well-being of people has kept covarying with the pandemic pressure.<sup>17,36</sup> Given that the pandemic pressure is relatively low at the time of writing, it seems reasonable to assume that its negative influence on psychological well-being is correspondingly low.

In conclusion, based on analysis of longitudinal data from 7808 runners from 86 countries, this study substantiates the notion that the COVID-19 pandemic has had a negative impact on the psychological well-being of the affected populations. As the COVID-19 pandemic is ongoing and may develop further due to occurrence of new viral variants, these findings are concerning from a global mental health perspective.

## Author affiliations

<sup>1</sup>Department of Affective Disorders, Aarhus University Hospital, Aarhus, Denmark

<sup>2</sup>Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

<sup>3</sup>Department of Political Science, Aarhus University, Aarhus, Denmark

<sup>4</sup>Centre for the Experimental-Philosophical Study of Discrimination, Aarhus University, Aarhus, Denmark

<sup>5</sup>Department of Political Science, University of Copenhagen, Copenhagen, Denmark



<sup>6</sup>Department of Political Science, University College London, London, UK

<sup>7</sup>Sport Sciences, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

<sup>8</sup>Department of Public Health, Aarhus University, Aarhus, Denmark

<sup>9</sup>Research Unit for General Practice, Aarhus University, Aarhus, Denmark

**Acknowledgements** The authors are grateful to Garmin International, Olathe, USA for assisting with recruitment of study participants and facilitating interpretation of running metrics from Garmin Connect.

**Contributors** The study was designed in collaboration between all authors. The analyses were carried out by HTV and KMS. The results were interpreted by all authors. HTV, PTD, KMS and SDØ wrote the first draft of the manuscript, which was subsequently revised for important intellectual content by RBKB and RØN. All authors approved the final version of the manuscript prior to submission. HTV is the guarantor.

**Funding** The study is supported by unconditional grants from the Novo Nordisk Foundation to SDØ (Grant number: NNF20SA0062874) and from Aarhus University Research Foundation (grant number: AUFF-E-2015-FLS-9-9) and the Danish Rheumatism Association to RØN (grant number: R160-A5157). SDØ is further supported by grants from the Lundbeck Foundation (grant numbers: R358-2020-2341 and R344-2020-1073), the Danish Cancer Society (grant number: R283-A16461), the Central Denmark Region Fund for Strengthening of Health Science (grant number: 1-36-72-4-20), the Danish Agency for Digitisation Investment Fund for New Technologies (grant number 2020-6720), Independent Research Fund Denmark (grant number: 7016-00048B).

**Disclaimer** These funders had no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

**Competing interests** SDØ received the 2020 Lundbeck Foundation Young Investigator Prize. Furthermore, SDØ owns units of mutual funds with stock tickers DKIGI and WEKAFKI, as well as units of exchange traded funds with stock tickers TRET and EUNL. The remaining authors report no conflicts of interest.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants but all participants in the Garmin-RUNSAFE Running Health Study completed an online informed consent form prior to enrolment. As this was an observational study, the local ethics committee in the Central Denmark Region waived registration (Request number: 227/2016- Record number: 1-10-72-189-16) in accordance with the Danish Act on Research Ethics Review of Health Research Projects, Section 14, no. 2. The Danish Data Protection Agency approved the study (the Danish Data Protection Agency's record number: 2015-57-0002; Aarhus University's record number: 62908, serial number 309), including the data collection procedure and data storage.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** The data used for the present study cannot be shared as the informed consent specifies that they will be stored only at servers at Aarhus University, Denmark.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Helene Tilma Vistisen <http://orcid.org/0000-0003-4929-943X>

Kim Mannemar Sønderskov <http://orcid.org/0000-0002-3550-0772>

Peter Thisted Dinesen <http://orcid.org/0000-0003-0526-9087>

René Børge Korsgaard Brund <http://orcid.org/0000-0002-7246-4811>

Rasmus Østergaard Nielsen <http://orcid.org/0000-0001-5757-1806>

Søren Dinesen Østergaard <http://orcid.org/0000-0002-8032-6208>

## REFERENCES

- Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry* 2020;7:547-60.
- WHO. Mental health and psychosocial considerations during the COVID-19 outbreak, 2020. Available: <https://www.who.int/docs/default-source/coronaviruse/mental-health-considerations.pdf>
- van Zyl LE, Rothmann S, Zondervan-Zwijnenburg MAJ. Longitudinal trajectories of study characteristics and mental health before and during the COVID-19 Lockdown. *Front Psychol* 2021;12:633533.
- Evans S, Alkan E, Bhargava JK, et al. Effects of the COVID-19 lockdown on mental health, wellbeing, sleep, and alcohol use in a UK student sample. *Psychiatry Res* 2021;298:113819.
- Kivi M, Hansson I, Bjälkebring P. Up and about: older adults' well-being during the COVID-19 pandemic in a Swedish longitudinal study. *Journals Gerontol Ser B* 2021;76:e4-9.
- Lizana PA, Vega-Fernandez G, Gomez-Bruton A, et al. Impact of the COVID-19 pandemic on teacher quality of life: a longitudinal study from before and during the health crisis. *Int J Environ Res Public Health* 2021;18:18073764. doi:10.3390/ijerph18073764
- Rantanen T, Eronen J, Kauppinen M, et al. Life-space mobility and active aging as factors underlying quality of life among older people before and during COVID-19 lockdown in Finland-a longitudinal study. *J Gerontol A Biol Sci Med Sci* 2021;76:e60-7.
- Kwong ASF, Pearson RM, Adams MJ, et al. Mental health before and during the COVID-19 pandemic in two longitudinal UK population cohorts. *Br J Psychiatry* 2021;218:334-43.
- Ramiz L, Contrand B, Rojas Castro MY, et al. A longitudinal study of mental health before and during COVID-19 lockdown in the French population. *Global Health* 2021;17:29.
- Savage MJ, Hennis PJ, Magistro D, et al. Nine months into the COVID-19 pandemic: a longitudinal study showing mental health and movement behaviours are impaired in UK students. *Int J Environ Res Public Health* 2021;18:2930.
- Ejiri M, Kawai H, Kera T, et al. Exercise as a coping strategy and its impact on the psychological well-being of Japanese community-dwelling older adults during the COVID-19 pandemic: a longitudinal study. *Psychol Sport Exerc* 2021;57:102054.
- Koppert TY, Jacobs JW, Geenen R. The psychological impact of the COVID-19 pandemic on Dutch people with and without an inflammatory rheumatic disease. *Rheumatology* 2021;60:3709-15.
- Sønderskov KM, Dinesen PT, Santini ZI, et al. The depressive state of Denmark during the COVID-19 pandemic. *Acta Neuropsychiatr* 2020;32:226-8.
- Sønderskov KM, Dinesen PT, Santini ZI, et al. Increased psychological well-being after the apex of the COVID-19 pandemic. *Acta Neuropsychiatr* 2020;32:277-9.
- Sønderskov KM, Dinesen PT, Santini ZI, et al. Increased psychological well-being after the apex of the COVID-19 pandemic. *Acta Neuropsychiatr* 2020;32:277-9.
- Sønderskov KM, Dinesen PT, Vistisen HT, et al. Variation in psychological well-being and symptoms of anxiety and depression during the COVID-19 pandemic: results from a three-wave panel survey. *Acta Neuropsychiatr* 2021;33:156-9.
- Vistisen HT, Sønderskov KM, Dinesen PT, et al. Psychological well-being and symptoms of depression and anxiety across age groups during the second wave of the COVID-19 pandemic in Denmark. *Acta Neuropsychiatr* 2021;33:331-4.
- O'Connor RC, Wetherall K, Cleare S. Mental health and well-being during the COVID-19 pandemic: longitudinal analyses of adults in the UK COVID-19 Mental Health & Wellbeing study. *Br J Psychiatry* 2020;1-8.
- Pedersen MT, Andersen TO, Clotworthy A, et al. Time trends in mental health indicators during the initial 16 months of the COVID-19 pandemic in Denmark. *BMC Psychiatry* 2022;22:25.
- Hansen BT, Sønderskov KM, Hageman I, et al. Daylight savings time transitions and the incidence rate of unipolar depressive episodes. *Epidemiology* 2017;28:346-53.
- Nielsen Rasmus Østergaard, Bertelsen ML, Ramskov D, et al. The Garmin-RUNSAFE running health study on the aetiology of running-related injuries: rationale and design of an 18-month

- prospective cohort study including runners worldwide. *BMJ Open* 2019;9:e032627.
- 22 Topp CW, Østergaard SD, Søndergaard S, *et al.* The WHO-5 well-being index: a systematic review of the literature. *Psychother Psychosom* 2015;84:167–76.
  - 23 Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* 2020;20:533–4.
  - 24 Rohde C, Jefsen OH, Nørremark B, *et al.* Psychiatric symptoms related to the COVID-19 pandemic. *Acta Neuropsychiatr* 2020;32:274–6.
  - 25 Enevoldsen KC, Danielsen AA, Rohde C, *et al.* Monitoring of COVID-19 pandemic-related psychopathology using machine learning. *Acta Neuropsychiatr* 2022;34:148–52.
  - 26 Johansson F, Côté P, Hogg-Johnson S, *et al.* Depression, anxiety and stress among Swedish university students during the second and third waves of COVID-19: a cohort study. *Scand J Public Health* 2021;49:750–4.
  - 27 Bendau A, Plag J, Kunas S, *et al.* Longitudinal changes in anxiety and psychological distress, and associated risk and protective factors during the first three months of the COVID-19 pandemic in Germany. *Brain Behav* 2021;11:e01964.
  - 28 Fancourt D, Steptoe A, Bu F. Trajectories of anxiety and depressive symptoms during enforced isolation due to COVID-19 in England: a longitudinal observational study. *Lancet Psychiatry* 2021;8:141–9.
  - 29 World Health Organization (WHO). Mental health and COVID-19. Available: <https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/publications-and-technical-guidance/noncommunicable-diseases/mental-health-and-covid-19> [Accessed 15 Mar 2022].
  - 30 White RL, Babic MJ, Parker PD, *et al.* Domain-Specific physical activity and mental health: a meta-analysis. *Am J Prev Med* 2017;52:653–66.
  - 31 Teychenne M, Ball K, Salmon J. Physical activity and likelihood of depression in adults: a review. *Prev Med* 2008;46:397–411.
  - 32 Schuch FB, Vancampfort D, Firth J, *et al.* Physical activity and incident depression: a meta-analysis of prospective cohort studies. *Am J Psychiatry* 2018;175:631–48.
  - 33 Mammen G, Faulkner G. Physical activity and the prevention of depression: a systematic review of prospective studies. *Am J Prev Med* 2013;45:649–57.
  - 34 Worldometer. Worldometer COVID-19 data. Available: <https://www.worldometers.info/coronavirus/about/> [Accessed 15 Mar 2022].
  - 35 Balmford B, Annan JD, Hargreaves JC, *et al.* Cross-Country comparisons of Covid-19: policy, politics and the price of life. *Environ Resour Econ* 2020;76:525–51.
  - 36 Joensen A, Danielsen S, Andersen PK, *et al.* The impact of the initial and second national COVID-19 lockdowns on mental health in young people with and without pre-existing depressive symptoms. *J Psychiatr Res* 2022;149:233–42.
  - 37 Prati G, Mancini AD. The psychological impact of COVID-19 pandemic lockdowns: a review and meta-analysis of longitudinal studies and natural experiments. *Psychol Med* 2021;51:201–11.