

## **Aalborg Universitet**

Detection of microorganisms involved in airway infection of cystic fibrosis patients by standard culturing and molecular methods

Rudkjøbing, Vibeke Børsholt; Nielsen, Per Halkjær; Johansen, Ulla: Høiby, Niels; Thomsen, Trine Rolighed; Bjarnsholt, Thomas

Publication date: 2010

Document Version Early version, also known as pre-print

Link to publication from Aalborg University

Citation for published version (APA):

Rudkjøbing, V. B., Nielsen, P. H., Johansen, U., Høiby, N., Thomsen, T. R., & Bjarnsholt, T. (2010). *Detection of microorganisms involved in airway infection of cystic fibrosis patients by standard culturing and molecular methods*. Abstract from Biofilm IV International Conference, 1-3 september 2010, Winchester, United Kingdom.

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
   You may freely distribute the URL identifying the publication in the public portal -

Take down policy
If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

## Biofilms4 international conference. 1-3 September, 2010. Winchester, UK

Poster: Detection of microorganisms involved in airway infection of cystic fibrosis patients by cultivation techniques and molecular methods.

Vibeke B Rudkjøbing1, Trine R Thomsen1,2, Per H Nielsen1, Morten Alhede3, Ulla R Johansen4, Niels Høiby4 and Thomas Bjarnsholt4,5

Patients suffering from the genetic disease cystic fibrosis (CF), develop chronic lung infection. This infection persists due to highly viscous mucus occurring as a result of the disease, in which bacteria form biofilms. Diagnostic tools for CF rely on culture based techniques performed on expectorated sputum samples, and most studies are centered on this sample type. It is however problematic to investigate biofilm-residing microorganisms, and further problems may occur since it is possible that the samples may be contaminated by oral flora during expectoration.

In this study tissue and sputum samples (n=24) from explanted lungs of four Danish CF patients were examined to circumvent possible oral flora contamination. Samples were examined by standard culturing techniques, including aerobic and anaerobic growth, at Rigshospitalet, Denmark. These findings were compared to results obtained by 16S rRNA gene analysis (16S rRNA gene amplification, cloning, sequencing and phylogenetic analysis) performed blinded of the growth results and quantification of the *oprL* gene of *Pseudomonas aeruginosa* by quantitative PCR at Aalborg University, Denmark.

The microorganism detected most often by 16S rRNA gene analysis in a sample was also detected by standard culturing techniques (which gave monomicrobial results). 16S rRNA gene analysis suggested that samples contained polymicrobial infection. A correspondence between the frequent detection of *P. aeruginosa* by the above methods and the number of *P. aeruginosa* cells in most samples were found by quantitative PCR. This suggests that detection of *P. aeruginosa* by culturing and 16S rRNA gene analysis is not caused by biases in the techniques.