

Nationwide study on trends in unplanned hospital attendance and deaths during the 7 weeks after the onset of the COVID-19 pandemic in Denmark

Bogh, Søren Bie; Fløjstrup, Marianne; Nissen, Søren Kabell; Hanson, Stine; Bech, Mickael; Johnsen, Søren Paaske; Kristensen, Mette Rahbek; Laugesen, Line Emilie; Søndergaard, Jens; Folkestad, Lars; Christensen, Erika Frischknecht; Henriksen, Daniel Pilsgaard; Hsia, Renee Y; Graham, Colin A; Lindskou, Tim Alex; Byg, Keld-Erik; Søvsø, Morten Breinholt; Laugesen, Henrik; Hallas, Peter; Mikkelsen, Søren; Olsen, Kim Rose; Thygesen, Lau Caspar; Gamst-Jensen, Hejdi; Brabrand, Mikkel

Published in:
BMJ Quality and Safety

DOI (link to publication from Publisher):
[10.1136/bmjqs-2020-012144](https://doi.org/10.1136/bmjqs-2020-012144)

Creative Commons License
CC BY-NC 4.0

Publication date:
2021

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Bogh, S. B., Fløjstrup, M., Nissen, S. K., Hanson, S., Bech, M., Johnsen, S. P., Kristensen, M. R., Laugesen, L. E., Søndergaard, J., Folkestad, L., Christensen, E. F., Henriksen, D. P., Hsia, R. Y., Graham, C. A., Lindskou, T. A., Byg, K.-E., Søvsø, M. B., Laugesen, H., Hallas, P., ... Brabrand, M. (2021). Nationwide study on trends in unplanned hospital attendance and deaths during the 7 weeks after the onset of the COVID-19 pandemic in Denmark. *BMJ Quality and Safety*, 30(12), 986-995. Article 012144. <https://doi.org/10.1136/bmjqs-2020-012144>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: December 04, 2025

A nationwide study on trends in unplanned hospital attendance and deaths during the seven weeks after the onset of the COVID-19 pandemic in Denmark

Word count

2939 words

ABSTRACT

Background:

The impact of a pandemic on unplanned hospital attendance has not been extensively examined before. The aim of this study is to report the nationwide consequences of the COVID-19 pandemic on unplanned hospital attendances in Denmark for seven weeks after a 'shelter at home' order was issued for the whole nation.

Methods:

We merged data from national registries (the Civil Registration System and Patient Registry) to conduct an observational study of unplanned (excluding outpatient visits and elective surgery from the patient registry) hospital-based healthcare seeking and mortality of all Danish residents. Using data for the first seven weeks after the 'shelter at home' order was issued, the incidence rate of unplanned hospital attendances per week in 2020 was compared to corresponding weeks in 2017-2019 extracted from the national patient register. The main outcome was hospital attendances per week as incidence rate ratios (IRR). Secondary outcomes were general population mortality and risk of death in-hospital, reported as weekly mortality rate ratios (MRR).

Results:

From 2,438,286 attendances in the study period, overall unplanned attendances decreased by up to 21%; attendances excluding COVID-19 were reduced by 31%; non-psychiatric attendances by 31%; and psychiatric attendances by 30%. Out of the five most common diagnoses expected to remain stable, only schizophrenia and myocardial infarction remained

stable, while COPD exacerbation, hip fracture and urinary tract infection fell significantly.

The nationwide general population MRR rose in six of the recorded weeks, while MRR excluding COVID-19 positive patients only increased in two.

Conclusion:

The COVID-19 pandemic and a governmental national 'shelter at home' order was associated with a marked reduction in unplanned hospital attendances with an increase in MRR for the general population in two of seven weeks, despite exclusion of COVID-19 patients. The findings should be taken into consideration when planning for public information campaigns during future pandemics.

Keywords:

Pandemic

Attendance

Mortality

COVID-19

Public Health

INTRODUCTION

At the end of 2020, more than 83 million people have been confirmed to have COVID-19 infection and 1.8 million have died worldwide.¹ To reduce the transmission and mortality from COVID-19 – i.e., “flatten the curve” – many countries, including Denmark (supplementary figure 1), have issued 'shelter at home' orders involving the whole nation temporarily shutting down daycare, schools, industries, and most public gatherings.² Such actions have not been taken before in modern times and the downstream effects are therefore unknown.

The effects of such drastic measures are not only confined to patients with suspected or confirmed COVID-19 infection: these actions affect the entire population. "Shut-down" measures and the intense focus on a single disease could increase public concerns about the risk of contamination by infected persons.³⁻⁵ This may cause the general public to believe that the concentration of infected persons is higher in healthcare settings than in society as a whole, and that the capacity of the health care system is not sufficient to handle non-COVID-19 patients. Consequently, people may avoid hospitals or delay seeking medical care with possible detrimental effects. Indeed, there have been reports of reduction in hospital and general practitioner (GP) visits during the COVID-19 pandemic from several countries.^{6,7} Several countries have reported increases in overall mortality during the pandemic,⁸ while some, including Denmark, until now have shown little change or a reduction of the mortality rate.⁹

To date, no scientific study has documented unplanned hospital attendances of an entire population during a pandemic. We report the nationwide consequences of the COVID-19 pandemic on unplanned hospital attendance patterns and population mortality of Denmark for

the first seven weeks after the national 'shelter at home' order (11 March 2020) was issued. Specifically, this study compares the incidence rate of unplanned non-psychiatric and psychiatric hospital attendances on a weekly basis before and during the 'shelter at home' period between 2017-2019 and 2020; it shows these patterns for the most common discharge diagnoses of diseases expected to remain stable during the pandemic; and it describes the risk of death-in-hospital and the general population mortality during these periods.

METHODS

We conducted a nationwide observational study of hospital-based healthcare seeking and mortality of all people living in Denmark, excluding Greenland. We used the Danish nationwide registers (the Danish Civil Registration System and the Danish National Patient Registry, see below) to identify all unplanned hospital attendances and calculated the incidence rate per week in 2020 (9 March-26 April), comparing it to corresponding weeks in 2017-2019.

Setting

Denmark is a high-income country with high population density (137 per km²) and approximately 5.8 million citizens. The Danish healthcare system is publicly funded by taxation and is free at the point of care to all residents with universal coverage, except for some co-payments for prescription drugs and dentistry. GPs act as gatekeepers to secondary healthcare and almost all residents are assigned a GP.¹⁰ Emergency Department (ED) attendance requires a medical referral or activation of the emergency medical services. Psychiatric hospitals have separate EDs and some are open for walk-in patients.¹¹

Participants

We included all people registered as living in Denmark between 1 January 2017 and 26 April 2020.

Data sources

Data on age, sex, and mortality were extracted from the Danish Civil Registration System (DCRS).¹² Information on hospital attendance (date of arrival and discharge), diagnoses, and procedures were extracted from the Danish National Patient Registry (DNPR),¹³ including non-psychiatric and psychiatric attendance. Both registries are considered to have high data quality and low frequency of missing data,¹² but DNPR was delivered in a temporary form, made available by the government expressly due to the COVID-19 pandemic for research. Follow-up ended on 26 April 2020. Unplanned hospital attendance were identified using the allocated variables (`c_patttype` = 1 or 3, `c_indm` = 1 and `prioritet` = ATA1) describing the indication of the priority of the patient and the patient's administrative relationship with the hospital.

DNPR is based on individual hospital attendances, and any transfer between departments was coded in the database as individual attendances. For the purpose of assessing attendance as an entire admission rather than individual attendances, we combined instances of multiple attendances within four-hours into one attendance.¹⁴ Only unplanned hospital attendances were included. We did not include elective surgery and outpatient clinic attendances. The number of inhabitants in Denmark was retrieved from Statistics Denmark that collects, compiles, and publishes statistics on the Danish society ().

Outcomes

The primary outcome was the incidence rate ratio of unplanned hospital attendances per week, calculated by comparing unplanned incidence rates during 'shelter at home' with corresponding periods from the prior three years.

Secondary outcomes were mortality rate of the general population and risk of death-in-hospital on a weekly basis.

Mortality rate for the general population was calculated for the entire population based on DCRS data, while risk of death-in-hospital was calculated for any person who died during admission or on the day of discharge based on DCRS (for date of death) and DNPR (for whether the person was admitted at that time or not) data. The yearly increase in the Danish population was taken into account by using the mean population number per period.

Variables

Age was grouped into four categories, 0-17, 18-49, 50-69, and 70+ years.

Comorbidity was presented using the Charlson Comorbidity Index (CCI).¹⁵ CCI was calculated by us based on hospital diagnoses in the ten years before the hospital attendance. CCI was coded three levels: low (score 0), moderate (score 1–2), and high (score ≥ 3).

In addition, we predefined 29 diagnoses (supplementary table 1) for which the incidence rate was expected to remain unaffected by the 'shelter at home' orders. We plotted these on a weekly basis for the years 2017-2019 (data not presented) and selected the five most common discharge diagnoses that remained stable across 2017-2019 to examine changes in 2020.

Ethics

Under Danish law, observational studies do not require ethical approval.¹⁶ The study was approved by the Danish Patient Safety Authority (file number 20/18426). Data are presented in accordance with the STROBE guidelines.¹⁷

Data availability

According to Danish law, data cannot be shared. However, the data sources are available for other researchers pending approval from the Danish Health Data Authority.

Statistics

Data are presented in absolute numbers of attendances (percent) and differences between groups assessed with the Pearson's chi-squared test. To account for any holidays that might have implications for week-to-week variance, we calculated the weekly mean number of the unplanned hospital attendances for the control years (2017-2019). To analyze the changes in death-in-hospital and mortality rate after 'shelter at home' was ordered, we calculated the incidence rate ratio and mortality rate ratio using Poisson regression. The impact model calculated the weekly incidence rate ratio and mortality rate ratio during the home shelter period (9 March-26 April 2020) with all other weeks as a reference. We adjusted for the year-by-year increase in unplanned hospital attendances and the weekly variance. We did not adjust for patient-level characteristics to describe the actual changes in the unplanned hospital attendances. However, analyses were stratified according to age, sex, psychiatric/non-psychiatric, and the five most common discharge diagnoses expected to remain unaffected during the pandemic. Analyses were done on attendance level. All analyses were carried out with Stata 15 (Stata Corp LP, College Station, TX).

RESULTS

A total of 2,438,286 unplanned hospital attendances were registered, 247,567 of these happened during the 'shelter at home' period and 763,538 in corresponding weeks in the previous three years (Table 1). The case-mix in the 'shelter at home period' differed compared to the previous years' corresponding weeks. More were female, older, and more likely to have a medium score in Charlson comorbidity index after the 'shelter at home' order.

Table 1 – Characteristics of patients from the control years in 2017-19, the prior period in 2020 (30 December 2019-8 March 2020) and during the 'shelter at home' (9 March-26 April) order was issued on 11 March 2020. The “five most stable diagnoses” designate the five most common of 29 pre-specified diagnoses we expected to remain stable during the ‘shelter at home’ period based on data from 2017 to 2019.

	2017-19		2020		P-value ^c
	Corresponding to before ‘shelter at home’	Corresponding to ‘shelter at home’	Before ‘shelter at home’	During ‘shelter at home’	
	N=1 050 805	N=763 538	N=376 376	N=247 567	
Sex					<0.001
Female	547,437 (52.1%)	395,541 (51.8%)	201,070 (53.4%)	137,194 (55.4%)	
Male	503,368 (47.9%)	367,997 (48.2%)	175,306 (46.6%)	110,373 (44.6%)	
Overall, age					<0.001
<18	247,445 (23.5%)	179,604 (23.5%)	80,062 (21.3%)	43,008 (17.4%)	
18-49	356,228 (33.9%)	263,684 (34.5%)	132,994 (35.3%)	92,072 (37.2%)	
50-69	214,652 (20.4%)	155,647 (20.4%)	77,617 (20.6%)	58,646 (23.7%)	
70+	232,480 (22.1%)	164,603 (21.6%)	85,703 (22.8%)	53,841 (21.7%)	
Charlson Comorbidity Index					<0.001
Low	691,649 (65.8%)	506,847 (66.4%)	246,049 (65.4%)	161,049 (65%)	
Medium	240,639 (22.9%)	173,122 (22.7%)	88,328 (23.5%)	59,798 (24.2%)	
High	118,515 (11.3%)	83,569 (10.9%)	41,993 (11.2%)	26,701 (10.8%)	
Five most stable diagnoses ^b					<0.001
Schizophrenia	7,151 (0.7%)	5,370 (0.7%)	3,064 (0.8%)	2,006 (0.8%)	
COPD ^a exacerbation	11,245 (1.1%)	7,392 (1.0%)	3,529 (0.9%)	1,871 (0.8%)	

Hip fracture	4,301 (0.4%)	2,902 (0.4%)	1,524 (0.4%)	870 (0.4%)	
Myocardial infarction	3,018 (0.3%)	2,022 (0.3%)	942 (0.3%)	600 (0.2%)	
Urinary tract infection	10,034 (1.0%)	7,562 (1.0%)	3,510 (0.9%)	1,728 (0.7%)	

^a COPD: Chronic obstructive pulmonary disease

^b Percentage of all patients

^c P values based on Pearson's chi-squared test representing the probability that the control period in 2017-2019 is equal to 9 March-26 April 2020

Immediately after the 'shelter at home' order was issued, there was a clear drop in unplanned hospital attendances (figure 1A). Within three weeks, however, the number of attendances rapidly increased to near normal numbers, but this was due to suspected or confirmed COVID-19 cases (figure 1A-D). The drop in non-COVID-19 attendances were seen for both non-psychiatric and psychiatric diagnoses (figure 1C and 1D). The mean number of unplanned hospital attendance in 2020 was 646 (SD 12.3) before and 607 (SD 61) after the 'shelter at home' order. In the corresponding periods in 2017-2019 the mean number of unplanned hospital attendances were 606 (SD 16.6) and 628.6 (SD 9), respectively. In relative numbers, expressed as incidence rate ratios, unplanned visits decreased significantly after the 'shelter at home' order (table 2), both overall (16-22 March, IRR 0.79 CI 95% 0.77; 0.80), excluding COVID-19 related attendances (16-22 March IRR 0.69 CI 95% 0.68; 0.71), for non-psychiatric attendances (16-29 March IRR 0.69 CI 95% 0.68; 0.71), and psychiatric attendances (6-12 April IRR 0.70 CI 95% 0.65; 0.75).




Figure 1 Number of unplanned hospital visits per 100 000 population per week. Solid lines indicate weeks in 2020, dashed lines mean number of visits per week in 2017–2019. (A) Number of unplanned hospital visits overall. (B) Number of unplanned hospital visits excluding COVID-19-related visits. (C) Number of unplanned non-psychiatric hospital visits excluding COVID-19-related visits. (D) Number of unplanned psychiatric hospital visits excluding COVID-19 related visits.

Children's (< 18 years old) unplanned attendances were the age group most affected with a 41 % reduction from 30 March-5 April (IRR 0.59 CI 95% 0.57; 0.61). However, this drop for children was not as pronounced for unplanned psychiatric attendances with reduced use in a single week (6-12 April) immediately followed by an increased use the following week.

Of the five diagnoses expected to remain stable, COPD exacerbation was the most affected with an unexpected decline in all weeks. Also, hip fracture and urinary tract infections had unexpected drops in unplanned attendances with a 32% (IRR 0.68 CI 95% 0.53; 0.88) and 54% (IRR 0.46 CI 95% 0.47; 0.67) drop 23-29 April, respectively.

Table 2 – Incidence rate ratios (IRR) for unplanned hospital visits in the period following the 'shelter at home' order in Denmark. The incidence rate of attendances in the control period from 2017-2019 was used as reference. Data are excluding COVID-19 related attendances unless otherwise specified. The “five most stable diagnoses” designate the five of 29 pre-specified diagnoses we expected to remain stable during a pandemic.

	9-15 March	16-22 March	23-29 March	30 March- 5 April	6-12 April	13-19 April	20-26 April
	IRR ^a (95% CI, p-value)	IRR ^a (95% CI, p-value)	IRR ^a (95% CI, p-value)	IRR ^a (95% CI, p-value)	IRR ^a (95% CI, p-value)	IRR ^a (95% CI, p-value)	IRR ^a (95% CI, p-value)
Overall – including COVID ^b 19 related attendances	0.86 (0.84;0.87, <0.001)	0.79 (0.77;0.80, <0.001)	0.85 (0.83;0.86, <0.001)	1.01 (0.99;1.02, 0.30)	0.86 (0.85;0.88, <0.001)	0.99 (0.97;1.00, 0.08)	0.99 (0.98;1.01, 0.29)
Overall – excluding COVID 19 related attendances	0.82 (0.80;0.83, <0.001)	0.69 (0.68;0.71, <0.001)	0.70 (0.68;0.71, <0.001)	0.71 (0.70;0.72, <0.001)	0.73 (0.72;0.74, <0.001)	0.84 (0.83;0.85, <0.001)	0.81 (0.80;0.82, <0.001)
Non-psychiatric – excluding COVID 19 related attendances	0.81 (0.80;0.82, <0.001)	0.69 (0.68;0.70, <0.001)	0.69 (0.68;0.70, <0.001)	0.70 (0.69;0.72, <0.001)	0.73 (0.72;0.74, <0.001)	0.83 (0.82;0.85, <0.001)	0.80 (0.79;0.81, <0.001)
Psychiatric – excluding COVID 19 related attendances	0.93 (0.86;0.99, 0.03)	0.79 (0.73;0.85, <0.001)	0.83 (0.77;0.89, <0.001)	0.82 (0.76;0.88, <0.001)	0.70 (0.65;0.75, <0.001)	1.01 (0.94;1.09, 0.70)	0.94 (0.88;1.01, 0.09)
Overall, age							
<18	0.80 (0.78;0.83, <0.001)	0.65 (0.62;0.67, <0.001)	0.61 (0.59;0.64, <0.001)	0.59 (0.57;0.61, <0.001)	0.69 (0.66;0.71, <0.001)	0.74 (0.71;0.76, <0.001)	0.64 (0.62;0.66, <0.001)
18-49	0.80 (0.78;0.82, <0.001)	0.68 (0.66;0.70, <0.001)	0.67 (0.65;0.69, <0.001)	0.70 (0.68;0.72, <0.001)	0.73 (0.71;0.75, <0.001)	0.81 (0.78;0.83, <0.001)	0.80 (0.78;0.82, <0.001)
50-69	0.86 (0.83;0.89, <0.001)	0.75 (0.72;0.77, <0.001)	0.76 (0.73;0.78, <0.001)	0.78 (0.75;0.80, <0.001)	0.75 (0.72;0.78, <0.001)	0.91 (0.88;0.94, <0.001)	0.89 (0.86;0.92, <0.001)
70+	0.81 (0.79;0.84, <0.001)	0.72 (0.69;0.74, <0.001)	0.76 (0.73;0.79, <0.001)	0.78 (0.76;0.81, <0.001)	0.75 (0.72;0.77, 0.001)	0.94 (0.91;0.98, <0.001)	0.90 (0.87;0.93, <0.001)
Non-psychiatric, age							
<18	0.79 (0.76;0.82, <0.001)	0.64 (0.61;0.66, <0.001)	0.60 (0.58;0.63, <0.001)	0.58 (0.56;0.61, <0.001)	0.69 (0.67;0.71, <0.001)	0.72 (0.70;0.75, <0.001)	0.62 (0.60;0.65, <0.001)
18-49	0.79 (0.77;0.81, <0.001)	0.68 (0.66;0.70, <0.001)	0.66 (0.64;0.67, <0.001)	0.69 (0.67;0.71, <0.001)	0.73 (0.71;0.75, <0.001)	0.79 (0.77;0.81, <0.001)	0.80 (0.77;0.82, <0.001)
49-69	0.85 (0.82;0.89, <0.001)	0.74 (0.71;0.77, <0.001)	0.75 (0.73;0.78, <0.001)	0.77 (0.74;0.80, <0.001)	0.75 (0.72;0.78, <0.001)	0.91 (0.87;0.94, <0.001)	0.88 (0.85;0.91, <0.001)
70+	0.81 (0.79;0.84, <0.001)	0.72 (0.69;0.74, <0.001)	0.76 (0.73;0.79, <0.001)	0.78 (0.75;0.81, <0.001)	0.75 (0.72;0.77, 0.001)	0.94 (0.91;0.98, <0.001)	0.90 (0.87;0.93, <0.001)
Psychiatric, age							

<18	1.06 (0.86;1.31, 0.56)	0.95 (0.76;1.18, 0.63)	0.89 (0.71;1.11, 0.31)	0.91 (0.73;1.15, 0.43)	0.65 (0.50;0.85, 0.001)	1.55 (1.20;2.01, 0.001)	1.03 (0.83;1.27, 0.79)
18-49	0.91 (0.83;0.99, 0.04)	0.72 (0.66;0.79, <0.001)	0.80 (0.73;0.88, <0.001)	0.76 (0.69;0.84, <0.001)	0.72 (0.65;0.79, <0.001)	0.99 (0.90;1.09, 0.91)	0.89 (0.82;0.98, 0.02)
50-69	0.91 (0.78;1.06, 0.20)	0.84 (0.72;0.98, 0.03)	0.84 (0.72;0.99, 0.04)	0.88 (0.75;1.02, 0.10)	0.71 (0.60;0.83, <0.001)	0.94 (0.81;1.10, 0.45)	1.00 (0.86;1.16, 0.99)
70+	0.77 (0.58;1.02, 0.07)	0.86 (0.65;1.13, 0.28)	0.82 (0.62;1.08, 0.15)	0.96 (0.73;1.27, 0.79)	0.68 (0.52;0.90, 0.006)	0.92 (0.70;1.21, 0.54)	0.92 (0.70;1.20, 0.53)
Risk of death in-hospital excluding Covid-19 related attendances for the 5 most common diagnosed expected to be stable without Covid-19 pandemic							
Schizophrenia	0.89 (0.75;1.06, 0.21)	0.86 (0.72;1.02, 0.08)	0.91 (0.76;1.90, 0.33)	0.89 (0.74;1.06, 0.19)	0.69 (0.57;0.83, <0.001)	0.98 (0.82;1.17, 0.85)	0.88 (0.74;1.04, 0.15)
COPD ^c exacerbation	0.77 (0.66;0.89, 0.001)	0.74 (0.63;0.87, <0.001)	0.70 (0.59;0.82, <0.001)	0.77 (0.66;0.90, <0.001)	0.68 (0.57;0.80, <0.001)	0.87 (0.74;1.02, 0.09)	0.80 (0.68;0.94, 0.006)
Hip fracture	0.85 (0.66;1.09, 0.20)	0.74 (0.58;0.97, 0.03)	0.68 (0.53;0.88, 0.003)	0.84 (0.65;1.09, 0.19)	0.84 (0.66;1.09, 0.19)	1.00 (0.77;1.28, 0.99)	1.01 (0.79;1.30, 0.91)
Myocardial Infarction	0.70 (0.51;0.96, 0.03)	0.92 (0.68;1.24, 0.57)	1.06 (0.77;1.44, 0.74)	1.13 (0.86;1.53, 0.40)	0.85 (0.62;1.18, 0.33)	1.12 (0.82;1.51, 0.44)	0.91 (0.68;1.22, 0.52)
Urinary tract infection	0.71 (0.59;0.85, <0.001)	0.58 (0.48;0.70, <0.001)	0.56 (0.47;0.67, <0.001)	0.55 (0.46;0.65, <0.001)	0.89 (0.76;1.03, 0.12)	0.58 (0.49;0.68, <0.001)	0.71 (0.60;0.84, <0.001)

^aIRR = Incidence rate ratio

^bCOVID = SARS-CoV2 virus disease

^cCOPD = Chronic obstructive pulmonary disease

There was an immediate increase in the number of hospital attendances (figure 1A-D) after the Danish National Board of Health issued a statement reminding the public to seek help if needed on 13 April.

The overall crude mortality rate in the general population decreased in the observed weeks of 2020 (figure 2). However, in relative terms, (adjusting for years, individual weeks, and weeks after issuing the 'shelter at home' order), we observed increased mortality rate ratio after the 'shelter at home' order in the general population for all weeks except week 11 (Table 3). When excluding suspected or confirmed COVID-19 infections, the increase in general population

mortality rate ratio was only significantly increased for two weeks, from 30 March-5 April (MRR 1.10 CI 95% 1.01; 1.20) and 13-19 April (MRR 1.10 CI 95% 1.01; 1.21) (Table 3).




Figure 2 Crude death rate (population and in-hospital) per 100 000 population per week in 2020. Solid lines indicate weeks in 2020 and dashed lines mean number of visits per week in 2017–2019.

Table 3 – Mortality rate ratios (MRR) for unplanned hospital attendances in the weeks following the 'shelter at home' order in Denmark. The incidence rate of attendances in the control period from 2017-2019 was used as reference.

The “five most stable diagnoses” designate the five of 29 pre-specified diagnoses we expected to remain stable during a pandemic.

NA signifies numbers that could not be calculated due to too few events. CI = confidence interval.

	9-15 March	16-22 March	23-29 March	30 March-5 April	6-12 April	13-19 April	20-26 April
	MRR ^a (95% CI, P-value)	MRR ^a (95% CI, P-value)	MRR ^a (95% CI, P-value)	MRR ^a (95% CI, P-value)	MRR ^a (95% CI, P-value)	MRR ^a (95% CI, P-value)	MRR ^a (95% CI, P-value)
Population mortality – including COVID ^b 19 related attendances	1.04 (0.96;1.14, 0.35)	1.09 (1.00;1.19, 0.046)	1.11 (1.02;1.21, 0.02)	1.20 (1.10;1.31, <0.001)	1.10 (1.00;1.20, 0.04)	1.17 (1.07;1.27, 0.001)	1.15 (1.05;1.26, 0.003)

Population mortality – excluding COVID 19 related attendances	1.04 (0.95;1.13, 0.38)	1.06 (0.98;1.16, 0.16)	1.05 (0.96;1.15, 0.28)	1.10 (1.01;1.20, 0.03)	1.01 (0.92;1.11, 0.80)	1.10 (1.01;1.21, 0.04)	1.08 (0.98;1.18, 0.11)
Risk of death-in-hospital – including COVID ^b 19 related attendances	0.96 (0.83;1.11, 0.59)	0.91 (0.79;1.06, 0.22)	0.91 (0.78;1.06, 0.24)	1.03 (0.89;1.20, 0.68)	1.03 (0.89;1.20, 0.66)	1.02 (0.88;1.19, 0.79)	1.01 (0.87;1.18), 0.87
Risk of death-in-hospital – excluding COVID 19 related attendances	0.96 (0.83;1.10, 0.54)	0.85 (0.73;0.99, 0.04)	0.79 (0.67;0.93, 0.004)	0.86 (0.73;1.01, 0.06)	0.86 (0.74;1.01, 0.07)	0.92 (0.79;1.08, 0.31)	0.90 (0.77;1.06, 0.20)
Risk of death-in-non-psychiatric-hospital	0.96 (0.83;1.11, 0.58)	0.86 (0.74;1.00, 0.04)	0.79 (0.67;0.93, 0.004)	0.86 (0.73;1.00, 0.06)	0.87 (0.74;1.01, 0.07)	0.92 (0.79;1.08, 0.30)	0.90 (0.77;1.06, 0.20)
Population mortality by age							
<18	1.18 (0.52;2.69, 0.69)	0.84 (0.30;2.36, 0.74)	0.94 (0.33;2.68, 0.91)	0.19 (0.02;1.45, 0.11)	0.55 (0.19;1.65, 0.29)	1.05 (0.44;2.47, 0.19)	0.57 (0.19;1.74, 0.33)
18-49	0.92 (0.61;1.42, 0.74)	0.88 (0.58;1.35, 0.57)	0.58 (0.37;0.90, 0.02)	1.03 (0.69;1.56, 0.88)	0.63 (0.40;0.98, 0.04)	1.38 (0.95;2.02, 0.09)	1.00 (0.66;1.53, 0.98)
49-69	1.15 (0.98;1.36, 0.08)	1.00 (0.85;1.17, 0.98)	0.91 (0.76;1.09, 0.29)	1.04 (0.88;1.23, 0.63)	1.06 (0.90;1.26, 0.49)	1.02 (0.87;1.21, 0.77)	0.99 (0.83;1.18, 0.92)
70+	1.02 (0.94;1.10, 0.67)	1.12 (1.03;1.22, 0.004)	1.18 (1.10;1.29, <0.001)	1.26 (1.16;1.36, <0.001)	1.14 (1.05;1.24, 0.002)	1.20 (1.10;1.30, <0.001)	1.20 (1.11;1.31, <0.001)
Risk of death in-hospital excluding Covid-19 related attendances for the 5 most common diagnosed expected to be stable without Covid-19 pandemic							
Schizophrenia	NA	NA	NA	NA	NA	NA	NA
COPD ^c exacerbation	1.06 (0.47;2.45, 0.88)	0.49 (0.19;1.26, 0.14)	0.68 (0.23;2.04, 0.49)	0.63 (0.23;1.70, 0.36)	0.36 (0.12;1.12, 0.08)	1.47 (0.59;3.68, 0.41)	0.49 (0.15;1.59, 0.24)
Hip fracture	1.08 (0.28;4.15, 0.90)	0.73 (0.17;3.10, 0.67)	1.45 (0.13;16.7, 0.77)	0.75 (0.10;5.40, 0.76)	1.09 (0.09;12.50, 0.95)	0.54 (0.29;5.83, 0.50)	1.31 (0.29;5.84, 0.73)
Myocardial Infarction	1.13 (0.27;4.60)	NA	1.35 (0.21;8.72)	0.33 (0.04;3.2)	4.53 (0.81;25.20)	2.41 (0.50;11.73)	0.81 (0.18;3.72)
Urinary tract infection	NA	NA	NA	NA	NA	NA	NA

^aMRR = Mortality rate ratio

^bCOVID = SARS-CoV2 virus disease

^cCOPD = Chronic obstructive pulmonary disease

The crude risk of death-in-hospital per week decreased in both 2020 and 2017-2019 (figure 2). The mean number of death-in-hospital in 2020 was 6.7 (SD 0.5) before and 6.2 (SD 0.4) per 100,000 after the 'shelter at home'. In the corresponding periods in 2017-2019 were mean number 7 (SD 0.5) and 6.6 (SD 0.5). Adjusting the risk of death-in-hospital as mortality rate ratio only showed a significantly reduced risk in the two weeks from 16-29 March (MRR 0.85 CI 95% 0.73; 0.99, MRR 0.79 CI 95% 0.67; 0.93, respectively), and only when excluding patients not hospitalized with suspected or confirmed COVID-19 infection. The reduction in the risk of death-in-hospital was mostly accounted for by the elderly (Table 3).

Out of the 29 pre-specified diagnoses expected to remain stable during a pandemic (supplementary table 1), the five most common were schizophrenia, COPD exacerbation, hip fracture, myocardial infarction, and urinary tract infection. Of these, schizophrenia, and myocardial infarction were unchanged, while the other three decreased over the seven weeks (supplementary figure 2 and table 2). The mortality rate ratio could not be calculated for schizophrenia and urinary tract infection due to too few events and was unchanged for the other three (table 3).

DISCUSSION

In this nationwide observational study, we describe a marked decrease in the incidence of unplanned hospital attendances, both for non-psychiatric and psychiatric patients, after a government-issued shelter at home order. We found a decrease of up to 31% for non-psychiatric patients and up to 30% for psychiatric patients after excluding visits due to suspected or confirmed COVID-19 infection. Crude mortality rates for the general Danish population decreased after the shelter at home order was issued, whereas the adjusted

mortality rate was moderately increased in some but not all weeks. Death-in-hospital also decreased after issuing the shelter at home order in crude analysis but adjusting for expected mortality rates as mortality rate ratios we found the decrease to only be significant if excluding patients with confirmed or suspected COVID-19 infection.

Our observed reduction in patients with non-COVID-19 related disease is similar to what has been reported elsewhere. However, the general population's overall mortality rate appears to be less marked than in many other countries.¹

When designing the study, our initial concern was, that patients would opt out of seeking healthcare during the pandemic due to the risk of nosocomial infection and a wish not to burden an already overstretched system. While choosing to opt out of healthcare perhaps can be of little consequence for some, other patients could put themselves in danger by doing so. In order to plan for future pandemics, decision-makers in healthcare must have an indication of the behavior of the public. Who will, and who will not, attend hospitals and primary care and why? Our findings show that, fortunately, mortality rates did not increase substantially despite a general decline in hospital attendance. Thus, perhaps, we should not be worried about patients with relevant complaints not seeking care during a full-scale pandemic in a country with free universal healthcare.

During this pandemic, other studies have also reported a reduction in hospital attendances by patients not suffering from COVID-19,¹⁸ but with a significant concomitant increase in mortality.^{7 19} Using the unique Danish registries, we have been able to examine the effects of the COVID-19 pandemic for an entire nation and all unplanned hospital attendances. We show a marked decrease in unplanned hospital attendances after the 'shelter at home' order

was issued, and demonstrate an 11-20% increase in the general population's mortality rate ratio in six of seven weeks. Excluding patients admitted with suspected or confirmed COVID-19, the excess mortality rate was lowered to 10% and only present in two out of seven weeks and with confidence intervals very close to 1.00.

In contrast to the findings of this study, other countries have reported reductions in stroke patients,²⁰ myocardial infarctions,²¹ and primary percutaneous coronary intervention.²² While not all of these entities were part of our analyses, we could not confirm the reduction in myocardial infarction for any of the weeks included. There are likely to be several reasons for this difference. Perhaps most importantly, Denmark was quick to issue a 'shelter at home' order; as a result, the healthcare system was never overrun, and vital emergency services were continuously provided. In addition, several widespread public information campaigns have taken place over recent years (even as late as early 2020), reminding the public to remember to seek help in case of symptoms relevant to stroke or myocardial infarction. It is also important to note that Denmark has a universal healthcare system, which decreases the barriers to seeking care. The Danish population has a high degree of health literacy and a general trust in authorities (both healthcare and governmental) and may have a stronger tendency to seek help when needed.²³ This is also evident by the rapid increase in hospital visits after the National Board of Health's reminder. While we expected COPD to remain stable in attendances based on 2017-2019 numbers, we, in fact, saw a marked reduction. We can only speculate to the cause, but suspect that a general reduction in transmittable diseases (including viral pathogens) also controlled by the national lockdown, probably decreased proportionally.

A situation, where an entire population could choose to opt out of healthcare services, has never been seen before in the Western world.²⁴ In fact, increased healthcare utilization, both in general and in relation to influenza, was seen in prior influenza epidemics.²⁵ During the recent Ebola epidemic in Africa, one study reported a decrease in hospital use in those most affected by Ebola.²⁶

While we did observe indications of increased general population mortality rates, this did not seem to be directly correlated with the steep decrease in unplanned hospital attendances, especially when excluding suspected or confirmed COVID-19 infections. As we only have data until 26 April 2020, it could be that we simply have not yet seen the full effect of the lower hospital attendance. An alternative explanation may be potential overuse of some healthcare services prior to the pandemic.²⁷ While previous studies reporting on hospital strikes have indicated a reduced in-hospital mortality,²⁸ no solid causal arguments yet have been made on adverse effects of lower hospital attendance. Denmark has a strong tradition for GP service and all Danes have access to a GP around the clock. Therefore, it is also plausible that patients could have contacted their GP (when they would normally seek hospital care), but as we do not have access to data on GP usage, we can only speculate.

Our study has limitations. First, we only have data until 26 April 2020. Therefore, any increase in delayed mortality or hospitalization will not be apparent in our study and additional follow-up time will be required. Second, while data were extracted from national registries, the patient registry was only delivered in a temporary form, made available by the government expressly due to the COVID-19 pandemic for research. Hence, the data was not validated to the usual level of the Danish databases, and there is a risk of incomplete data.²⁹ However (see supplementary figure 3), we have no indication of any reduction in data quality.

As most data in the national registers are automatically extracted from hospital registries, we expect that data quality should not be affected. Indeed, our thorough data validation did not provide any indication that this should be the case. Third, as we use only primary discharge diagnoses and not supplementary diagnoses, we rely on the treating physicians' coding quality. Given that we have compared 2017-2019 with 2020, there is most likely only a minimal risk of coding practice changing over time. Fourth, we only include data for attendances and one patient might have been included multiple times, even with related readmissions. Fifth, there is the possibility that the 'shelter at home' order resulted in a general reduction in communicable diseases which could reduce other admissions. Unfortunately, we are currently unable to study this with the existing dataset. While we only have register data, we do indeed have data on a truly national level and not only for hospital use. Our results as such do represent the general population and not only for residents using the healthcare system. And last, our results might not be generalizable as the Danish healthcare system is unique and the health-literacy of the population is high.

CONCLUSION

The COVID-19 pandemic and the governmental shelter at home order in Denmark led to a marked reduction in the incidence rate of unplanned hospital attendances with a significant decrease in the absolute number of fatalities in the general population but with a moderate increase in adjusted mortality rate in four of seven weeks in the general population.

STATEMENTS

Funding source

This study was not funded by external sources.

Conflicts of interests

All authors have no conflicts of interests to declare.

REFERENCES

1. World Health Organization. Coronavirus disease (COVID-2019) situation reports [Internet]. WHO; 2020 [Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports> accessed 16 July 2020.
2. Vrangbæk KP, Kjeld Møller. Denmark's Response to the Coronavirus Pandemic: Cambridge CoreCambridge Core blog; 2020 3 April [Available from: <https://www.cambridge.org/core/blog/2020/04/03/denmarks-response-to-the-coronavirus-pandemic/> accessed 15 May 2020.
3. Thornton J. Covid-19: A&E visits in England fall by 25% in week after lockdown. *BMJ* 2020 369:m1401.
4. Lazzerini M, Barbi E, Apicella A, et al. Delayed access or provision of care in Italy resulting from fear of COVID-19. *Lancet Child Adolesc Health* 2020 4(5):e10-e11.
5. Morning Consult. Public poll: Emergency care concerns amidst COVID-19: American College of Emergency Physicians; 2020 Apr [Available from: <https://www.emergencyphysicians.org/globalassets/emphysicians/all-pdfs/acep-mc-covid19-april-poll-analysis.pdf> accessed 15 May 2020.
6. Krumholz HM. Where Have All the Heart Attacks Gone? [Internet]. The New York Times; 2020 6 April [Available from: <https://www.nytimes.com/2020/04/06/well/live/coronavirus-doctors-hospitals-emergency-care-heart-attack-stroke.html> accessed 12 May 2020.
7. Appleby J. What is happening to non-covid deaths? *BMJ* 2020;369:m1607.
8. Baldi E, Sechi GM, Mare C, et al. Out-of-Hospital Cardiac Arrest during the Covid-19 Outbreak in Italy. *N Engl J Med* 2020;383(5):496-98.
9. Euromomo. Graphs and maps [Internet]. Euromomo; 2020 [Available from: <https://www.euromomo.eu/graphs-and-maps> accessed 7 May 2020.
10. Olejaz M, Nielsen AJ, Rudkjøbing A, et al. Denmark: Health system review [Internet]. Health Systems in Transition; 2012;14(2):1-192 [Available from: http://www.euro.who.int/__data/assets/pdf_file/0004/160519/e96442.pdf?ua=1 accessed 15 May 2020.
11. Fløjstrup M, Bogh SB, Henriksen DP, et al. Increasing emergency hospital activity in Denmark, 2005–2016: a nationwide descriptive study. *BMJ open* 2020;10(2):e031409.
12. Schmidt M, Pedersen L, Sørensen HT. The Danish Civil Registration System as a tool in epidemiology. *Eur J Epidemiol* 2014 29(8):541-49.

13. Schmidt M, Schmidt SAJ, Sandegaard JL, et al. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol* 2015;7:449-90.
14. Gubbels S, Nielsen KS, Sandegaard J, et al. The development and use of a new methodology to reconstruct courses of admission and ambulatory care based on the Danish National Patient Registry. *Int J Med Inform* 2016;95:49-59.
15. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40(5):373-83.
16. Thygesen LC, Daasnes C, Thaulow I, et al. Introduction to Danish (nationwide) registers on health and social issues: structure, access, legislation, and archiving. *Scand J Public Health* 2011;39(Suppl 7):12-16.
17. Von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370(9596):1453-57.
18. Nelson LM, Simard JF, Oluyomi A, et al. US Public Concerns About the COVID-19 Pandemic From Results of a Survey Given via Social Media. *JAMA Intern Med* 2020;180(7):1020-22.
19. Rosenbaum L. The Untold Toll—The Pandemic's Effects on Patients without Covid-19. *N Engl J Med* 2020;382(24):2368-71.
20. Kansagra AP, Goyal MS, Hamilton S, et al. Collateral Effect of Covid-19 on Stroke Evaluation in the United States. *N Engl J Med* 2020;383(4):400-01.
21. Garcia S, Albaghdadi MS, Meraj PM, et al. Reduction in ST-segment elevation cardiac catheterization laboratory activations in the United States during COVID-19 pandemic. *J Am Coll Cardiol* 2020;75(22):2871-72.
22. Piccolo R, Bruzzese D, Mauro C, et al. Population Trends in Rates of Percutaneous Coronary Revascularization for Acute Coronary Syndromes Associated with the COVID-19 Outbreak. *Circulation* 2020;141(24):2035-37.
23. Bo A, Friis K, Osborne RH, et al. National indicators of health literacy: ability to understand health information and to engage actively with healthcare providers-a population-based survey among Danish adults. *BMC Public Health* 2014;14:1095.
24. Feral-Pierssens A-L, Claret P-G, Chouihed T. Collateral damage of the COVID-19 outbreak: expression of concern. *Eur J Emerg Med* 2020;27(4):233-34.
25. McDonnell WM, Nelson DS, Schunk JE. Should we fear "flu fear" itself? Effects of H1N1 influenza fear on ED use. *Am J Emerg Med* 2012;30(2):275-82.

26. Elston J, Moosa A, Moses F, et al. Impact of the Ebola outbreak on health systems and population health in Sierra Leone. *J Public Health (Oxf)* 2016;38(4):673-78.
27. Berwick DM. Avoiding overuse—the next quality frontier. *Lancet* 2017 390(10090):102-04.
28. Cunningham SA, Mitchell K, Narayan KV, et al. Doctors' strikes and mortality: a review. *Soc Sci Med* 2008;67(11):1784-88.
29. The Danish Health Data Authority. LPR3 datakvalitetsrapport. April 12 ed, 2019:1-49.

FIGURE LEGENDS

Figure 1 – Number of unplanned hospital visits per 100 000 population per week. Solid lines indicate weeks in 2020, dashed lines mean number of visits per week in 2017-2019. A – Number of unplanned hospital visits overall. B – Number of unplanned hospital visits excluding COVID-19 related visits. C – Number of unplanned non-psychiatric hospital visits excluding COVID-19 related visits. D – Number of unplanned psychiatric hospital visits excluding COVID-19 related visits.

Figure 2 – Crude death rate (population and in-hospital) per 100 000 population per week in 2020. Solid lines indicate weeks in 2020, dashed lines mean number of visits per week in 2017-2019.