Aalborg Universitet



Multifocal tDCS targeting the motor network modulates event-related cortical responses during prolonged pain

Gregoret, Luisina; Zamorano, Anna M; Graven-Nielsen, Thomas

Published in: The Journal of Pain

DOI (link to publication from Publisher): 10.1016/j.jpain.2022.09.010

Creative Commons License CC BY 4.0

Publication date: 2023

Document Version Publisher's PDF, also known as Version of record

Link to publication from Aalborg University

Citation for published version (APA): Gregoret, L., Zamorano, A. M., & Graven-Nielsen, T. (2023). Multifocal tDCS targeting the motor network modulates event-related cortical responses during prolonged pain. *The Journal of Pain*, *24*(2), 226-236. https://doi.org/10.1016/j.jpain.2022.09.010

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.





Original Report

Multifocal tDCS Targeting the Motor Network Modulates Event-Related Cortical Responses During Prolonged Pain



Luisina Gregoret, *,[†] Anna M. Zamorano, * and Thomas Graven-Nielsen *

^{*} Center for Neuroplasticity and Pain (CNAP), Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark, [†]iMotions A/S, Copenhagen, Denmark

Abstract: Multifocal transcranial direct current stimulation (tDCS) targeting several brain regions is promising for inducing cortical plasticity. It remains unknown whether multifocal tDCS aimed at the resting-state motor network (network-tDCS) can revert N2-P2 cortical responses otherwise attenuated during prolonged experimental pain. Thirty-eight healthy subjects participated in 2 sessions separated by 24 hours (Day1, Day2) of active (n = 19) or sham (n = 19) network-tDCS. Experimental pain induced by topical capsaicin was maintained for 24 hours and assessed using a numerical rating scale. Electrical detection and pain thresholds, and N2-P2 evoked potentials (electroencephalography) to noxious electrical stimulation were recorded before capsaicin-induced pain (Day1-baseline), after capsaicin application (Day1-post-cap), and after 2 sessions of network-tDCS (Day2). Capsaicin induced moderate pain at Day1post-cap, which further increased at Day2 in both groups (P = .01). Electrical detection/pain thresholds did not change over time. N2-P2 responses were reduced on Day1-post-cap compared to Day1-baseline (P = .019). At Day2 compared with Day1-post-cap, N2-P2 responses were significantly higher in the Active network-tDCS group (P < .05), while the sham group remained inhibited. These results suggest that tDCS targeting regions associated with the motor network may modulate the late evoked brain responses to noxious peripheral stimulation otherwise initially inhibited by capsaicin-induced pain.

Perspective: This study extends the evidence of N2-P2 reduction due to capsaicin-induced pain from 30 minutes to 24 hrs. Moreover, 2 sessions of tDCS targeting the motor network in the early stage of nociceptive pain may revert the inhibition of N2-P2 associated with capsaicin-induced pain.

© 2022 The Author(s). Published by Elsevier Inc. on behalf of United States Association for the Study of Pain, Inc. This is an open access article under the CC BY license

(http://creativecommons.org/licenses/by/4.0/)

Key words: tDCS, multifocal tDCS, motor network, non-invasive brain stimulation, event-related evoked potentials, event-related potentials, experimental pain, capsaicin.

The authors declare no conflict of interest.

Pain is an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage.⁵⁷ At the cortical level, pain triggers a dynamic interaction between widespread brain networks that include regions such as somatosensory, operculoinsular, and cingulate cortices.^{4,14,36,68} Maladaptive pain-related neuroplasticity in those regions may contribute to the development of clinical pain conditions.^{3,5,14,36,61} In this context, 1 way to probe the pathways also involved in pain^{10,41} is the N2-P2 components of event-related potentials (ERPs). They represent an electrophysiological outcome of the cortical integration, cognitive evaluation, and modulation of salient stimulation, and, when

Received May 11, 2022; Revised September 8, 2022; Accepted September 13, 2022.

Center for Neuroplasticity and Pain (CNAP) is supported by the Danish National Research Foundation (DNRF121). This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 754465.

Address reprint requests to Thomas Graven-Nielsen, PhD, Center for Neuroplasticity and Pain (CNAP), Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Fredrik Bajers Vej 7D-3, 9220 Aalborg El, Denmark. E-mail: tgn@hst.aau.dk 1526-5900/\$36.00

^{© 2022} The Author(s). Published by Elsevier Inc. on behalf of United States Association for the Study of Pain, Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/) https://doi.org/10.1016/j.jpain.2022.09.010

noxious stimulation is delivered, they have been reported to correlate with pain perception^{12,38,65} and its associated saliency.⁴⁹ Indeed, pain is an intrinsically salient experience.³⁵

Modulating such processing and potentially antagonizing pain symptoms may be done by non-invasive brain stimulation (NIBS), eg transcranial direct current stimulation over the motor cortex (M1 tDCS), which is a feasible, tolerable, patient-friendly, and safe⁵³ treatment option.⁵² An overall rationale for using M1 tDCS to modulate the effects of pain is to restore the balance of the endogenous inhibitory pain pathways and to prevent or revert the maladaptive plasticity associated to persistent pain²³ through the modulation of the cortical and subcortical activity of the thalamus, anterior cingulate, and prefrontal cortices.⁵¹ However, low effect sizes and inconclusive findings in experimental⁵⁰ and clinical settings³⁹ of traditional and high definition M1 tDCS have motivated the use of new methodologies examining the stimulation of functionally associated areas through multifocal tDCS^{21,28} using a number of smallsized electrodes in order to facilitate cortical and corticospinal responses.^{11,20,27,28,46} Those studies report higher effect sizes than traditional tDCS montages. In fact, even though some tDCS studies targeting solely the M1 have shown an effect, ³⁹ M1 does not function in isolation; it interacts within the motor network.^{25,32} Stimulation of more regions of this network have resulted in increased corticospinal excitability and aftereffects persisted for a longer period compared with traditional M1 tDCS in healthy individuals.²⁰ Therefore, if there is a certain effect of M1 tDCS on the maladaptive plasticity associated to persistent pain²³ and pain perception processes,³⁹ it is likely that to stimulate all the network could increase its restorative effects.

In a healthy population, anodal tDCS of the left primary motor cortex facilitates early sensory ERPs.⁴⁷ Conflicting evidence on late ERPs after NIBS has been reported. Whereas anodal M1 tDCS was found to increase the event-related N2-P2 responses compared to baseline,²⁶ other studies show that anodal M1 tDCS¹³ and M1 repetitive transcranial magnetic stimulation⁹ did not modulate these potentials in healthy individuals. Moreover, NIBS-induced modulation has been speculated being state-specific, promoting antagonizing effects in sensitized pain pathways⁴⁰ rather than in pain-free non-sensitized subjects.⁹

Experimental models of prolonged pain, in contrast to clinical pain, allow measures before pain induction. Indeed, experimental pain models often induce relevant clinical characteristics such as dysfunctional pain mechanisms and cortical responses. For example, prolonged topical application of capsaicin produces ongoing pain, long-lasting hyperalgesia,^{7,33} reduced corticomotor excitability,^{19,20} impaired conditioned pain modulation (CPM),⁶ and reduced N2-P2 ERPs although without affecting latencies.^{64,67} At present, no studies have tried to normalize the reduced cortical ERPs during prolonged pain through sessions of tDCS.

The present exploratory study aimed at investigating whether the effects of experimental pain during

24 hours on behavior and its associated cortical processing may be reverted by 2 sessions of multifocal tDCS targeting the resting state motor network (network-tDCS). It was hypothesized, that 1) experimental pain for 24 hours would increase pain scores, reduce detection and pain thresholds, and inhibit the amplitude of the N2-P2 ERPs in the sham group, and 2) active networktDCS in contrast with sham tDCS would reduce pain scores, and modulate the reduction of detection and pain electrical thresholds, as well as revert the amplitude reduction of the N2-P2 ERPs responses.

Methods

Participants

Healthy, right-handed subjects (N = 38, 16 females) between 21 and 36 years old were recruited in this parallel, double-blinded, and randomized study. Before experiments, all subjects were informed about the procedures in writing and orally, and completed a tDCS safety screen questionnaire.² Subjects were randomly assigned into 1 of 2 independent and sexmatched groups by employing a stratified randomization (www.random.org): Active (n = 19; 8 females; age: 26.0±4.2 years; height: 1.74±.12 m; weight: 79.8 \pm 12.8 kg) and Sham network-tDCS (n = 19; 8 females; age: 27.1±2.7 years; height: 1.75±.09 m; weight: 73.1±15.3 kg). All subjects reported to be free of chronic pain or acute pain at the time of the experiment, chili (capsaicin) allergies, pregnancy as well as neurological, musculoskeletal, and mental conditions. These data are secondary outcomes from a previous study and the sample size estimation was performed based on primary outcome parameters (corticomotor excitability to single-pulse TMS)²⁴ and previous publications using a similar approach.^{29,34} The protocol was approved by the local ethics committee (VN-20180092), registered in clinicaltrials.gov (NCT04165980), and procedures were in conformity with the declaration of Helsinki.

Experimental Design

The effect of 24 hours of capsaicin-induced pain and 2 daily network-tDCS treatments (active/sham) were studied in 2 sessions on successive days (Fig 1) at the Center for Neuroplasticity and Pain (CNAP), Denmark. All participants were familiarized with the testing procedures before baseline assessments. The detection and pain threshold to electrical stimulation at the right volar wrist area, as well as ERPs evoked by painful electrical stimulation and their perceived pain intensity were recorded at the beginning of Day1 (Day1-baseline). Topical capsaicin was then applied on the right hand to induce experimental pain lasting 24 hours. Given that approximately 1 hour of topical capsaicin induces a robust pain and changes in cortical networks,¹ the assessment of the initial impact of capsaicin was conducted similarly to Day1baseline approximately 50 minutes after the capsaicin application (Day1-post-cap). Subsequently, the first



Figure 1. Experimental design of the study. After familiarization, the detection and pain thresholds to electrical stimulation as well as the event-related potentials (ERPs) to noxious electrical stimulation and self-reported pain scores due to electrical pulses delivered during ERPs recordings were registered at the right volar wrist area at the beginning of Day1 (Day1-baseline), 50 minutes after the first capsaicin application on the right hand (Day1-post-cap), and 50 minutes after the second capsaicin application on Day2 (Day2). Immediately after Day1-post-cap and after 24 hours, 20 minutes of network-tDCS (active or sham) was applied. NRS pain scores due to capsaicin were reported every 20 minutes at the lab after first and second capsaicin patch application on Day1 and Day2, respectively.

session of network-tDCS was applied. The next networktDCS session took place 24 hours after. Five minutes before the end of the second session of network-tDCS, capsaicin was reapplied on the right hand (beside the original capsaicin site) to ensure that pain-related responses would not be affected by the phenomenon of habituation. After 50 minutes of the second capsaicin application (Fig 1) the last assessment of ERPs and pain sensitivity were done (Day2, identical to the Day1-baseline and Day1-post-cap).

Electrical Stimulation and Related Sensations

Electrical stimulation was delivered through a stainless steel cutaneous pin electrode⁴³ placed on the right volar wrist at the level of the styloid process (proximal to application of experimental pain). Stimuli were delivered in 3pulse trains (each pulse duration was 200 μ s) and interpulse period of 5 ms⁵⁴ using an isolated electrical stimulator (NoxiTest IES 230; NoxiTest, Aalborg, Denmark). The electrical detection thresholds (EDT) and electrical pain thresholds (EPT) were assessed using an ascending method-of-limits.^{54,60} An ascending ramp of stimulus intensities was delivered using steps sizes of 0.02 mA and an inter-stimulus interval of 1 seconds. EDT was defined as the minimum current value at which the participant reported a tingling sensation. EPT was defined as the minimum current value at which the participant reported a painful pricking sensation. Three trials were run for both EDT and EPT, respectively, and the resulting values were averaged for further use. Forty ERPs were recorded with variable inter-stimulus intervals between 8 to 12 seconds to avoid habituation to the frequency of the stimulation, and the stimulus intensity was fixed as 2 times the EPT⁵⁴ measured at Day1-baseline to keep stimulus intensity constant across days. At the end of the stimulation for ERPs, subjects rated the electrically-induced pain on a numerical rating scale (NRS; anchored at 0: no pain, and at 10: worst pain imaginable). EDT and EPT values were normalized by computing the ratio between the individual values and the group mean value at Day1-post-cap and used for further analysis.

Recordings of Evoked Potentials

Electroencephalographic (EEG) activity was acquired through a 32-channel system (Starstim 32, Neuroelectrics, Spain) in the 10-10 international configuration using an EEG neoprene cap (NE056 Headcap R, Neuroelectrics, Spain) with Ag/AgCl electrodes (Neuroelectrics, Spain). Thirty-one EEG channels were used for EEG acquisition referenced to the right earlobe and the remaining channel (O2) was used to convey trigger information in order to synchronize EEG activity to electrical stimulation. Electrode impedance was secured below 5 k Ω . Sampling frequency was set at 500 Hz and a band-pass filtering was applied between .5 Hz to 40 Hz. The position of the EEG neoprene cap was systematically fixed considering the nasion-to-inion and tragus-to-tragus distances of each individual, and monitored throughout the experiment. During EEG recordings, participants sat relaxed in an armchair and kept their eyes open during the electrical stimulation (approximately 7-8 minutes).

Analysis of Evoked Potentials

Offline analysis was performed using EEGLAB v14.1.2b¹⁷ running under MATLAB (The Mathworks, Natick, MA). Visual inspection was done to identify noisy channels and movement artifacts. A linear bandpass filter with cutoff frequencies of .5 to 40 Hz was applied. Independent component analysis (ICA) of the filtered recordings was performed and this was followed by thorough inspection of ICA maps and ICA continuous recordings to extract eye movement, eye blinks, cardiac, muscle, and stimulation artifacts. Data were epoched with a duration of 3 seconds each, stimuluslocked from -1 to 2 seconds with time 0 corresponding to stimulus onset. Baseline correction was made using the -1 to 0 seconds window. The resultant baseline-corrected epochs were averaged to extract the ERPs of interest and stored for further analysis. The N2 and P2 components were identified with the recommended central-earlobe montage (Cz-A1), where the N2 was defined as the first major negative deflection after stimulus onset, and P2 was defined as the first major positive

Gregoret et al

deflection. The N2-P2 peak-to-peak amplitude was determined by combining the voltage of each peak of the biphasic component. Latencies were measured at the peak of each component. N2-P2 amplitudes as well as N2 and P2 latencies were normalized by calculating the ratio of each individual value and their group mean value at Day1-post-cap. Grand average (across all subjects) of N2-P2 responses and scalp maps of averaged N2 and P2 responses considering the range of mean latency (± SD) were performed for illustrative purposes.

Capsaicin-Induced Pain

Sustained pain was provoked for 24 hours, by applying a 4 cm x 4 cm patch of topical capsaicin (8%, Qutenza, Germany) on the distal skin area of the first metacarpus²⁰ on Day1 and on the center of third metacarpus on Day2 on the right hand. Both patches were kept on the same positions until the completion of the experiment on Day2. During the sessions at Day1 and Day2, subjects reported current pain intensity ratings due to topical capsaicin every 20 minutes. At the end of Day1 assessments, participants received a pain diary and were instructed to rate their pain every hour (off-lab hours) until they returned to the laboratory the following day (sleep hours excluded). Averaged NRS pain ratings were quantified at 6 different time points: using current pain NRS ratings reported during 1) first 50 minutes of capsaicin application on Day 1 (Day1-50minutes-cap), 2) after the 50 minutes of capsaicin application and before 1st tDCS session (Day1-before1st-tDCS), 3) during off-lab hours on Day1 (Day1-evening), 4) during morning hours on Day2 before the experiment (Day2-morning), 5) after 2nd tDCS session and 2nd capsaicin application (Day2-50minutescap), and 6) during the final 30 minutes of Day2 (Day2final30minutes).

Transcranial Direct Current Stimulation

Network-tDCS delivered an anodal current over bilateral motor cortices (C1, C2, C3, C4, and T8) and a cathodal current over posterior parietal (P3 and P4) and frontal (Fz) cortices using the electrode configuration designed in a previous study.²⁰ Using the 10-10 international EEG system, 3.14 cm² Ag/AgCl circular electrodes (PiStim, Neuroelectrics, Spain) were placed on the following positions administering the indicated currents: C1 = 872 μ A, C2 = 888 μ A, C3 = 1135 μ A, C4 = 922 μ A, F_Z = -1843 $\mu A, P3$ = -1121 $\mu A, P4$ = -1036 μA and T8 = 183 μ A.²⁰ Sessions of double-blinded tDCS were conducted using tDCS system (Starstim 32, Neuroelectrics, Spain) and specific software (NIC2, Neuroelectrics, Spain). Following evidence-based guidelines on the clinical use of tDCS, ^{39,59} 2 consecutive daily sessions of tDCS were delivered. Administration of active and sham network-tDCS lasted 20 minutes per session during which subjects were requested to keep their eyes open and remain relaxed. Active network-tDCS applied a constant current during the 20-minute period, whereas the sham network-tDCS administered a ramped current during the first 30 seconds and the last 30 seconds of the 20minute period while in-between no current was delivered. The examiner doing all psychophysical and electrophysiological recordings were blinded to the type of stimulation the subjects received.

Statistics

Data are presented as mean and standard deviation (SD) in text, figures and tables. Statistical analysis were carried out using SPSS (SPSS, v25.0, IBM) and significance was set at P<.05. Analysis of normal distribution was performed using visual inspection and Shapiro-Wilks' test. Data that did not show normal distribution (EDTs) were log-transformed and normality assessments were reconducted to apply parametric tests. Current intensity for eliciting ERPs in the 2 groups (Active and Sham network-tDCS) was compared by an independent t-test. Changes in averaged capsaicin-induced pain NRS scores were analyzed with a 2-way analysis of variance (ANOVA) with time (Day1-50minutes-cap, Day1-before1sttDCS, Day1-evening, Day2-morning, Day2-50minutes-cap, and Day2-final30minutes) as repeated factor, and group (Active or Sham networktDCS) as the between-group factor. The impact of topical capsaicin on EDT, EPT, latencies and peak-to-peak amplitudes of N2-P2 responses, and pain NRS ratings due to electrical stimulation during EP recordings, was tested by a 2-way ANOVA performed with the factors, time (Day1-baseline and Day1-post-cap) as the repeated factor and group (Active or Sham networktDCS) as the between-group factor. The impact of 2 sessions of tDCS on EDT, EPT, latencies and peak-to-peak amplitudes of N2-P2 responses, pain NRS ratings due to electrical stimulation during EP recordings, and capsaicin-induced current NRS pain scores was evaluated by a 2-way ANOVA with time (Day1-post-cap and Day2) as the repeated factor and group (Active or Sham network-tDCS) as the between group factor. When ANOVA factors or interactions were significant, posthoc analysis was performed using Bonferroni corrections for multiple comparison.

Results

Capsaicin-Induced Pain

Current pain NRS ratings were zero at Day1-baseline, 3.6 ± 1.8 and 4.3 ± 2.5 at Day1-post-cap, and 4.4 ± 2.2 and 5.5 ± 2.6 at Day2 in the active and sham network-tDCS group, respectively. ANOVA of current pain NRS ratings revealed a main effect of *time* (*P*<.01) showing that the current pain was increased on Day2 compared with Day1-post-cap. No main effect of *group* nor interactions were observed.

ANOVA of averaged pain NRS ratings (Table 1) revealed a main effect of *time*. Compared to Day1-post-cap, the averaged NRS pain ratings were increased at all time points (P<.001). There was no significant interaction, and no significant main effect of *group*.

Table 1. Mean (\pm SD) Pain Numerical Rating Scale (NRS) Ratings Following Capsaicin Application at Day1

	Active Network-tDCS	Sham Network-tDCS
Day1-post-cap	2.3±1.3	2.6±1.7
Day1-beforetDCS	4.2±1.8	5.1±2.5
off-lab hours on Day1	4.7±1.9	6.1±2.6
off-lab hours on Day2	4.6±1.9	5.7±2.8
Day2-after2ndtDCS	3.9±1.7	4.9±2.3
Day2-final30min	4.6±2.0	6.1±2.5

Electrical Stimulation

The ANOVAs of EDT and EPT (Table 2) showed no significant main effect of *time*, *group* nor interactions (Table 3). Likewise, the ANOVA of pain NRS scores during electrical stimulation for ERPs (Table 2) showed no significant main effect of *time*, *group*, or interactions (Table 3).

N2-P2 Components of Evoked Potentials

Recordings from 2 subjects in the sham group were excluded due to technical trigger issues. Therefore, recordings from 19 participants in the active and 17 participants in the sham group were analyzed. Stimulation intensity was 0.56 ± 0.16 mA and 0.63 ± 0.25 mA in the active and sham group, respectively, without significant differences between groups (t=-1.44, P = .260). The topographical maps of the evoked responses (Fig 2A-D) show the scalp distributions with maximum activity on central electrodes. The N2 and P2 ERPs were clearly detectable in the active and sham network-tDCS groups (Fig 2E, F)

The ANOVA of the N2-P2 peak-to-peak amplitude at Day1-baseline and Day1-post-cap showed a significant main effect of *time* (Table 3, Fig 2E-H). No significant main effect of *group* or interaction was observed. The ANOVA of the N2-P2 peak-to-peak amplitude at Day1-post-cap and Day2 showed a significant time-x-group interaction (P<.025, Table 3). Post-hoc analysis showed a significant N2-P2 increment at Day2 compared to Day1-post-cap (P<.05) only in the active tDCS group. Posthoc results also revealed significant group differences at Day2 (P<.05), showing the impact of active network-tDCS on N2P2 amplitudes, compared to sham network-tDCS.

Latency of N2-P2 Components

ANOVA of the N2 latency as well as P2 latency showed no significant effects over *time*, group and interactions

Table 2. Mean (\pm SD) Raw Electrical Detection Thresholds (EDT), Electrical Pain Thresholds (EPT), Latencies of the N2 and P2, as Well as Pain Numerical Rating Scale (NRS) Scores Due to the Electrical Stimulation at Day1-Baseline, Day1-Post-Cap, and Day2 in the Active and Sham Network-tDCS Groups

	Active Network-TDCS			Sham Network-tDCS		
	Day1-Baseline	Day1-Post-Cap	Day2	Day1-Baseline	Day1-Post-Cap	DAY2
EDT (mA)	.12±.06	.11±.03	.12±.04	.10±.05	.10±.05	.11±.04
EPT (mA)	.28±.08	.32±.10	.30±.12	.31±.13	.33±.16	.30±.14
N2 Latency (ms)	162.8±31.7	159.1±34.5	150.8±47.6	160.2±32.6	160.0±34.3	170.9±30.2
P2 latency (ms) Pain NRS during stim. (0–10)	302.6±26.2 3.6±1.3	304.3±39.2 4.0±1.2	297.1±41.7 3.9±1.6	302.8±23.1 4.4±1.8	305.8±22.6 4.6±1.9	294.8±31.7 4.3±2.0

Table 3. Results of ANOVA (F and p values) for Normalized Electrical Detection Thresholds (EDT), Electrical Pain Thresholds (EPT), Latencies and Peak-to-Peal Amplitudes of N2-P2 Event-Related Potentials, N2P2 Amplitudes, and Pain Numerical Rating Scale (NRS) Scores Due to the Electrical Stimulation at Day1-Baseline, Day1-Post-Cap, and Day2 in the Active and Sham Network-tDCS Groups

	RESULTS OF ANOVA FROM DAY1-BASELINE TO DAY1-POST-CAP			RESULTS OF ANOVA FROM DAY1-POST-CAP TO DAY2		
	Main Effect of Time	MAIN EFFECT OF GROUP	Time x Group Interaction	Main Effect of Time	Main Effect of Group	Time x Group Interaction
EDT	F = .01 <i>P</i> = .928	F = 2.04 <i>P</i> = .161	F = . 00 <i>P</i> = .996	F = .41 <i>P</i> = .524	F=2.39 P = .131	F = .21 <i>P</i> = .652
EPT	F = 3.56 <i>P</i> = .067	F = .11 <i>P</i> = .741	F = .47 <i>P</i> = .496	F = 2.54 <i>P</i> = .120	F = .07 <i>P</i> = .791	F = .52 <i>P</i> = .475
Latency N2	F = .30 <i>P</i> = .588	F = .03 <i>P</i> = .870	F = .23 <i>P</i> = .633	F = .04 <i>P</i> = .846	F = .81 <i>P</i> = .375	F = 2.23 P = .144
Latency P2	F = .21 <i>P</i> = .650	F = .01 <i>P</i> = .942	F = .01 <i>P</i> = .908	F = 2.72 <i>P</i> = .108	F = .03 <i>P</i> = .858	F = .12 <i>P</i> = .735
N2P2 amplitude	F = 6.08 P = .019	F = .40 <i>P</i> = .529	F = 1.03 <i>P</i> = .318	F = .07 <i>P</i> = .793	F = 1.33 P = .258	F = 5.61 P = .024
Pain NRS during stimulation	F = 2.76 <i>P</i> = .105	F = .07 <i>P</i> = .791	F = .51 <i>P</i> = .480	F = .97 <i>P</i> = .330	F = .02 <i>P</i> = .885	F = .15 <i>P</i> = .704



Figure 2. Scalp maps of averaged N2 (**A**,**B**) between 130 and 190 ms and P2 (**C**,**D**) between 275 and 325 ms in the active (left) and sham (right) Network-tDCS groups, respectively, at Day1-baseline, Day1-post-cap and Day 2. Grand-average evoked responses (**E**,**F**) and the mean (standard deviation; **G**,**H**) of normalized N2-P2 peak-to-peak amplitudes of evoked potentials in the active (left) and sham (right) network-tDCS at baseline (continuous black line), after capsaicin application (continuous gray line) and after active or sham tDCS (dashed line). Individual responses are illustrated in gray. Significantly decreased compared with Day1-baseline (*, *P*<.05). Significantly increased compared to Day1-post-cap (#<.05). Significantly higher compared to the sham network-tDCS group at Day2 (¤, *P*<.05).

at Day1-baseline and Day1-post-cap (Table 3). Likewise, no significant modulation of those latencies was observed over *time, group* nor interactions at Day1post-cap and Day2 (Table 3).

Discussion

The current study explored changes in cutaneous sensitivity and event-related potentials in response to electrical noxious stimulation before and after 2 sessions of multifocal tDCS over the motor network while experimental prolonged pain developed through a period of 24 hours. The results show that topical capsaicin induced sustained pain for 24 hours and exerted a reduction of peak-to-peak amplitude of N2-P2 ERPs already after 50 minutes of pain. Compared with a sham condition, network-tDCS did not elicit significant changes on the pain intensity elicited by topical capsaicin and by electrical noxious stimulations. With basis in the reduced N2-P2 ERPs after 50 minutes pain, subjects in the sham network-tDCS group demonstrated a sustained inhibition in the N2-P2 ERPs after 24 hours, while subjects in the active network-tDCS group showed a relative facilitation (towards normalization).

Effects of Network-tDCS on Capsaicin-Induced Pain and Electrical Sensory and Pain Sensitivity

Detection and pain thresholds to electrical skin stimulations were not significantly modulated by topical capsaicin for 50 minutes in line with previous studies where widespread hyperalgesia was not found during capsaicin application for up to 1 hour,⁶⁷ 3 hours, and 24 hours.⁶ The results of electrical cutaneous sensitivity as well as capsaicin-induced current pain and average pain intensity indicate that two sessions of network-tDCS did not influence these outcomes significantly. Evidence of tDCSdriven pain reduction and hypoalgesia is debated.^{23,39,53} In line with the results of the current study, a lack of modulation of sensory and pain sensitivity was reported when applying a battery of quantitative sensory testing (mechanical pain thresholds, pressure pain thresholds, warm detection and heat pain thresholds) after 3 consecutive sessions of either M1, M1 and dorsolateral prefrontal cortex (DLPFC), and DLPFC HD-tDCS³⁴ in pain-free individuals compared to sham stimulation. However, Boggio and colleagues reported increased electrical detection and pain thresholds when delivering 1 session of anodal M1 tDCS in healthy pain-free individuals.⁸ An explanation for the different findings between Boggio's study with the present work may rely on the application of 1 session of traditional anodal M1 tDCS (versus network tDCS), the size and materials of the electrodes (35 cm² sponge vs 3 cm² vs Ag/AgCl circular electrodes), the use of surface electrodes (versus pin electrodes) to deliver the current stimulation, the type of phasic pain stimulation in pain-free individuals, and, as suggested by previous authors,⁹ the antagonizing effects in sensitized pain pathways compared to pain-free non-sensitized subjects. Even though both network-tDCS and traditional anodal M1 tDCS facilitate corticomotor excitability,^{20,37} these paradigms could possibly induce differential behavioral responses. Network-tDCS stimulates a number of interconnected regions (left and right M1, medial prefrontal cortex, and posterior parietal cortices), whereas classical anodal M1 tDCS applies an anodal current on the left motor cortex and a cathodal current on the right supraorbital area. However, findings with classical anodal M1 tDCS,⁵⁸ M1 HD-tDCS,²² and network-tDCS²⁴ support the notion of activation of descending inhibitory pathways using tDCS on the M1. Indeed, network-tDCS improved

conditioned pain modulation and normalized corticomotor inhibition induced due to prolonged capsaicininduced pain in the present study cohort but published elsewhere.²⁴

Reduction of N2-P2 Responses During Topical Capsaicin-Induced Pain

The present findings extend the evidence of N2-P2 reduction due to capsaicin-induced pain from 30 minutes^{29,67} to at least 50 minutes, and, in the sham group, 24 hours. Moreover, the association between the N2-P2 amplitudes with the pain NRS scores during electrical stimulation across all participants at Day1-baseline and at Day1-post-cap, and only in the sham group at Day2 suggests a pain-related nature of this modulation. Accordingly, previous studies found reduced N2-P2 amplitude when elicited by stimulations in the topical capsaicin-induced secondary hyperalgesic skin area.^{29,67} Since self-reported pain during the stimulation were not significantly altered, this N2-P2 reduction was attributed to spinothalamic tract (STT) inhibition.⁶⁷ In contrast, intradermal capsaicin amplifies N2-P2 ERPs when based on stimulations in the secondary hyperalgesic area,⁴⁴ showing variability of N2-P2 responses among different pain models or pain severity. Equally unexpected are findings based on early somatosensory ERPs, where increased activity in the primary sensory cortex is produced during experimental muscle soreness¹⁶ and clinical disorders, 12 but decreased amplitude of early sensory ERPs were observed in other pain models eg injection of hypertonic saline inducing a short duration but intense pain.⁶² A descending inhibitory control effect is unlikely since pain scores during electrical stimulation did not change significantly over time nor between groups. Thus, as a working hypothesis it could be that less intense pain for longer time (topical capsaicin) reduces the amplitude of noxious ERPs whereas more severe pain models for shorter time (intradermal capsaicin) could amplify the noxious ERPs. Since N2-P2 ERPs are mainly generated in the anterior cingulate cortex and the operculoinsular area,¹² they are influenced by salient and attentional factors,¹⁵ and these factors could possibly explain the present results.^{45,66} Reduction of N2-P2 amplitudes has been attributed to downregulating saliency of noxious electrical stimulation when delivering nociceptive stimulation at a constant stimulation frequency in comparison to a variable 1.³⁰ Even though the present work aimed to minimize this effect by applying a variable stimulation frequency, current results were comparable with the study by lannetti et al,³⁰ indicating that such reduction may be attributed to a decrement of saliency to electrical stimuli due to attentional reorientation towards the topical capsaicin pain. Moreover, further work elicited N2-P2 ERPs³⁸ as a result of administrating nociceptive and non-nociceptive sensory, visual and auditory stimuli, suggesting that N2-P2 ERPs are non-specific to pain but correlated with the subjective degree of saliency across all modalities.⁴⁹ Therefore, N2-P2 potentials may be an indicator of the

Gregoret et al

level of saliency of an incoming stimulus,^{30,49} in this case affected by the capsaicin-induced pain rather than the perception of electrical-induced pain per se. Future studies should investigate the interplay between attentional reorientation⁶⁶ (using eg attention scales) towards the conditioning (in this case capsaicin) and testing stimulus (electrical pulses) when delivering salient noxious and salient sensory stimulation. Indeed, concurrent sensory and/or painful inputs⁶³ "compete to be represented in the neural system".^{42,41}

Effects of Network-tDCS on Cortical Pain Responses During Prolonged Pain

The present findings showed a significant modulation of N2-P2 amplitudes after network-tDCS. Although traditional anodal tDCS on M1 has previously demonstrated inconsistent effects on the N2-P2 amplitude, 1 study showed similar increase of the event-related N2-P2 responses, compared to baseline (not compared to sham).²⁶ It has been suggested that anodal M1 tDCS induces corticothalamic inhibition and endogenous opioid release as mechanisms of tDCS-induced analgesia. For instance, anodal M1 tDCS has produced a significant increment of functional coupling between the M1 and the ipsilateral thalamus in pain free individuals.⁵⁵ There is also evidence of decreased μ -opioid receptor binding after a session of anodal M1 tDCS in the thalamus, nucleus accumbens, ACC and insula,¹⁸ regions that are crucial for processing and integration of pain. Other studies show a reduction in regional cerebral blood flow (rCBF) in anterior insula, an area associated to salience detection and considered to be one of the core nodes of the salience network along with the dorsal ACC,48 after anodal M1 tDCS compared to cathodal M1 tDCS during heat pain stimulation.³¹ However, the anterior insula has also been associated to decision-making relative to salience detection. Future studies are therefore warranted to explore whether tDCS-driven modulation is non-specific to pain intensity but it rather elicits changes on the stimulus saliency, or on the other side, if such modulation is attributed to decision-making or salience detection.

Limitations

Attention and saliency were not evaluated (eg through attentional and saliency scales) during or after

References

1. Alhajri N, Boudreau SA, Graven-Nielsen T: Angular gyrus connectivity at alpha and beta oscillations is reduced during tonic pain – Differential effect of eye state. Neuro-image Clin 33:102907, 2022

2. Antal A, Alekseichuk I, Bikson M, Brockmöller J, Brunoni AR, Chen R, Cohen LG, Dowthwaite G, Ellrich J, Flöel A, Fregni F, George MS, Hamilton R, Haueisen J, Herrmann CS, Hummel FC, Lefaucheur JP, Liebetanz D, Loo CK, McCaig CD, Miniussi C, Miranda PC, Moliadze V, Nitsche MA, Nowak R, Padberg F, Pascual-Leone A, Poppendieck

The Journal of Pain 233

delivering electrical pulses (for ERP recordings), limiting the conclusions of capsaicin-induced reduction of N2-P2 potentials as well as the lack of significant modulation of self-reported pain due to tDCS. Second, the aim of this study was to investigate tDCS-driven modulation of N2P2 amplitudes following noxious stimulation during prolonged pain to understand cortical integration of noxious stimuli on sensitized individuals. Given that N2P2 responses are not restricted to nociceptive pathways and that cutaneous pin electrodes may co-activate in some circumstances large myelinated $A\beta$ fibers,⁵⁶ future research should investigate the modulatory effects of tDCS on N2P2 ERPs following nociceptive (laser) stimulation. Third, recruited subjects are young healthy individuals. It has been demonstrated that N2-P2 ERPs are age-dependent.⁵⁴ Therefore, the results of this study may not be transposable to individuals in different age ranges. The present study evaluated the modulatory effects of 2 sessions of tDCS; more specifically 50 minutes after the delivery of the second tDCS session on each subject since previous research shows significantly increased corticomotor output at 1 hour compared with 30 and 15 minutes after a network-tDCS session.²⁰ Future studies should evaluate though the immediate effects of tDCS on day 1 on ERPs amplitudes to understand the temporal dynamics of tDCS modulation during pain. Finally, since this is an exploratory study and the sample size was calculated based on primary outcome parameters (corticomotor excitability to single-pulse TMS).²⁴ future studies are needed to validate the current findings based on a priori sample size estimations.

Conclusions

The present study indicates that tDCS targeting regions associated to the motor network does not induce analgesia in experimental prolonged pain during 24 hours but is effective in the modulation of late event-related potentials in response to noxious peripheral stimulation. These findings extend the basic insight into the neurophysiological effects of tDCS and support further investigation of multifocal tDCS as a tool for modulating cortical evoked-responses. Further basic and clinical studies are necessary to prove the usefulness of tDCS in patients.

W, Priori A, Rossi S, Rossini PM, Rothwell J, Rueger MA, Ruffini G, Schellhorn K, Siebner HR, Ugawa Y, Wexler A, Ziemann U, Hallett M, Paulus W: Low intensity transcranial electric stimulation: Safety, ethical, legal regulatory and application guidelines. Clin Neurophysiol 128:1774-1809, 2017

3. Antal A, Terney D, Kühnl S, Paulus W: Anodal transcranial direct current stimulation of the motor cortex ameliorates chronic pain and reduces short intracortical inhibition. J Pain Symptom Manage 39:890-903, 2010

234 The Journal of Pain

4. Apkarian AV, Bushnell MC, Treede RD, Zubieta JK: Human brain mechanisms of pain perception and regulation in health and disease. Eur J Pain 9:463-484, 2005

5. Apkarian AV, Flor H: Functional Reorganization of the Brain With Chronic Pain. Chicago, Illinois, Wolters Kluwer Health Adis (ESP), 2015

6. Bement MH, Petersen KK, Sørensen LB, Andersen HH, Graven-Nielsen T: Temporal aspects of endogenous pain modulation during a noxious stimulus prolonged for 1 day. Eur J Pain (United Kingdom) 24:752-760, 2020

7. Binder A: Human surrogate models of neuropathic pain : Validity and limitations. Pain 157:48-52, 2016

8. Boggio PS, Zaghi S, Lopes M, Fregni F: Modulatory effects of anodal transcranial direct current stimulation on perception and pain thresholds in healthy volunteers. Eur J Neurol 15:1124-1130, 2008

9. Bradley C, Perchet C, Lelekov-Boissard T, Magnin M, Garcia-Larrea L: Not an aspirin: No evidence for acute anti-nociception to laser-evoked pain after motor cortex rTMS in healthy humans. Brain Stimul 9:48-57, 2016

10. Bromm B, Lorenz J: Neurophysiological evaluation of pain. Electroencephalogr Clin Neurophysiol 107:227-253, 1998

11. Chen C, Fang Y, Wang X, Bao SC, Tang Z, Tong RKY: Excitation comparison between multi-site stimulation using network-based tDCS and focal stimulation using high-definition tDCS. Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS IEEE; 1:6884–7, 2019.

12. Cruccu G, Aminoff MJ, Curio G, Guerit JM, Kakigi R, Mauguiere F, Rossini PM, Treede R, Garcia-larrea L, Lyon D: Recommendations for the clinical use of somatosensory-evoked potentials. Clin Neurophysiol 119:1705-1719, 2008

13. Csifcsak G, Antal A, Hillers F, Levold M: Modulatory effects of transcranial direct current stimulation. Pain Med 10:122-132, 2009

14. Dancey E, Yielder P, Murphy B: The interactive effect of tonic pain and motor learning on corticospinal excitability. Brain Sci 9:1-17, 2019

15. Davis KD, Moayedi M: Central mechanisms of pain revealed through functional and structural MRI. J Neuroimmune Pharmacol 8:518-534, 2013

16. De Martino E, Petrini L, Schabrun S, Graven-Nielsen T: Cortical somatosensory excitability is modulated in response to several days of muscle soreness. J Pain 19:1296-1307, 2018

17. Delorme A, Makeig S: EEGLAB : An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods 134:9-21, 2004

18. Dossantos MF, Love TM, Martikainen IK, Nascimento TD, Fregni F, Cummiford C, Deboer MD, Zubieta J, Dasilva AFM: Immediate effects of tDCS on the μ -opioid system of a chronic pain patient. Front Psychiatry 3:1-6, 2012

19. Farina S: Transient inhibition of the human motor cortex by capsaicin-induced pain. A study with transcranial magnetic stimulation. Neurosci Lett 314:97-101, 2001

20. Farina S, Tinazzi M, Le Pera D, Valeriani M: Pain-related modulation of the human motor cortex. Neurol Res 25:130-142, 2003

21. Fischer DB, Fried PJ, Ruffini G, Ripolles O, Salvador R, Banus J, Ketchabaw WT, Santarnecchi E, Pascual-Leone A, Fox MD: Multifocal tDCS targeting the resting state motor network increases cortical excitability beyond traditional tDCS targeting unilateral motor cortex. Neuroimage 157:34-44, 2017

22. Flood A, Waddington G, Cathcart S: High-definition transcranial direct current stimulation enhances conditioned pain modulation in healthy volunteers: a randomized trial. J Pain 17:600-605, 2016

23. Giannoni-Luza S, Pacheco-Barrios K, Cardenas-Rojas A, Mejia-Pando PF, Luna-Cuadros MA, Barouh JL, Gnoatto-Medeiros M, Candido-Santos L, Barra A, Caumo W, Fregni F: Non-invasive motor cortex stimulation effects on quantitative sensory testing (QST) in healthy and chronic pain subjects. Pain 161(9):1955-1975, 2020

24. Gregoret L, Zamorano AM, Graven-Nielsen T: Effects of multifocal transcranial direct current stimulation targeting the motor network during prolonged pain. Eur J Pain 25:1241-1253, 2021

25. Hanakawa T, Dimyan MA, Hallett M: Motor planning, imagery, and execution in the distributed motor network: A time-course study with functional MRI. Cereb Cortex 18:2775-2788, 2008

26. Hansen N, Obermann M, Poitz F, Holle D, Diener HC, Antal A, Paulus W, Katsarava Z: Modulation of human trigeminal and extracranial nociceptive processing by transcranial direct current stimulation of the motor cortex. Cephalalgia 31:661-670, 2011

27. Hill AT, Rogasch NC, Fitzgerald PB, Hoy KE: Effects of prefrontal bipolar and high-definition transcranial direct current stimulation on cortical reactivity and working memory in healthy adults. Neuroimage Elsevier 152:142-157, 2017

28. Hill AT, Rogasch NC, Fitzgerald PB, Hoy KE: Effects of single versus dual-site High-Definition transcranial direct current stimulation (HD-tDCS) on cortical reactivity and working memory performance in healthy subjects. Brain Stimul 1033-1043, 2018

29. Hüllemann P, Watfeh R, Shao Y, Nerdal A, Binder A, Baron R: Peripheral sensitization reduces laser-evoked potential habituation Réduction de l ' habituation des potentiels évoqués au laser. Neurophysiol Clin 45:457-467, 2015

30. Iannetti GD, Hughes NP, Lee MC, Mouraux A: Determinants of laser-evoked EEG responses: Pain perception or stimulus saliency. J Neurophysiol 100:815-828, 2008

31. Ihle K, Rodriguez-Raecke R, Luedtke K, May A: TDCS modulates cortical nociceptive processing but has little to no impact on pain perception. Pain 155:2080-2087, 2014

32. Jurkiewicz MT, Crawley AP, Mikulis DJ: Is rest really rest? resting-state functional connectivity during rest and motor task paradigms. Brain Connect 8:268-275, 2018

33. Klein T, Magerl W, Rolke R, Treede R: Human surrogate models of neuropathic pain. Pain 115:227-233, 2005

34. Kold S, Graven-Nielsen T: The effect of anodal high definition transcranial direct current stimulation on the pain sensitivity in a healthy population. Pain 162:1659-1668, 2021

35. Kucyi A, Davis KD: The dynamic pain connectome. Trends Neurosci 38:86-95, 2015

The Journal of Pain 235

Gregoret et al

36. Kuner R, Flor H: Structural plasticity and reorganisation in chronic pain. Nat Rev Neurosci 18:20-30, 2016

37. Kuo HI, Bikson M, Datta A, Minhas P, Paulus W, Kuo MF, Nitsche MA: Comparing cortical plasticity induced by conventional and high-definition 4×1 ring tDCS: A neurophysiological study. Brain Stimul 6:644-648, 2013

38. Lee MC, Mouraux A, lannetti GD: Characterizing the cortical activity through which pain emerges from nociception. J Neurosci 29:7909-7916, 2009

39. Lefaucheur JP, Antal A, Ayache SS, Benninger DH, Brunelin J, Cogiamanian F, Cotelli M, De Ridder D, Ferrucci R, Langguth B, Marangolo P, Mylius V, Nitsche MA, Padberg F, Palm U, Poulet E, Priori A, Rossi S, Schecklmann M, Vanneste S, Ziemann U, Garcia-Larrea L, Paulus W: Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tDCS). Clin Neurophysiol 128:56-92, 2017

40. Lefaucheur JP, Jarry G, Drouot X, Ménard-Lefaucheur I, Keravel Y, Nguyen JP: Motor cortex rTMS reduces acute pain provoked by laser stimulation in patients with chronic neuropathic pain. Clin Neurophysiol 121:895-901, 2010. Available from: https://doi.org/10.1016/j.clinph.2009.12.028.

41. Legrain V, Mancini F, Sambo CF, Torta DM, Ronga I, Valentini E: Cognitive aspects of nociception and pain. Bridging neurophysiology with cognitive psychology. Neurophysiol Clin 42:325-336, 2012

42. Legrain V, Van Damme S, Eccleston C, Davis KD, Seminowicz DA, Crombez G: A neurocognitive model of attention to pain: Behavioral and neuroimaging evidence. Pain 144:230-232, 2009

43. Lelic D, Mørch C, Hennings K, Andersen O, Drewes A: Differences in perception and brain activation following stimulation by large versus small area cutaneous surface electrodes. Eur J Pain 827-837, 2012

44. Liang M, Lee MC, O'Neill J, Dickenson AH, lannetti GD: Brain potentials evoked by intraepidermal electrical stimuli reflect the central sensitization of nociceptive pathways. J Neurophysiol 116:286-295, 2016

45. Lorenz J, Garcia-Larrea L: Contribution of attentional and cognitive factors to laser evoked brain potentials. Neurophysio Clin 33:293-301, 2003

46. Maezawa H, Vicario CM, Kuo M-F, Hirata M, Mima T, Nitsche MA: Effects of bilateral anodal transcranial direct current stimulation over the tongue primary motor cortex on cortical excitability of the tongue and tongue motor functions. Brain Stimul 13:2019-2021, 2019

47. Matsunaga K, Nitsche MA, Tsuji S, Rothwell JC: Effect of transcranial DC sensorimotor cortex stimulation on somatosensory evoked potentials in humans. Clin Neurophysiol 115:456-460, 2004

48. Menon V, Uddin LQ: Saliency, switching, attention and control: A network model of insula function. Brain Struct Funct 214:655-667, 2010

49. Mouraux A, lannetti GD: Nociceptive laser-evoked brain potentials do not reflect nociceptive-specific neural activity. J Neurophysiol 101:3258-3269, 2009

50. Mylius V, Borckardt JJ, Lefaucheur J: Noninvasive cortical modulation of experimental pain. Pain 153:1350-1363, 2012

51. Naegel S, Biermann J, Theysohn N, Kleinschnitz C, Diener HC, Katsarava Z, Obermann M, Holle D: Polarity-specific modulation of pain processing by transcranial direct current stimulation - A blinded longitudinal fMRI study. J Headache Pain 19, 2018

52. Nitsche MA, Paulus W: Transcranial direct current stimulation - Update 2011. Restor Neurol Neurosci 29:463-492, 2011

53. O'Neill F, Sacco P, Bowden E, Asher R, Burnside G, Cox T, Nurmikko T: Patient-delivered tDCS on chronic neuropathic pain in prior responders to TMS (A randomized controlled pilot study). J Pain Res 11:3117-3128, 2018

54. Özgül ÖS, Maier C, Enax-Krumova EK, Vollert J, Fischer M, Tegenthoff M, Höffken O: High test-retest-reliability of pain-related evoked potentials (PREP) in healthy subjects. Neurosci Lett 647:110-116, 2017

55. Polanía R, Nitsche MA, Paulus W: Modulating functional connectivity patterns and topological functional organization of the human brain with transcranial direct current stimulation. Hum Brain Mapp 32:1236-1249, 2011

56. Poulsen AH, Tigerholm J, Meijs S, Andersen OK, Mørch CD: Comparison of existing electrode designs for preferential activation of cutaneous nociceptors. J Neural Eng 17:1-16, 2020

57. Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, Keefe FJ, Mogil JS, Ringkamp M, Sluka KA, Song XJ, Stevens B, Sullivan MD, Tutelman PR, Ushida T, Vader K: The revised international association for the study of pain definition of pain: Concepts, challenges, and compromises. Pain NLM (Medline) 161:1976-1982, 2020

58. Reidler JS, Mendonca ME, Santana MB, Wang X, Lenkinski R, Motta AF, Marchand S, Latif L, Fregni F: Effects of motor cortex modulation and descending inhibitory systems on pain thresholds in healthy subjects. J Pain 13:450-458, 2012

59. Ribeiro H, Sesterhenn RB, De Souza A, De Souza AC, Alves M, Machado JC, Burger NB, Torres IL da S, Stefani LC, Fregni F, Caumo W: Preoperative transcranial direct current stimulation: Exploration of a novel strategy to enhance neuroplasticity before surgery to control postoperative pain. A randomized sham-controlled study. PLoS One 12:1-17, 2017

60. Sang CN, Max MB, Gracely RH: Stability and reliability of detection thresholds for human A-Beta and A-Delta sensory afferents determined by cutaneous electrical stimulation. J Pain Symptom Manage 25:64-73, 2003

61. Schabrun SM, Christensen S, Mrachacz-kersting N, Graven-Nielsen T: Motor cortex reorganization and impaired function in the transition to sustained muscle pain. Cereb Cortex 26:1878-1890, 2016

62. Schabrun SM, Jones E, Kloster J, Hodges PW: Temporal association between changes in primary sensory cortex and corticomotor output during muscle pain. Neuroscience 235:159-164, 2013

63. Squintani G, Rasera A, Segatti A, Concon E, Bonetti B, Valeriani M, Tinazzi M: Conditioned pain modulation affects the N2P2 complex but not the N1 wave: A pilot study with laser-evoked potentials. Eur J Pain 25:550-557, 2021

236 The Journal of Pain

64. Tamura Y, Hoshiyama M, Inui K, Nakata H, Qiu Y, Ugawa Y, Inoue K, Kakigi R: Facilitation of A[delta]-fibermediated acute pain by repetitive transcranial magnetic stimulation. Neurology 62:2176-2181, 2004

65. Tiemann L, Hohn VD, Ta-Dinh S, May ES, Nickel MM, Gross J, Ploner M: Distinct patterns of brain activity mediate perceptual and motor and autonomic responses to noxious stimuli. Nat Commun 9:1-12, 2018

66. Torta DM, Jure FA, Andersen OK: Biurrun Manresa JAB: Intense and sustained pain reduces cortical responses to auditory stimuli: Implications for the interpretation of the effects of heterotopic noxious conditioning stimulation in humans. Eur J Neurosci 50:1-10, 2019

67. Valeriani M, Arendt-Nielsen L, Le Pera D, Restuccia D, Rosso T, De Armas L, Maiese T, Fiaschi A, Tonali P, Tinazzi M: Short-term plastic changes of the human nociceptive system following acute pain induced by capsaicin. Clin Neurophysiol 114:1879-1890, 2003

68. Zheng W, Woo C, Yao Z, Goldstein P, Atlas LY, Roy M, Schmidt L, Krishnan A, Jepma M, Hu B, Wager TD: Painevoked reorganization in functional brain networks. Cereb Cortex 00:1-19, 2019