

Contemporary management of atrial fibrillation and the predicted vs. absolute risk of ischaemic stroke despite treatment

a report from ESC-EHRA EORP-AF Long-Term General Registry

Ding, Wern Yew; Blomström-Lundqvist, Carina; Fauchier, Laurent; Marin, Francisco; Potpara, Tatjana S.; Boriani, Giuseppe; Lip, Gregory Y. H.; ESC-EHRA EORP-AF Long-Term General Registry Investigators; Joensen, A. Marni; Gammelmark, A.; Rasmussen, L. Hvilsted; Dinesen, P.; Riahi, S.; Venø, S. Krogh; Sorensen, B.; Korsgaard, A.; Andersen, K.; Hellum, C. Fragtrup

Published in:
Europace

DOI (link to publication from Publisher):
[10.1093/europace/euac214](https://doi.org/10.1093/europace/euac214)

Creative Commons License
CC BY-NC 4.0

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Ding, W. Y., Blomström-Lundqvist, C., Fauchier, L., Marin, F., Potpara, T. S., Boriani, G., Lip, G. Y. H., ESC-EHRA EORP-AF Long-Term General Registry Investigators, Joensen, A. M., Gammelmark, A., Rasmussen, L. H., Dinesen, P., Riahi, S., Venø, S. K., Sorensen, B., Korsgaard, A., Andersen, K., & Hellum, C. F. (2023). Contemporary management of atrial fibrillation and the predicted vs. absolute risk of ischaemic stroke despite treatment: a report from ESC-EHRA EORP-AF Long-Term General Registry. *Europace*, 25(2), 277-282. Article euac214. <https://doi.org/10.1093/europace/euac214>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.








- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: December 04, 2025

Contemporary management of atrial fibrillation and the predicted vs. absolute risk of ischaemic stroke despite treatment: a report from ESC-EHRA EORP-AF Long-Term General Registry

Wern Yew Ding ¹, Carina Blomström-Lundqvist ², Laurent Fauchier ³, Francisco Marin ⁴, Tatjana S. Potpara ^{5,6}, Giuseppe Boriani ^{7†}, and Gregory Y. H. Lip ^{1,8*†}; on behalf of the ESC-EHRA EORP-AF Long-Term General Registry Investigators

¹Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest Hospital, Liverpool, UK; ²Department of Cardiology, School of Medical Sciences, Faculty of Medicine and Health, Örebro University, Örebro, Sweden; ³Service de Cardiologie, Centre Hospitalier Universitaire Trousseau, Tours, France; ⁴Department of Cardiology, Hospital Universitario Virgen de la Arrixaca, IMIB-Arrixaca, University of Murcia, CIBERCV, Murcia, Spain; ⁵School of Medicine, University of Belgrade, Belgrade, Serbia; ⁶Intensive Arrhythmia Care, Cardiology Clinic, Clinical Center of Serbia, Belgrade, Serbia; ⁷Cardiology Division, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Policlinico di Modena, Modena, Italy; and ⁸Aalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

Received 28 June 2022; accepted after revision 30 September 2022; online publish-ahead-of-print 25 November 2022

Background

Risk stratification in patients with atrial fibrillation (AF) is important to facilitate guideline-directed therapies. The Calculator of Absolute Stroke Risk (CARS) scheme enables an individualized estimation of 1-year absolute risk of stroke in AF. We aimed to investigate the predicted and absolute risks of ischaemic stroke, and evaluate whether CARS (and CHA₂DS₂-VASc score) may be useful for identifying high risk patients with AF despite contemporary treatment.

Methods

We utilized the EORP-AF General Long-Term Registry which prospectively enrolled patients with AF from 250 centres across 27 participating European countries. Patients with sufficient data to determine CARS and CHA₂DS₂-VASc score, and reported outcomes of ischaemic stroke were included in this analysis. The primary outcome of ischaemic stroke was recorded over a 2-year follow-up period.

Results

A total of 9444 patients were included (mean age 69.1 [±11.4] years; 3776 [40.0%] females). There was a high uptake (87.9%) of anticoagulation therapy, predominantly with vitamin K antagonist (50.0%). Over a mean follow-up period of 24 months, there were a total of 101 (1.1%) ischaemic stroke events. In the entire cohort, the median CARS and absolute annual risks of ischaemic stroke were 2.60 (IQR 1.60–4.00) and 0.53% (95%CI 0.43–0.64%), respectively. There was no statistical difference between the predictive performance of CARS and CHA₂DS₂-VASc score (0.621 [95%CI 0.563–0.678] vs. 0.626 [95%CI 0.573–0.680], $P = 0.725$).

Conclusion

Contemporary management of AF was associated with a low risk of ischaemic stroke. CARS and CHA₂DS₂-VASc score may be useful to identify high risk patients despite treatment who may benefit from more aggressive treatment and follow-up.

Keywords

Atrial fibrillation • Predictive stroke risk • Absolute stroke risk • Ischaemic stroke • CARS • CHA₂DS₂-VASc score • EORP-AF

* Corresponding author. Tel: +0151 794 9020. E-mail address: gregory.lip@liverpool.ac.uk

† Joint senior authors.

© The Author(s) 2022. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

What’s new?

- Contemporary management, including anticoagulation, among patients with atrial fibrillation was associated with a reduction of 86% of the absolute risk of ischaemic stroke as predicted by CARS.
- Both CARS and CHA₂DS₂-VASc score had modest predictive capability for absolute ischaemic stroke events among anticoagulated patients with atrial fibrillation.
- The predictive capability of CARS and CHA₂DS₂-VASc score was comparable for ischaemic stroke events.
- The predictive capability of CARS was similar in subgroups by CHA₂DS₂-VASc risk category and anticoagulation therapy.

Introduction

Atrial fibrillation (AF) is associated with a significant risk of thromboembolism and a doubling of the mortality rate, with increased hospitalizations and healthcare costs.^{1,2} Management of patients with AF includes a detailed stroke risk assessment, typically performed using the CHA₂DS₂-VASc score.³ The information derived from this assessment may be used to guide the prescription of anticoagulation therapy,^{4,5} which was traditionally the only established treatment associated with a prognostic benefit in AF. However, there is emerging evidence to suggest that appropriate characterization and evaluation^{6,7} followed by a more holistic or integrated care management approach⁸ as well as early rhythm control^{9,10} may also offer a survival advantage. Hence, it is important to determine the residual risk of stroke in patients receiving contemporary treatment for AF to identify those who may benefit from additional forms of intervention. Yet, there is currently no validated model for this purpose.

Recently, Lee et al. proposed the Calculator of Absolute Stroke Risk (CARS) which enables an individualized estimation of 1-year absolute risk of stroke in AF.¹¹ CARS was derived from a large real-world retrospective Danish cohort of non-anticoagulated patients from 1997 to 2015 with first-diagnosed AF. It utilizes the same risk factors as the CHA₂DS₂-VASc score but analyses age as a continuous variable and considers the individual contribution of risk factors. The role of CARS in a prospective cohort of patients receiving treatment for AF remains to be determined.

Herein, we aimed to investigate the predicted and absolute risks of ischaemic stroke, and evaluate whether CARS (and CHA₂DS₂-VASc score) may be useful for identifying high risk patients in a prospective cohort of patients with AF receiving contemporary management. We investigated this using the European Society of Cardiology-European Heart Rhythm Association (ESC-EHRA) EURObservational Research Programme EORP-AF General Long-Term Registry.

Methods

The EORP-AF General Long-Term Registry is a prospective, observational, large-scale multicentre registry from 250 centres across 27 participating European countries. A detailed description of the study design has previously been provided.¹² In brief, patients with AF who presented to cardiology services were enrolled between October 2013 and September 2016. All patients were over 18 years old and had electrocardiographic confirmation of AF within 12 months of enrolment. Institutional review board approval of the study protocol was obtained for every institution, and the study was performed in accordance to the European Union Note for Guidance on Good Clinical Practice CPMP/ECH/135/95 and the Declaration of Helsinki. For the purposes of this analysis, we included only patients with sufficient data to calculate their individual CARS and CHA₂DS₂-VASc score, and without missing information on the outcome of ischaemic stroke.

Table 1 Baseline characteristics

Baseline characteristics	n = 9444
Age (years), mean (±SD)	69.1 (±11.4)
Female sex, n (%)	3776 (40.0%)
BMI (kg/m ²), mean (±SD)	28.3 (±5.1)
eGFR (mL/min/1.73 m ²), mean (±SD)	69.4 (±21.2)
LA diameter (mm), mean (±SD)	45.4 (±8.2)
LV ejection fraction (%), mean (±SD)	53.7 (±13.0)
Left ventricular hypertrophy, n (%)	2181 (27.5%)
AF classification, n (%)	
First-detected	1452 (15.6%)
Paroxysmal	2468 (26.6%)
Persistent	1850 (19.9%)
Long-standing persistent	420 (4.5%)
Permanent	3099 (33.4%)
EHRA classification, n (%)	
I	4283 (45.4%)
II	3373 (35.7%)
III	1592 (16.9%)
IV	196 (2.1%)
Comorbidities, n (%)	
Cardiomyopathy	1367 (14.5%)
Congenital heart disease	103 (1.1%)
COPD	815 (8.7%)
Coronary artery disease	2551 (28.3%)
Diabetes mellitus	2142 (22.7%)
Heart failure	3569 (37.8%)
Hypertension	5843 (61.9%)
Liver disease	236 (2.5%)
Peripheral vascular disease	748 (7.9%)
Previous haemorrhagic event	501 (5.3%)
Previous thromboembolic event	1093 (11.6%)
Previous ischaemic stroke	572 (6.1%)
Sleep apnoea	440 (4.8%)
Hyperthyroidism	168 (1.8%)
Hypothyroidism	511 (5.5%)
Valvular heart disease	4629 (49.7%)
Prior catheter AF ablation, n (%)	457 (6.0%)
CHA ₂ DS ₂ -VASc score, mean (±SD)	3.1 (±1.8)
HAS-BLED score, mean (±SD)	1.6 (±1.1)

AF, atrial fibrillation; BMI, body mass index; COPD, chronic obstructive pulmonary disease; eGFR, estimated glomerular filtration rate; EHRA, European Heart Rhythm Association; LA, left atrium; LV, left ventricle; SD, standard deviation.

Data on demographics, comorbidities and medications were collected at baseline. These were used to determine the predicted stroke risk by CARS¹¹ and CHA₂DS₂-VASc score. Estimated glomerular filtration rate (eGFR) was assessed using the Chronic Kidney Disease-Epidemiology Collaboration (CKD-EPI) equation. Patients with a CHA₂DS₂-VASc score of 0, 1–3 and more than 3 were designated as low risk, high risk and very high risk, respectively. The predicted residual 1-year stroke risk

Table 2 Baseline medication use

Baseline medication use	n = 9444
Anti-thrombotic, n (%)	8876 (94.0%)
Any antiplatelet	1769 (18.7%)
Acetylsalicylic acid	1588 (16.8%)
Clopidogrel	547 (5.8%)
Oral anticoagulants, n (%)	8296 (87.9%)
Vitamin K antagonist	4721 (50.0%)
Any NOAC	3429 (36.3%)
Apixaban	956 (10.1%)
Dabigatran	756 (8.0%)
Edoxaban	94 (1.0%)
Rivaroxaban	1623 (17.2%)
Heparin	146 (1.5%)
Anti-arrhythmics, n (%)	2610 (27.7%)
Amiodarone	1638 (17.4%)
Flecainide	341 (3.6%)
Propafenone	325 (3.5%)
Sotalol	275 (2.9%)
Other medications, n (%)	
ACE inhibitors	3944 (41.9%)
Aldosterone blockers	1673 (17.8%)
Angiotensin receptor blocker	1843 (19.6%)
Beta-blockers	6529 (69.4%)
Dihydropyridine CCB	1617 (17.2%)
Digoxin	1321 (14.0%)
Diuretics	4761 (50.6%)
Insulin	510 (5.4%)
Oral antidiabetics	1425 (15.1%)
Statins	3999 (42.5%)

ACE, angiotensin converting enzyme; CCB, calcium-channel blocker; eGFR, estimated glomerular filtration rate; NOAC, non-vitamin K antagonist oral anticoagulant.

('mCARS') was estimated as previously described.¹³ For this analysis, the primary study outcome was ischaemic stroke. This was recorded over a 2-year follow-up period.

Statistical analyses

Continuous variables were described with mean and standard deviation (SD), and tested for differences with *t*-test if normally distributed. Non-parametric variables are shown as median (IQR, interquartile range). Categorical variables were described with count and percentage, and tested for differences with chi-squared test. Absolute stroke risk at 1-year was expressed as incidence risk with 95% confidence intervals (CI). The predictive capability of CARS for absolute stroke risk at 1-year was investigated using receiver-operating characteristic (ROC) curves and area under the curve (AUC) was used to represent its ability to predict ischaemic stroke events. The performance of CARS was tested against the CHA₂DS₂-VASc score using DeLong's test.¹⁴ Further subgroup analyses were performed according to stroke risk profile by CHA₂DS₂-VASc score and oral anticoagulation agent (vitamin K antagonist [VKA] or non-vitamin K antagonist oral anticoagulant [NOAC]). A two-sided *p* value of less than 0.05 was considered statistically significant. Statistical analyses were performed using SPSS

Table 3 Predicted vs. absolute stroke risk

CARS, %	
Mean (±SD)	3.81 (±4.06)
Median (IQR)	2.60 (1.60–4.00)
Range	0.10–24.10
mCARS, %	
Mean (±SD)	1.37 (±1.46)
Median (IQR)	0.94 (0.58–0.94)
Range	0.04–8.68
Absolute stroke risk	
Number of events, n	101
Annual risk (95% CI), %	0.53 (0.43–0.64)

CARS, Calculator of Absolute Stroke Risk; CI, confidence interval; IQR, interquartile range; SD, standard deviation.

version 24 (IBM Corp, Armonk, NY) and R version 4.0.3 (R Foundation for Statistical Computing, Vienna, Austria).

Results

A total of 9444 patients were included in the present analysis, corresponding to 85.1% of the original cohort of 11 096 patients. The mean age of patients was 69.1 (±11.4) years with 3776 (40.0%) females (Table 1). Majority of patients had permanent AF (33.4%), and the mean left atrial diameter and eGFR were 45.4 (±8.2) mm and 69.4 (±21.2) mL/min/1.73 m², respectively. Few patients (19.0%) had severe symptoms based on European Heart Rhythm Association symptom scale (III–IV). The most common comorbidities were hypertension (61.9%), valvular heart disease (49.7%), heart failure (37.8%), coronary artery disease (28.3%), diabetes mellitus (22.7%), and cardiomyopathy (14.5%). Mean CHA₂DS₂-VASc score was 3.1 (±1.8).

There was a high uptake (87.9%) of anticoagulation therapy, predominantly with VKA (50.0%). The use of anticoagulation therapy according to CHA₂DS₂-VASc score is shown in [Supplementary material online, Table S1](#). Rivaroxaban was the most frequently (17.2%) prescribed NOAC, followed by apixaban (10.1%), dabigatran (8.0%), and edoxaban (1.0%). Other medications used at baseline can be found in Table 2.

Predicted vs. absolute stroke risk

Over a mean follow-up period of 24 months, there were a total of 101 (1.1%) ischaemic stroke events. In the entire cohort, the absolute annual risk of ischaemic stroke was 0.53% (95%CI 0.43–0.64%) (Table 3). The median CARS and mCARS were 2.60 (IQR 1.60–4.00) and 0.94 (IQR 0.58–0.94), respectively. Based on the absolute risk of ischaemic stroke, there was an 86% reduction from predicted stroke risk in this cohort. The relative risk reduction appeared to be more pronounced in patients with greater CHA₂DS₂-VASc score (Table 4).

Predictive performance of CARS and CHA₂DS₂-VASc score

Receiver-operating characteristics curves for absolute stroke events by CARS and CHA₂DS₂-VASc score are shown in Figure 1. The AUCs for CARS and CHA₂DS₂-VASc score were 0.621 (95%CI 0.563–0.678) vs. 0.626 (95%CI 0.573–0.680), respectively. There was no statistical

Table 4 Predicted vs. absolute stroke risk stratified by CHA₂DS₂-VASc score

CHA ₂ DS ₂ -VASc score	CARS (%)		mCARS (%)		Absolute stroke risk	
	Mean (\pm SD)	Range	Mean (\pm SD)	Range	Number of events, n	Annual risk (95% CI), %
CHA ₂ DS ₂ -VASc score 0 (n = 627)	0.68 (\pm 0.80)	0.10–10.60	0.25 (\pm 0.29)	0.04–3.82	3	0.24 (0–0.51)
CHA ₂ DS ₂ -VASc score 1 (n = 1225)	1.15 (\pm 0.85)	0.10–12.00	0.42 (\pm 0.30)	0.04–4.32	6	0.24 (0.05–0.44)
CHA ₂ DS ₂ -VASc score 2 (n = 1736)	2.00 (\pm 1.37)	0.10–17.10	0.72 (\pm 0.49)	0.04–6.16	14	0.40 (0.19–0.61)
CHA ₂ DS ₂ -VASc score 3 (n = 1959)	3.04 (\pm 2.10)	0.10–15.90	1.09 (\pm 0.76)	0.04–5.72	17	0.43 (0.23–0.64)
CHA ₂ DS ₂ -VASc score 4 (n = 1871)	4.35 (\pm 3.03)	0.40–20.10	1.57 (\pm 1.09)	0.14–7.24	22	0.59 (0.34–0.83)
CHA ₂ DS ₂ -VASc score 5 (n = 1171)	5.90 (\pm 4.16)	1.00–20.60	2.12 (\pm 1.50)	0.36–7.42	22	0.94 (0.55–1.33)
CHA ₂ DS ₂ -VASc score 6 (n = 550)	9.45 (\pm 5.66)	2.40–22.70	3.40 (\pm 2.04)	0.86–8.17	11	1.00 (0.41–1.59)
CHA ₂ DS ₂ -VASc score 7 (n = 219)	13.69 (\pm 5.26)	3.80–24.10	4.93 (\pm 1.89)	1.37–8.68	5	1.14 (0.14–2.14)
CHA ₂ DS ₂ -VASc score 8 (n = 74)	16.90 (\pm 1.99)	13.60–22.20	6.08 (\pm 0.72)	4.90–7.99	1	0.68 (0–2.02)
CHA ₂ DS ₂ -VASc score 9 (n = 12)	17.63 (\pm 1.62)	15.90–21.00	6.35 (\pm 0.58)	5.72–7.56	0	NA

CARS, Calculator of Absolute Stroke Risk; CI, confidence interval; IQR, interquartile range; NA, not available; SD, standard deviation.

difference between the predictive performance of CARS and CHA₂DS₂-VASc score, $P = 0.725$.

Subgroup analysis by CHA₂DS₂-VASc score and risk category

The predicted and absolute stroke risks stratified by CHA₂DS₂-VASc score are displayed in Table 4. The mean CARS ranged from 0.68% to 17.63% while the absolute annual stroke risk ranged from 0.24% to 1.14%. In general, both CARS and absolute annual stroke risk were greater with increasing CHA₂DS₂-VASc score (Figure 2). Patients in the low risk category (score 0) had an absolute annual stroke risk of 0.24% compared with 0.38% in the high risk category (score 1–3) and 0.78% in the very high risk category (score >3) (see [Supplementary material online, Table S2](#)).

Receiver-operating characteristic curves for absolute stroke events by CARS according to CHA₂DS₂-VASc risk category are shown in [Supplementary material online, Figure S1](#). The AUCs for CARS in low risk, high risk and very high risk patients were 0.595 (95%CI 0.361–0.829, 0.512 (95%CI 0.432–0.592) and 0.623 (95%CI 0.553–0.693), respectively. There were no statistical difference between the AUCs for each risk category (high risk vs. low risk, $P = 0.444$; very high risk vs. low risk, $P = 0.822$).

Subgroup analysis by anticoagulation therapy

The baseline characteristics of subgroups treated with either VKA or NOAC are shown in [Supplementary material online, Table S3](#). Patients in the VKA subgroup were older with higher prevalence of permanent AF, worse echocardiographic parameters, and a greater burden of comorbidities. As a result, the mean CHA₂DS₂-VASc and HAS-BLED scores were increased among patients treated with VKA compared to NOAC (CHA₂DS₂-VASc: 3.4 vs. 3.1, $P = 0.022$; HAS-BLED 1.8 vs. 1.4, $P < 0.001$). The mean CARS in the VKA subgroup was 4.02 (\pm 4.08) and NOAC subgroup was 3.78 (\pm 4.11) (see [Supplementary material online, Table S4](#)). The absolute annual stroke risk in the VKA and NOAC subgroups were 0.49 (95%CI 0.35–0.63) % and 0.24 (95%CI 0.16–0.32) %, respectively.

Receiver-operating characteristic curves for absolute stroke events in patients receiving VKA vs. NOAC therapy are shown in [Supplementary material online, Figure S2](#). The AUCs for CARS in the

VKA subgroup was 63.3% (95%CI 54.9–71.7%) and NOAC subgroup was 67.1% (95%CI 57.5–76.7%). There was no statistical difference between the predictive performance of CARS in either of these subgroups with $P = 0.557$.

Discussion

The main findings from this study of predominantly anticoagulated patients with AF in a prospective multicentre registry across 27 participating European countries were as follows: (1) absolute risk of ischaemic stroke was 86% lower than the predicted risk by CARS among this cohort; (2) both CARS and CHA₂DS₂-VASc score had modest predictive capability for absolute ischaemic stroke events among anticoagulated patients; (3) predictive capability of CARS and CHA₂DS₂-VASc score was comparable for ischaemic stroke events; and (4) predictive capability of CARS was similar in subgroups by CHA₂DS₂-VASc risk category and anticoagulation therapy.

In this contemporary cohort of treated patients with AF, the risk of ischaemic stroke in this cohort was lower compared with the predicted risk by CARS. Furthermore, the relative risk reduction of predicted vs. absolute ischaemic stroke risk appeared to be greater in patients with increased CHA₂DS₂-VASc score. The reduction in risk of ischaemic stroke was likely driven by the high uptake of oral anticoagulation therapy which is known to be beneficial for this purpose in AF. Nonetheless, the observed risk reduction was greater than 64%, as previously reported in a meta-analysis comparing OAC use against controls,¹⁵ suggesting the interplay of other factors. Although our study was not designed to compare the effects of contemporary vs. historical treatment in AF patients, it may be postulated that better understanding of the underlying pathophysiological mechanisms of AF,¹⁶ alongside the implementation of newer oral anticoagulation agents¹⁷ may have led to improved outcomes in this condition.

In this study, we evaluated the use of CARS and CHA₂DS₂-VASc score to predict ischaemic stroke risk in real-world patients. Although the CHA₂DS₂-VASc score had a marginally better predictive score, this was not statistically significant. The advantage of CARS over the CHA₂DS₂-VASc score in this situation is that it may be modified (mCARS) to provide an individual risk estimate of absolute risk that is easy to interpret and may be useful to facilitate patient-centred discussions. Furthermore, CARS is a more dynamic tool that allows for the adjustment of age in a continuous manner.




Figure 1 Receiver-operating characteristic curves for absolute stroke events by CARS vs. CHA₂DS₂-VASc score.




Figure 2 Plot of CARS vs. absolute annual stroke risk stratified by CHA₂DS₂-VASc score.

Using CARS and CHA₂DS₂-VASc score, patients with a high residual risk of ischaemic stroke despite treatment^{18,19} may be identified for more intensive follow-up and holistic management strategies. Indeed, adherence to the guideline-recommended Atrial fibrillation Better Care (ABC) pathway is associated with a 45% reduction in ischaemic stroke, compared to non-adherence.⁸ To reinforce the possible role of CARS, we also demonstrated that the (individualized) predictive capability of this tool was maintained regardless of CHA₂DS₂-VASc risk category and choice of anticoagulation therapy. Additionally, the calculated baseline absolute risk with CARS could be used to estimate the impact of specific treatments in various study cohorts. It may also be

incorporated with reported outcomes to determine the NNTnet (number needed to treat for net effect), which combines the benefit-and-harm of an intervention or therapy.²⁰

Limitations

The major limitation of this study relates to its observational nature with potential for misclassification bias. Furthermore, as the EORP-AF General Long-Term Registry was based exclusively on cardiology practices, our findings may not be generalizable to other cohorts. The annual absolute stroke risk was determined by assuming a linear event rate over the 2 year follow-up period. There were few ischaemic stroke events in patients with a high CHA₂DS₂-VASc score which may have influenced the results of our subgroup analysis. Moreover, we used somewhat arbitrary risk categories to classify patients according to their CHA₂DS₂-VASc score.

Conclusions

Contemporary management of patients with AF across cardiology practices in Europe was associated with an 86% relative risk reduction in ischaemic stroke. CARS and CHA₂DS₂-VASc score may be useful to identify high risk patients despite treatment who may benefit from more aggressive treatment and intensive follow-up. CARS may facilitate future research using real-world cohorts and registries to determine the impact of specific treatments or intervention on the baseline absolute risk of ischaemic stroke in AF.

Supplementary material

Supplementary material is available at *Europace* online.

Acknowledgements

EORP Oversight Committee, Executive, and Steering Committees (National Coordinators) of the EURObservational Research Programme (EORP)—Atrial Fibrillation General Long-Term (EORP-AFGen LT) Registry of the European Society of Cardiology (ESC). Data collection was conducted by the EORP department by Patti-Ann McNeill as Project Officer, Viviane Missiamenou as Data Manager. Overall activities were co-ordinated and supervised by Doctor Aldo P. Maggioni (EORP Scientific Coordinator).

Funding

Since the start of EORP, the following companies have supported the programme: Abbott Vascular Int. (2011–21), Amgen Cardiovascular (2009–18), AstraZeneca (2014–21), Bayer (2009–18), Boehringer Ingelheim (2009–19), Boston Scientific (2009–12), The Bristol Myers Squibb and Pfizer Alliance (2011–16), The Alliance Daiichi Sankyo Europe GmbH and Eli Lilly and Company (2011–17), Edwards (2016–19), Gedeon Richter Plc. (2014–17), Menarini Int. Op. (2009–12), MSD-Merck & Co. (2011–14), Novartis Pharma AG (2014–20), ResMed (2014–16), Sanofi (2009–11), SERVIER (2010–21), Vifor (2019–22).

Conflict of interest: G.B.: small speaker's fees from Medtronic, Boston, Biotronik, Boehringer, and Bayer, outside of the submitted work. F.M.: receiving grants from Ferrer, and personal fees from Bayer, Pfizer/BMS. Boehringer-Ingelheim and Astra-Zeneca outside the submitted work. C.B.L.: receiving grants from Medtronic, Cardiome, and personal fees from Bayer, Sanofi, Boston Scientific, Philips, CathPrint, Organon, and Merck Sharp & Dohme outside the submitted work. T.S.P.: Consultant for Bayer and Pfizer, no fees. L.F.: consultant and speaker activities of small amounts for AstraZeneca, Bayer, BMS/Pfizer, Boehringer Ingelheim, Medtronic, Novartis, Novo, XO and Zoll. G.Y.H.L.: Consultant and speaker for BMS/Pfizer, Boehringer Ingelheim and Daiichi-Sankyo. No fees are received personally. Other authors declare no conflict of interest.

Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

Appendix 1

EURObservational Research Programme Atrial Fibrillation (EORP-AF) Long-Term General Registry Committees and Investigators (ESC Member Countries)

Executive committee: G.Boriani (Chair), G.Y.H. Lip, L. Tavazzi, A. P. Maggioni, G-A. Dan, T. Potpara, M. Nabauer, F. Marin, Z. Kalarus, L. Fauchier.

Steering Committee (National Coordinators): A. Goda, G. Mairesse, T.Shalganov, L. Antoniades, M. Taborsky, S. Riahi, P. Muda, I. García Bolao, O. Piot, M. Nabauer, K. Etsadashvili, E.Simantirakis, M. Haim, A. Azhari, J. Najafian, M. Santini, E. Mirrahimov, K.a Kulzida, A. Erglis, L. Poposka, M. Burg, H. Crijns, Ö. Erküner, D. Atar, R. Lenarczyk, M. Martins Oliveira, D. Shah, G-A. Dan, E. Serdechnaya, T. Potpara, E. Diker, G.Y.H. Lip, D. Lane.

Participants: **ALBANIA** Durrës: E. Zëra, Tirana: U. Ekmekçi, V. Paparisto, M. Tase, Tirana: H. Gjergo, J. Dragoti, A. Goda, **BELGIUM** Bastogne: M. Ciutea, N. Ahadi, Z. el Husseini, M. Raepers, Gilly: J. Leroy, P. Haushan, A. Jourdan, Haine Saint Paul: C. Lepiece, Hasselt: L. Desteghe, J. Vijgen, P. Koopman, G. Van Genechten, H. Heibuchel, Kortrijk: T. Boussy, M. De Coninck, H. Van Eeckhoutte, N. Bouckaert, La Louviere: A. Friart, J. Boreux, C. Arend, Liège: P. Evrard, Liège: L. Stefan, E. Hoffer, J. Herzet, M. Massoz, Liège: C. Celentano, M. Sprynger, L. Pierard, Liège: P. Melon, Overpelt: B. Van Hauwaert, C. Kuppens, D. Faes, D. Van Lie, A. Van Dorpe, Waremmes: A. Gerardy, Yvoir: O. Deceuninck, O. Xhaet, F. Dormal, E. Ballant, D. Blommaert, **BULGARIA** Pleven: D. Yakova, M. Hristov, T. Yncheva, N. Stancheva, S. Tisheva, Plovdiv: M. Tokmakova, F. Nikolov, D. Gencheva, Sofia: T. Shalganov, B. Kunev, M. Stoyanov, Sofia: D. Marchov, V. Gelev, V. Traykov, Varna: A. Kischeva, H. Tsvyatkov, R. Shtereva, S. Bakalska-Georgieva, S. Slavcheva, Y. Yotov, **CZECH REPUBLIC** Ústí nad Labem: M. Kubičková, **DENMARK** Aalborg: A. Marni Joensen, A. Gammelmark, L. Hvilsted Rasmussen, P. Dinesen, S. Riahi, S. Krogh Venø, B. Sorensen, A. Korsgaard, K. Andersen, C. Fragtrup Hellum, Esbjerg: A. Svenningsen, O. Nyvad, P. Wiggers, Herning: O. May, A. Aarup, B. Graversen, L. Jensen, M. Andersen, M. Svejgaard, S. Vester, S. Hansen, V. Lynggaard, Madrid: M. Ciudad, Tallinn: R. Vettus, Tartu: P. Muda, **ESTONIA** Elche, Alicante: A. Maestre, Toledo: S. Castaño, **FRANCE** Abbeville: S. Cheggour, Abbeville: J. Poulard, V. Mouquet, S. Leparrée, Aix-en-Provence: J. Bouet, J. Taieb, Amiens: A. Doucy, H. Duquenne, Angers: A. Furber, J. Dupuis, J. Rautureau, Aurillac: M. Font, P. Damiano, Avignon Cedex: M. Lacrimini, Brest: J. Abalea, S. Boismal, T. Menez, J. Mansourati, Chartres: G. Range, H. Gorka, C. Laure, C. Vassalière, Creteil: N. Elbaz, N. Lellouche, K. Djouadi, Montpellier: F. Roubille, D. Dietz, J. Davy, Nîmes: M. Granier, P. Winum, C. Leperchois-Jacquey, Paris: H. Kassim, E. Marijon, J. Le Heuzey, Paris: J. Fedida, C. Maupain, C. Himbert, E. Gandjbakhch, F. Hidden-Lucet, G. Duthoit, N. Badenco, T. Chastre, X. Waintraub, M. Oudihah, J. Lacoste, C. Stephan, Pau: H. Bader, N. Delarche, L. Giry, Pessac: D. Arnaud, C. Lopez, F. Boury, I. Brunello, M. Lefèvre, R. Mingam, M. Haissaguerre, Rennes: M. Le Bidan, D. Pavin, V. Le Moal, C. Leclercq, Saint Denis: O. Piot, T. Beitar, Saint Etienne: I. Martel, A. Schmid, N. Sadki, C. Romeyer-Bouchard, A. Da Costa, Tours: I. Arnault, M. Boyer, C. Piat, L. Fauchier, **FYR MACEDONIA** Bitola: N. Lozance, S. Nastevska, Ohrid: A. Doneva, B. Fortomarska Milevska, B. Sheshoski, K. Petroska, N. Taneska, N. Bakrecheski, Skopje: K. Lazarovska, S. Jovevska, V. Ristovski,

A. Antovski, Skopje: E. Lazarova, I. Kotlar, J. Taleski, L. Poposka, S. Kedev, Skopje: N. Zlatanovik, Štip: S. Jordanova, T. Bajraktarova Proseva, S. Doncovska, **GEORGIA** Tbilisi: D. Maisuradze, A. Esakia, E. Sagirashvili, K. Lartsuliani, N. Natelashvili, N. Gumberidze, R. Gvenetadze, Tbilisi: K. Etsadashvili, N. Gotonelia, N. Kuridze, Tbilisi: G. Papiashvili, I. Menabde, **GERMANY** Aachen: S. Glöggler, A. Napp, C. Leberherz, H. Romero, K. Schmitz, M. Berger, M. Zink, S. Köster, J. Sachse, E. Vonderhagen, G. Soiron, K. Mischke, Bad Reichenhall: R. Reith, M. Schneider, Berlin: W. Rieker, Biberach: D. Boscher, A. Taschareck, A. Beer, Boppard: D. Oster, Brandenburg: O. Ritter, J. Adamczewski, S. Walter, Chemnitz: A. Frommhold, E. Luckner, J. Richter, M. Schellner, S. Landgraf, S. Bartholome, Chemnitz: R. Naumann, J. Schoeler, Dachau: D. Westermeier, F. William, K. Wilhelm, M. Maerkl, Detmold: R. Oekinghaus, M. Denart, M. Kriete, U. Tebbe, Ebersbach: T. Scheibner, Erlangen: M. Gruber, A. Gerlach, C. Beckendorf, L. Anneken, M. Arnold, S. Lengerer, Z. Bal, C. Uecker, H. Förtsch, S. Fechner, V. Mages, Friedberg: E. Martens, H. Methe, Göttingen: T. Schmidt, Hamburg: B. Schaeffer, B. Hoffmann, J. Moser, K. Heitmann, S. Willems, S. Willems, Hartmannsdorf: C. Klaus, I. Lange, Heidelberg: M. Durak, E. Esen, Itzehoe: F. Mibach, H. Mibach, Kassel: A. Utech, Kirchzarten: M. Gabelmann, R. Stumm, V. Ländle, Koblenz: C. Gartner, C. Goerg, N. Kaul, S. Messer, D. Burkhardt, C. Sander, R. Orthen, S. Kaes, Köln: A. Baumer, F. Dodos, Königsbrück: A. Barth, G. Schaeffer, Leisnig: J. Gaertner, J. Winkler, Leverkusen: A. Fahrig, J. Aring, I. Wenzel, Limburg: S. Steiner, A. Kliesch, E. Kratz, K. Winter, P. Schneider, Ludwigsburg: A. Haag, I. Mutscher, R. Bosch, Markkleeberg: J. Taggeselle, S. Meixner, Meissen: A. Schnabel, Meppen: A. Shamalla, H. Hötze, A. Korinth, Merzig: C. Rheinert, Moosburg: G. Mehlretter, Mühldorf: B. Schön, N. Schön, A. Starflinger, E. Englmann, Munich: G. Baytok, T. Laschinger, G. Ritscher, Munich: A. Gerth, Münster: D. Dechering, L. Eckardt, Nienburg: M. Kuhlmann, N. Proskynitopoulos, Paderborn: J. Brunn, K. Foth, Pirna: C. Axthelm, H. Hohensee, K. Eberhard, S. Turbanisch, Plauen: N. Hassler, A. Koestler, Riesa: G. Stenzel, Riesa: D. Kschwan, M. Schwefer, S. Neiner, S. Hettwer, Rotenburg a.d. Fulda: M. Haeussler-Schuchardt, R. Degenhardt, S. Sennhenn, S. Steiner, Starnberg: M. Brendel, Westerstedde: A. Stoehr, W. Widjaja, S. Loehndorf, A. Logemann, J. Hoskamp, J. Grundt, Zorneding: M. Block, Zwiesel: R. Ulrych, A. Reithmeier, V. Panagopoulos, **ITALY** Bologna: C. Martignani, D. Bernucci, E. Fantecchi, I. Diemberger, M. Ziacchi, M. Biffi, P. Cimaglia, J. Frisoni, G. Boriani, Firenze: I. Giannini, S. Boni, S. Fumagalli, S. Pupo, A. Di Chiara, P. Mirone, Modena: E. Fantecchi, G. Boriani, F. Pesce, C. Zoccali, V.L.Malavasi, **KAZAKHSTAN** Almaty: A. Mussagaliyeva, B. Ahyt, Z. Salihova, K. Koshum-Bayeva, **KYRGYZSTAN** Bishkek: A. Kerimkulova, A. Bairamukova, E. Mirrahimov, **LATVIA** Riga: B. Lurina, R. Zuzans, S. Jegere, I. Mintale, K. Kupics, K. Jubele, A. Erglis, O. Kalejs, **MALTA** Birkirkara: K. Vanhear, M. Burg, M. Cachia, E. Abela, S. Warwicker, T. Tabone, R. Xuereb, **MONTENEGRO** Podgorica: D. Asanovic, D. Drakalovic, M. Vukmirovic, N. Pavlovic, L. Music, N. Bulatovic, A. Boskovic, **NETHERLANDS** Almere: H. Uiterwaal, N. Bijsterveld, Amsterdam: J. De Groot, J. Neefs, N. van den Berg, F. Piersma, A. Wilde, Delfzijl: V. Hagens, Enschede: J. Van Es, J. Van Opstal, B. Van Rennes, H. Verheij, W. Breukers, Heerenveen: G. Tjeerdsma, R. Nijmeijer, D. Wegink, R. Binnema, Hengelo: S. Said, Maastricht: Ö. Erküner, S. Philippens, W. van Doorn, H. Crijns, Rotterdam: T. Szili-Torok, R. Bhagwandien, P. Janse, A. Muskens, s-Hertogenbosch: M. van Eck, R. Gevers, N. van der Ven, Venlo: A. Duygun, B. Rahel, J. Meeder, **NORWAY** Oslo: A. Vold, C. Holst Hansen, I. Engset, D. Atar, **POLAND** Bytom: B. Dyduch-Fejkiowicz, E. Koba, M. Cichocka, Cieszyn: A. Sokal, A. Kubicius, E. Pruchniewicz, Gliwice: A. Kowalik-Sztylec, W. Czapl, Katowice: I. Mróz, M. Kozlowski, T. Pawlowski, M. Tendera, Katowice: A. Winiarska-Filipek, A. Fidyk, A. Slowikowski, M. Haberka, M. Lachor-Broda, M. Biedron, Z. Gasior, Kielce: M. Kolodziej, M. Janion, Kielce: I. Górczyca-Michta, B. Wozakowska-Kaplon, Łódź: M. Stasiak, P. Jakubowski, T. Ciurus,

- J. Drozd, Łódź: M. Simiera, P. Zajac, T. Wcislo, P. Zycinski, J. Kasprzak, Nysa: A. Olejnik, E. Harc-Dyl, J. Miarka, M. Pasieka, M. Ziemińska-Łuć, W. Bujak, Opoczno: A. Śliwiński, A. Grech, J. Morka, K. Petrykowska, M. Prasał, Opole: G. Hordyński, P. Feusette, P. Lipski, A. Wester, Radlin: W. Streb, Rzeszów: J. Romanek, P. Woźniak, M. Chlebuś, P. Szafarz, W. Stanik, Szczecin: M. Zakrzewski, J. Kaźmierczak, Szczecin: A. Przybylska, E. Skorek, H. Błaszczak, M. Stępień, S. Szabowski, W. Krysiak, M. Szymańska, Tarnów: J. Karasiński, J. Blicharz, M. Skura, Warsaw: K. Hałas, L. Michalczyk, Z. Orski, K. Krzyżanowski, A. Skrobowski, Warsaw: L. Zieliński, M. Tomaszewska-Kiecana, M. Dłużniewski, Warsaw: M. Kiliszek, M. Peller, M. Budnik, P. Balsam, G. Opolski, A. Tyminińska, K. Ozierański, A. Wancerz, Warsaw: A. Borowiec, E. Majos, R. Dabrowski, H. Szwed, Zabrze: A. Musialik-Lydk, Zabrze: A. Leopold-Jadczyk, E. Jedrzejczyk-Patej, M. Koziel, R. Lenarczyk, M. Mazurek, Z. Kalarus, Zabrze: K. Krzemien-Wolska, P. Starosta, E. Nowalany-Kozielska, Zakopane: A. Orzechowska, M. Szpot, M. Stasz, **PORTUGAL** Almada: S. Almeida, H. Pereira, L. Brandão Alves, R. Miranda, L. Ribeiro, Carnaxide Lisboa: F. Costa, F. Morgado, P. Carmo, P. Galvao Santos, R. Bernardo, P. Adragão, Santarém: G. Ferreira da Silva, M. Peres, M. Alves, M. Leal, Vila Real: A. Cordeiro, P. Magalhães, P. Fontes, S. Leão, Viseu: A. Delgado, A. Costa, B. Marmelo, B. Rodrigues, D. Moreira, J. Santos, L. Santos, **ROMANIA** Arad: A. Terchet, D. Darabantiu, S. Mercea, V. Turcin Halka, A. Pop Moldovan, Brasov: A. Gabor, B. Doka, G. Catanescu, H. Rus, L. Oboroceanu, E. Bobescu, Bucharest: R. Popescu, A. Dan, A. Buzea, I. Doha, G. Dan, I. Neuhoff, Bucharest: M. Baluta, R. Ploesteanu, N. Dumitrache, M. Vintila, Bucharest: A. Daraban, C. Japie, E. Badila, H. Tewelde, M. Hostiuc, S. Frunza, E. Tintea, D. Bartos, Bucharest: A. Ciobanu, I. Popescu, N. Toma, C. Gherghinescu, D. Cretu, N. Patrascu, C. Stoicescu, C. Udriou, G. Bicescu, V. Vintila, D. Vinereanu, M. Cinteza, R. Rimbis, Iași: M. Grecu, Oradea: A. Cozma, F. Boros, M. Ilie, O. Tica, R. Tor, A. Corina, A. Jeewooth, B. Maria, C. Georgiana, C. Natalia, D. Alin, D. Dinu-Andrei, M. Livia, R. Daniela, R. Larisa, S. Umaar, T. Tamara, M. Ioachim Popescu, Târgu Mureș: D. Nistor, I. Sus, O. Coborosan, Timișoara: N. Alina-Ramona, R. Dan, L. Petrescu, Timișoara: G. Ionescu, I. Popescu, C. Vacarescu, E. Goanta, M. Mangea, A. Ionac, C. Mornos, D. Cozma, S. Pescariu, **RUSSIAN FEDERATION** Arkhangelsk: E. Solodovnicova, I. Soldatova, J. Shutova, L. Tjuleneva, T. Zubova, V. Uskov, Arkhangelsk: D. Obukhov, G. Rusanova, Arkhangelsk: I. Soldatova, N. Isakova, S. Odinsonova, T. Arhipova, Arkhangelsk: E. Kazakevich, E. Serdechnaya, O. Zavyalova, Saint-Petersburg: T. Novikova, Saint-Petersburg: I. Riabaia, S. Zhigalov, Saint-Petersburg: E. Drozdova, I. Luchkina, Y. Monogarova, Vladivostok: D. Hegya, L. Rodionova, L. Rodionova, V. Nevzorova, Vladivostok: I. Soldatova, O. Lusanova, **SERBIA** Belgrade: A. Arandjelovic, D. Toncev, L. Vukmirovic, M. Radisavljevic, M. Milanov, N. Sekularac, Belgrade: M. Zdravkovic, S. Hinic, S. Dimkovic, T. Acimovic, J. Saric, S. Radovanovic, Belgrade: A. Kocijancic, B. Obrenovic-Kircanski, D. Kalimanovska Ostric, D. Simic, I. Jovanovic, I. Petrovic, M. Polovina, M. Vukicevic, M. Tomasevic, N. Mujovic, N. Radivojevic, O. Petrovic, S. Aleksandric, V. Kovacevic, Z. Mijatovic, B. Ivanovic, M. Tesic, T. Potpara, A. Ristic, B. Vujsic-Tesic, M. Nedeljkovic, Belgrade: A. Karadzic, A. Uscumlic, M. Prodanovic, M. Zlatar, M. Asanin, Belgrade: B. Bisenic, V. Vasic, Z. Popovic, Belgrade: D. Djikic, M. Sipic, V. Peric, B. Dejanovic, N. Milosevic, Belgrade: S. Backovic, A. Stevanovic, A. Andric, B. Pencic, M. Pavlovic-Kleut, V. Celic, Kragujevac: M. Pavlovic, M. Petrovic, M. Vuleta, N. Petrovic, S. Simovic, Z. Savovic, S. Milanov, G. Davidovic, V. Iric-Cupic, Niš: D. Djordjevic, M. Damjanovic, S. Zdravkovic, V. Topic, D. Stanojevic, M. Randjelovic, R. Jankovic-Tomasevic, V. Atanaskovic, S. Antic, M. Pavlovic, Niška Banja: D. Simonovic, M. Stojanovic, S. Stojanovic, V. Mitic, V. Ilic, D. Petrovic, M. Deljanin Ilic, S. Ilic, V. Stoickov, Pirot: S. Markovic, Šabac: A. Mijatovic, D. Tanasic, D. Petrovic, G. Radakovic, J. Peranovic, M. Pavlovic, N. Panic-Jelic, O. Vujadinovic, P. Pajic, S. Bekic, S. Kovacevic, **SPAIN** Alicante: A. García Fernandez, Benalmadena: A. Perez Cabeza, Córdoba: M. Anguita, Granada: L. Tercedor Sanchez, Huarte: E. Mau, J. Loayssa, M. Ayarra, M. Carpintero, Madrid: I. Roldán Rabadan, Murcia: M. Leal, Murcia: M. Gil Ortega, Murcia: A. Tello Montoliu, E. Orenes Piñero, S. Manzano Fernández, F. Marín, A. Romero Aniorte, A. Veliz Martínez, M. Quintana Giner, Pamplona: G. Ballesteros, M. Palacio, O. Alcalde, I. García-Bolao, San Juan de Alicante: V. Bertomeu Gonzalez, Santiago de Compostela: F. Otero-Raviña, J. García Seara, J. Gonzalez Juanatey, **SWITZERLAND** Geneva: N. Dayal, P. Maziarski, P. Gentil-Baron, D. Shah, **TURKEY** Adana: M. Koç, Afyon: E. Onrat, I. E. Dural, Ankara: K. Yilmaz, B. Özün, Ankara: S. Tan Kurklu, Y. Atmaca, Ankara: U. Canpolat, L. Tokgozoglu, Ankara: A. K. Dolu, B. Demirtas, D. Sahin, Ankara: O. Ozcan Celebi, E. Diker, Antalya: G. Gagirci, Bayraklı/Izmir: U.O.Turk, Bursa: H. Ari, Diyarbakır: N. Polat, N. Toprak, Gaziantep: M. Sucu, Görükle-Bursa: O. Akin Serdar, Istanbul: A. Taha Alper, Istanbul: A. Kepez, Istanbul: Y. Yuksel, Kurupelit - Samsun: A. Uzunseli, S. Yuksel, M. Sahin, Merkez/Düzce: O. Kayapinar, Mersin: T. Ozcan, Sivas: H. Kaya, M. B. Yilmaz, Trabzon: M. Kutlu, Yüreğir-Adana: M. Demir, **UNITED KINGDOM** Barnstaple: C. Gibbs, S. Kaminskiene, M. Bryce, A. Skinner, G. Belcher, J. Hunt, L. Stancombe, B. Holbrook, C. Peters, S. Tetterzell, Birmingham: A. Shantsila, D. Lane, K. Senoo, M. Proietti, K. Russell, P. Domingos, S. Hussain, J. Partridge, R. Haynes, S. Bahadur, R. Brown, S. McMahon, G. Y H Lip, Blackburn: J. McDonald, K. Balachandran, R. Singh, S. Garg, H. Desai, K. Davies, W. Goddard, Blackpool: G. Galasko, I. Rahman, Y. Chua, O. Payne, S. Preston, O. Brennan, L. Pedley, C. Whiteside, C. Dickinson, J. Brown, K. Jones, L. Benham, R. Brady, Carlisle: L. Buchanan, A. Ashton, H. Crowther, H. Fairlamb, S. Thornthwaite, C. Relph, A. McKean, U. Poultney, N. Kelsall, P. Rice, T. Wilson, Chertsey: M. Wrigley, R. Kaba, T. Patel, E. Young, J. Law, Cramlington: C. Runnett, H. Thomas, H. McKie, J. Fuller, S. Pick, Exeter: A. Sharp, A. Hunt, K. Thorpe, C. Hardman, E. Cusack, L. Adams, M. Hough, S. Keenan, A. Bowring, J. Watts, Great Yarmouth: J. Zaman, K. Goffin, H. Nutt, Harrogate: Y. Beerachee, J. Featherstone, C. Mills, J. Pearson, L. Stephenson, Huddersfield: S. Grant, A. Wilson, C. Hawke, I. Alam, M. Robinson, S. Ryan, Macclesfield: R. Egdell, E. Gibson, M. Holland, D. Leonard, Maidstone: B. Mishra, S. Ahmad, H. Randall, J. Hill, L. Reid, M. George, S. McKinley, L. Brockway, W. Milligan, Manchester: J. Sobolewska, J. Muir, L. Tuckis, L. Winstanley, P. Jacob, S. Kaye, L. Morby, Nottingham: A. Jan, T. Sewell, Poole: C. Boos, B. Wadams, C. Cope, P. Jefferey, Portsmouth: N. Andrews, A. Getty, A. Suttliff, C. Turner, K. Hudson, R. Austin, S. Howe, Redhill: R. Iqbal, N. Gandhi, K. Brophy, P. Mirza, E. Willard, S. Collins, N. Ndlovu, Rhy: E. Subkovas, V. Karthikeyan, L. Waggett, A. Wood, A. Bolger, J. Stockport, L. Evans, E. Harman, J. Starling, L. Williams, V. Saul, Salisbury: M. Sinha, L. Bell, S. Tudgay, S. Kemp, J. Brown, L. Frost, Shrewsbury: T. Ingram, A. Loughlin, C. Adams, M. Adams, F. Hurford, C. Owen, C. Miller, D. Donaldson, H. Tivenan, H. Button, South Shields: A. Nasser, O. Jhagra, B. Stidolph, C. Brown, C. Livingstone, M. Duffy, P. Madgwick, Southampton: P. Roberts, E. Greenwood, L. Fletcher, M. Beveridge, S. Earles, Taunton: D. McKenzie, D. Beacock, M. Dayer, M. Seddon, D. Greenwell, F. Luxton, F. Venn, H. Mills, J. Rewbury, K. James, K. Roberts, L. Tonks, Torquay: D. Felmeden, W. Taggu, A. Summerhayes, D. Hughes, J. Sutton, L. Felmeden, Watford: M. Khan, E. Walker, L. Norris, L. O'Donohoe, Weston-super-Mare: A. Mozid, H. Dymond, H. Lloyd-Jones, G. Saunders, D. Simmons, D. Coles, D. Cotterill, S. Beech, S. Kidd, Wolverhampton: B. Wrigley, S. Petkar, A. Smallwood, R. Jones, E. Radford, S. Milgate, S. Metherell, V. Cottam, Yeovil: C. Buckley, A. Broadley, D. Wood, J. Allison, K. Rennie, L. Balian, L. Howard, L. Pippard, S. Board, T. Pitt-Kerby.

References

1. Chugh SS, Havmoeller R, Narayanan K, Singh D, Rienstra M, Benjamin EJ et al. Worldwide epidemiology of atrial fibrillation: a global burden of disease 2010 study. *Circulation* 2014;**129**:837–47.
2. Burdett P, Lip GYH. Atrial fibrillation in the United Kingdom: predicting costs of an emerging epidemic recognising and forecasting the cost drivers of atrial fibrillation-related costs. *Eur Hear J Qual Care Clin Outcomes* 2022;**8**:187–94.
3. Lip GYH, Nieuwlaat R, Pisters R, Lane DA, Crijns HJGM. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: The Euro Heart Survey on atrial fibrillation. *Chest* 2010;**137**: 263–72.
4. Hindricks G, Potpara T, Dagres N, Arbelo E, Bax JJ, Blomström-Lundqvist C et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association of Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2021;**42**:373–498.
5. Chao TF, Joung B, Takahashi Y, Lim TW, Choi EK, Chan Y-H et al. 2021 Focused update consensus guidelines of the Asia Pacific heart rhythm society on stroke prevention in atrial fibrillation: executive summary. *Thromb Haemost* 2022;**122**:20–47.
6. Potpara TS, Lip GYH, Blomstrom-Lundqvist C, Boriani G, Van Gelder IC, Heidbuchel H et al. The 4S-AF scheme (stroke risk; symptoms; severity of burden; substrate): a novel approach to in-depth characterization (rather than classification) of atrial fibrillation. *Thromb Haemost* 2021;**121**:270–8.
7. Ding WY, Proietti M, Boriani G, Fauchier L, Blomström-Lundqvist C, Marin F et al. Clinical utility and prognostic implications of the novel 4S-AF scheme to characterize and evaluate patients with atrial fibrillation: a report from ESC-EHRA EORP-AF Long-Term General Registry. *EP Europace* 2022;**24**:721–8.
8. Romiti GF, Pastori D, Rivera-Caravaca JM, Ding WY, Gue YX, Menichelli D et al. Adherence to the “atrial fibrillation better care” pathway in patients with atrial fibrillation: impact on clinical outcomes-A systematic review and meta-analysis of 285,000 patients. *Thromb Haemost* 2022;**122**:406–14.
9. Kirchhof P, Camm AJ, Goette A, Brandes A, Eckardt L, Elvan A et al. Early rhythm-control therapy in patients with atrial fibrillation. *N Engl J Med* 2020;**383**:1305–16.
10. Ding WY, Calvert P, Gupta D, Huisman M V, Lip GYH. Impact of early ablation of atrial fibrillation on long-term outcomes: results from phase II/III of the GLORIA-AF registry. *Clin Res Cardiol* 2022;**111**:1057–68.
11. Lee CJY, Toft-Petersen AP, Ozenne B, Phelps M, Olesen JB, Ellinor PT et al. Assessing absolute stroke risk in patients with atrial fibrillation using a risk factor based approach. *Eur Hear J Cardiovasc Pharmacother* 2021;**9**:f3–f10.
12. Boriani G, Proietti M, Laroche C, Fauchier L, Marin F, Nabauer M et al. Contemporary stroke prevention strategies in 11 096 European patients with atrial fibrillation: a report from the EURObservational Research Programme on Atrial Fibrillation (EORP-AF) long-term general registry. *Europace* 2018;**20**:747–57.
13. Ding WY, Rivera-Caravaca JM, Marin F, Torp-Pedersen C, Roldán V, Lip GYH. Prediction of residual stroke risk in anticoagulated patients with atrial fibrillation: mCARS. *J Clin Med* 2021;**10**:3357.
14. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;**44**:837–45.
15. Hart RG, Pearce LA, Aguilar MI. Meta-analysis: antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med* 2007;**146**: 857–67.
16. Ding WY, Gupta D, Lip GYH. Atrial fibrillation and the prothrombotic state: revisiting Virchow's triad in 2020. *Heart* 2020;**106**:1463–8.
17. Ruff CT, Giugliano RP, Braunwald E, Hoffman EB, Deenadayalu N, Ezekowitz MD et al. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet* 2014;**383**: 955–62.
18. Ding WY, Lane DA, Gupta D, Huisman MV, Lip GYH. Incidence and risk factors for residual adverse events despite anticoagulation in atrial fibrillation: results from phase II/III of the GLORIA-AF registry. *J Am Heart Assoc* 2022;**11**:e026410.
19. Ding WY. Residual stroke risk in atrial fibrillation. *Arrhythmia Electrophysiol Rev* 2021;**10**: 147–53.
20. Ding WY, Rivera-Caravaca JM, Marin F, Li G, Roldán V, Lip GYH. Number needed to treat for net effect of anticoagulation in atrial fibrillation: real-world vs. clinical-trial evidence. *Br J Clin Pharmacol* 2022;**88**:282–9.