Aalborg Universitet



The iodine content of Faroese food items to support iodine nutrition in the North Atlantic

Johannesen, Herborg Líggjasardóttir; Andersen, Stig; Weihe, Pál; Petersen, Maria Skaalum; Strøm, Marin; Veyhe, Anna Sofía

Published in: International Journal of Circumpolar Health

DOI (link to publication from Publisher): 10.1080/22423982.2023.2233145

Creative Commons License CC BY-NC 4.0

Publication date: 2023

Document Version Publisher's PDF, also known as Version of record

Link to publication from Aalborg University

Citation for published version (APA):

Johannesen, H. L., Andersen, S., Weihe, P., Petersen, M. S., Strøm, M., & Veyhe, A. S. (2023). The iodine content of Faroese food items to support iodine nutrition in the North Atlantic. International Journal of Circumpolar Health, 82, Article 2233145. https://doi.org/10.1080/22423982.2023.2233145

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: July 03, 2025





International Journal of Circumpolar Health

ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/zich20

The iodine content of Faroese food items to support iodine nutrition in the North Atlantic

Herborg Líggjasardóttir Johannesen, Stig Andersen, Pál Weihe, Maria Skaalum Petersen, Marin Strøm & Anna Sofía Veyhe

To cite this article: Herborg Líggjasardóttir Johannesen, Stig Andersen, Pál Weihe, Maria Skaalum Petersen, Marin Strøm & Anna Sofía Veyhe (2023) The iodine content of Faroese food items to support iodine nutrition in the North Atlantic, International Journal of Circumpolar Health, 82:1, 2233145, DOI: 10.1080/22423982.2023.2233145

To link to this article: https://doi.org/10.1080/22423982.2023.2233145

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



0

View supplementary material 🖸

đ	1	ſ	1	
С				
⊢	н	_		

Published online: 21 Jul 2023.



🕼 Submit your article to this journal 🗗





View related articles 🗹



View Crossmark data 🗹

RESEARCH ARTICLE

OPEN ACCESS OPEN ACCESS

The iodine content of Faroese food items to support iodine nutrition in the North Atlantic

Herborg Líggjasardóttir Johannesen ^{a,b}, Stig Andersen^{c,d,e}, Pál Weihe^{b,f,g}, Maria Skaalum Petersen^{b,f,g}, Marin Strøm^{b,f} and Anna Sofía Veyhe^{b,f,g}

^aDepartment of Endocrinology and Medicine, The National Hospital of the Faroe Islands, Torshavn, Faroe Islands; ^bCenter of Health Science, Faculty of Health Science and Nursing, University of the Faroe Islands, Torshavn, Faroe Islands; ^cDepartment of Clinical Medicine, Aalborg University, Aalborg, Denmark; ^dArctic Health Research Centre, Aalborg University Hospital, Aalborg, Denmark; ^eGreenland Centre for Health Research, University of Greenland, Nuuk, Greenland; ^fCenter of Health Science, University of the Faroe Islands, Torshavn, Faroe Islands; ^gDepartment of Occupational Medicine and Public Health, The Faroese Hospital System, Torshavn, Faroe Islands

ABSTRACT

lodine is essential for human health, and dietary iodine sources are critical. Living isolated on islands in the North Atlantic Ocean fostered unique dietary traditions in the Faroe Islands. The dietary peculiarities led us to collect and measure the iodine content of local foods and beverages and calculate daily iodine intakes based on four Faroese dietary scenarios. Marine foods and birds had iodine contents from 300 μ g/kg and up, markedly higher than dairy, grains, terrestrial, vegetables, and liquid foods (p<0.001). Vegetables and beverages contained negligible iodine. Wind-dried fermented lamb meat iodine content varied by 100% depending on whether it was herded on an island with beaches (105 μ g/kg) or with cliffs (53 μ g/kg) facing the ocean. The daily iodine intake varied from around 50 μ g to just over 200 μ g with diets rich in fish and whales. In conclusion, the traditional diet has been a critical source of iodine on the Faroe Islands. The Faroese may be iodine replete with traditional diets but face the risk of iodine deficiency with dietary transition towards a more western diet. Locally produced Faroese foods are therefore relevant to promote health, and may additionally support food security, local involvement and knowledge, and long-term sustainability.

Introduction

lodine is an element that is essential for human growth and development [1] and plays a vital role in the occurrence of thyroid disorders [2]. Human iodine status is dependent on dietary sources, and the iodine content of food items determines the individual subject's iodine intake [3–5]. lodine-rich foods include marine mammals and fish, and milk and other dairy products [6,7]. Additionally, tap water may contribute to the iodine intake in some areas [8].

The Faroe Islands are a group of 18 islands in the North Atlantic Ocean, with no single spot being more than 5 kilometres from the sea [9]. The proximity to the sea may influence local foods' iodine content, and we hypothesise that the traditional Faroese foods are iodine rich. Local food items are those produced and consumed in the Faroe Islands, including food items that are sourced from both land and sea, and are a part of the traditional, local Faroese diet. A growing local movement is the initiative to explore the possibilities for the Faroese to be self-supplied. Thus, the word of the year in 2022 in the Faroe Islands was "stuttflutt", meaning "locally produced". Local productions are ample, and the Faroese cuisine includes local, traditional foods such as fermented and wind-dried lamb meat, seabirds, fish, and Faroese milk [10]. Pilot whale hunting, by driving and communal sharing, is an old tradition that was cherished for centuries and still is [11]. However, recommendations are to limit the intake of both whale meat and blubber or even not to eat it at all due to the content of environmental pollutants [12,13].

Food tables for iodine are lacking in the Faroe Islands, and foreign records such as the Danish database "Frida" [14] have been used for estimating intake from Faroese foods. One study based on diet interviews estimated mean iodine intake in the Faroe Islands at 244 μ g/day [15]. Faroese teenagers were iodine replete [16], while urine iodine concentration was 86 μ g/L in a cross-sectional study of Faroese women raising a concern for iodine

CONTACT Herborg Líggjasardóttir Johannesen Arighels.fo Department of Endocrinology and Medicine, The National Hospital Faroe Islands, J.c. Svabosgøta, Torshavn, FO 100, Faroe Islands

Supplemental data for this article can be accessed online at https://doi.org/10.1080/22423982.2023.2233145.

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

ARTICLE HISTORY

Received 23 January 2023 Revised 22 June 2023 Accepted 30 June 2023

KEYWORDS

lodine; traditional food; Faroe Island; source of iodine; nutrition; food security



This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (http://creativecommons.org/licenses/by-nc/4.0/), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

deficiency even in areas surrounded by the sea [17]. Thus, uncertainties have emerged about the iodine intake in the Faroese population and, particularly, the iodine content in Faroese food. These uncertainties led us to measure the iodine content in local food products. Based on the measured iodine content, we calculated the iodine intake from four Faroese dietary scenarios.

Methods

Foods and beverage samples were collected or purchased in local stores in November 2021. We kept food items frozen in iodine-free containers until analysis. We sampled two dummy tests and 89 Faroese food items: 34 dairy products, 15 marine-, 13 terrestrial-, 9 beverages-, 7 vegetables-, and 11 additional samples, as listed in Table 1. Dairy products included milk, yoghurt, cream, sour cream, cheese, and butter. Repeated sampling of milk and cream was performed to evaluate between-day fluctuations. Marine samples included local food items sold in stores: cod, halibut, and salmon. Local hunters supplied meat and blubber from pilot whale, fermented lamb, seabirds (common guillemot and puffin), and hare. We sampled lamb from farmers on three islands to assess the impact of geography. The islands were Hestur, Streymoy (Leynar), and Eysturoy (Glyvrar). We also measured iodine in samples from the lamb's organs: heart, kidney, liver, blood, tallow, and rolled seasoned meat (a sausage made of rolled mutton). We sampled wheat and rye bread from a local bakery, made home-baked rye bread (by H.L.J.), pizza, and traditional dry biscuit. We included five different types of locally purchased salt. The selection of Faroese vegetables, herbs and fruits is sparse, and only a few locally grown products are available. We sampled potatoes, carrots, brassica species, rhubarb jam, and parsley. Finally, we sampled beverages comprising three beers, one cider and three soft drinks, all of which were produced locally.

Samples were blinded by random order labelling before they were shipped for analysis to keep the laboratory blinded to the food items during sample handling, iodine measurement, and reporting of test results. Samples included two dummy samples: one of water with negligible iodine contents subsequently spiked with iodine as described elsewhere [18], and one with Icelandic lamb, with median iodine content of $32-52 \mu g/kg (0-127 \mu g/kg)$ [19].

Definition of food groups and calculation of iodine intake

A recent report described food classification for the UK biobank and recommended fifteen main food categories (13 food and two beverage categories) [20]. To support

our calculation of the iodine intake based on a daily dietary intake from the included foods in this study we chose to categorise the foods into six groups of local foods and one with beverages. These were logical to the food items available: marine-, seabirds, meat, grains, dairy products, offal, vegetables-, and salt, and beverages. We included a group of other food items ("other") when there was no or minimal commercial Faroese production of the food item. The individual samples' iodine content is shown in supplementary table S1.

We listed four typical Faroese dietary scenarios for the calculation of the iodine intakes of an individual using the local products. The meals were composed based on Faroese dietary reports for adults [10,15,21,22]. We based the calculation on the consumption of 600-800 grams of foods in three main meals and a daily fluid intake of 1600 ml, as listed in Table 2. The breakfast comprised of bread, butter, cheese, rhubarb jam, milk, and water for all four days. A biscuit with butter and cheese was included for a snack. Lunch comprised of bread, butter, kohlrabi pickled, and water all four days, and toppings of liver pate, sausage roll, wind-dried fermented sheep for two days, and fishcakes and salmon for two days. Dinners were either a meat meal with fermented sheep, kohlrabi raw and potato; cod twice with different milk consumptions, or whale meat and blubber with increasing the milk intake. These Faroese dietary scenarios can hence be summarised as follows: breakfast followed by four different food patterns for the day with (1) Meat dinner and Meat lunch; (2) Fish dinner and Meat lunch; (3) Fish dinner and Fish lunch; or (4) Whale dinner and Fish lunch. Details are listed in Table 2, and the iodine content of Faroese foods was multiplied by the intake of each food item and beverage. The sum of iodine contents of the food intakes added up to the calculated total daily iodine intake.

Assay procedures

The iodine concentration in food items was determined by the Sandell-Kolthoff reaction modified after Wilson and van Zyl [23]. The principle is detecting the catalytic role of iodine in reducing ceric ammonium sulphate in the presence of arsenous acid after pretreatment with alkaline ashing. The pre-treatment protocol was modified for conducting the measurement in solid items. The iodine laboratory at Aalborg University Hospital (Aalborg, Denmark) has vast experience with these procedures [24,25], and is included in the Ensuring the Quality of Urinary Iodine Procedures (EQUIP) program (Department of Health and Human Services, CDC, USA). We use a rigorous and validated

protocol for measuring the iodine content of solid and liquid items [6]. The measurement unit for iodine content in solid products was micrograms per kilogram $(\mu g/kg)$ and micrograms per litre $(\mu g/L)$ in liquids. Measuring iodine was complicated for certain solid samples with very high iodine concentrations. In these samples, we chose to provide the valid result, which was "high jodine concentration" reported as $>320 \mu g/kg$. The coefficient of variation for liquids at our laboratory is 9.2% (interval 1–4 μ g/L, n = 8); 8.7% (interval 5–9 μ g/L, n = 4); 4.2% (interval 10–15 μ g/L, n =4); 1.5% (interval 15–50 μ g/L, n = 5); 1.5% at higher levels after dilution of samples [18]. The analytical coefficient of variation was determined for solid items with the Faroese food items: 20.3% (interval 10–45 μ g/ kg, n = 8; 8.7% (interval 50-85 µg/kg, n = 5); 7.5% (interval 100–400 μ g/kg, n = 3). For solid food items we managed measurements up to 400 µg/kg, but for most samples with high iodine contents we the highest values were reported validly to be above 320 µg/kg.

The iodine concentration in the dummy samples were according to expected values. The water sample had low iodine content identical to a previous measurement [18], and the solid item (Icelandic lamb meat) had an iodine content as reported in Icelandic nutrient tables.

Statistics

The statistical analyses were performed in the R environment for statistical computing (version 4.2.1) [26,27]. Data are presented with numbers, percentages, mean, minimum, maximum, median, and interquartile range. We tested data for normality by visually inspection of QQ-plots. The model assumptions were that the observations from different food groups were independent and the individual observation within each group was independent. Differences between food groups were calculated using the One-way ANOVA. A p-value <0.05 indicated a statistically significant difference.

Results

Marine food items from the North Atlantic Ocean had high iodine contents of around 300 μ g/kg or higher. Beverages, including tap water, had consistently low iodine contents of <2 μ g/L. Figure 1 illustrates the differences in iodine content between food groups.

Table 1 lists the iodine contents of the collected Faroese food items. Marine foods had markedly higher iodine contents than other foods (p < 0.001), and the

iodine content was distinctly higher than terrestrial foods (p < 0.001). Seabirds had iodine contents mirroring that measured in fish. Store-bought bread and pizza had iodine contents above 100 µg/kg, while other grains were lower in iodine content (Table 1, supplementary table S1).

The median iodine concentration of sixteen samples of four kinds of milk collected over one month was 67 μ g/L [IQR 12], the lowest at 39 μ g/L and the highest at 89 μ g/L. The daily variation was limited with no time trend (p > 0.1). The iodine content was similar between the different types of milk products (p = 0.1). Other dairy products ranged from 42 to 189 μ g/L.

As for terrestrial foods, lamb had an average iodine content of 56 μ g/kg, while it was only 12 μ g/kg in vegetables. The five local salt products varied from 9–16 μ g/kg, with "Gestus Cornwall England Salt" being an exception having an iodine content of 72 μ g/kg. Finally, all nine beverages, brewed or produced based on the local water, had negligible iodine contents (<2 μ g/L).

Lamb meat iodine content varied by 100% depending on whether it was herded on an island with beaches (Figure 2) or with cliffs facing the ocean (Table 1). The four Faroese diets with varying consumption of meat and fish for main dishes, adding side dishes and drinks, are listed in Table 2. The results indicate a total daily iodine intake that spans from 50 μ g/day for meals based on terrestrial meat to above 200 μ g/day when meals were based on marine dishes (Table 2).

Discussion

These are the first Faroese data on iodine content of Faroese food items. Our study showed that Faroese marine foods were iodine rich, followed by grain and dairy products, while terrestrial food items had low iodine contents. In addition, local water-based products and vegetables had negligible iodine content. Our findings corroborate and details previous reports that marine foods generally have a higher iodine content than terrestrial food items [6,14]. The present data are interesting as maintaining life on the small, remote islands in the North Atlantic Ocean requires adaptation to the local environment. Such adaptation is reflected in dietary peculiarities, including fermented and wind-dried lamb meat and pilot whale, hunting and driving, contributing to both local culture and to the iodine intake.

Seabirds' high iodine content is interesting although not surprising, as they often feed on small fish. It illustrates that living on the Faroe Islands requires an adaptation to local opportunities and that these are related

4 😉 H. L. JOHANNESEN ET AL.

Food groups		Mean					
		Food items	Samples	μg/kg or <i>μg/L</i>	Min	Max	
Dairy products	Milk and yoghurt	Milk ^a	16	67	39	89	
, ,	, ,	Sour cream and yoghurt ^b	6	89	42	189	
		Cream ^c	7	49	39	64	
	Cheese	Butter	3	28	24	35	
		Cheese ^d	2	70			
Marine products	Fish products	Cod ^e	2	>320			
	·	Salmon	3	>320		403	
		Halibut	1	>320			
		Fish balls (homemade) ^f	1	>320			
		Fish balls (commercial)	1	84			
		Fish sausage	1	48			
	Whale	Whale meat ^g	1	>320			
		Whale blubber ^h	1	>320			
	Seaweed	Seaweed	2	>320			
	Sea birds	Common guillemont ⁱ (fat)	1	>320			
		Puffin (meat)	1	244			
Terrestrial products	Meat	Sausage roll (lamb)	2	25			
		Fermented meat (lamb)	3	76	53	105	
		Hare	1	20			
	Offal	Heart (lamb)	1	19			
		Kidney (lamb)	1	64			
		Liver (lamb) ^j	2	39			
		Blood (lamb)	1	125			
		Tallow ^e	2	21			
Grains		Bread ^k	2	178			
		Rve bread (home-baked)	1	39			
		Non-sweet biscuit	1	27			
		Pizza	1	128			
Vegetables	Vegetables	Kohlrabi root (raw)	1	12			
		Kohlrabi root (pickled)	1	8			
		Carrot (raw)	1	12			
	Potatoes	Potato (raw)	2	14			
		Potato (boiled)	1	23			
	Other	Parsley (domestic-grown)	1	209			
		Rhubarb iam	1	5			
Salt		Salt	5	24	9	72	
Beverages		Beer	3	1	-	, 2	
		Sparkling water	3	0			
		Cider	1	0			
		Liquor	2	0			

Table 1. lodine-content food items produced in the Faroe Islands.

^aIncludes four types of fat content: 0.1%; 0.5%; 1.5%; 3.5%. ^bIncludes sour cream and plain & fruity yoghurt. ^cFat content 38%. ^dOne type of cheese. ^e Fresh and fermented. ^fContent is nearly 100% fresh fish. ^gDried whale meat. ^hSalted whale blubber. ⁱUria aalge. ^jPaté and raw liver. ^kCommercially produced wheat and rye.

to the sea. Hence, the birds on the remote islands support life by using sea-based food substances, which contribute to the high iodine content of the birds. This finding may be a parallel to human nutrition in the same environment in the North Atlantic.

We previously reported that the frequency of fish meals was associated with urinary iodine concentration among Faroese [17]. People living in harsh coastal areas, an environment unfavourable to terrestrial food production, often rely on a diet dominated by marine food items. This is the case for people living on islands in the North Atlantic, such as the Faroes, Shetland, Orkney Islands, and Iceland. All face changes in dietary habits [21,22,28,29] with an influx of junk food, western-style foods, and restaurants. The fish intake decreased from 70 to 40 grams per day between 1982 and 2001 [22], and recent data illustrated an even lower fish

consumption today, especially among younger generations [21].

Fishcakes and cold cuts are mixed products containing a fillet of fish, often haddock, milk powder, cornflour, onion, herbs, and water. Adding water, spices, onion, and cornflour to marine food items lowered the iodine content of the composite dietary component. However, fishcakes are popular and seem to support the iodine intake in this population, where a dietary transition has implied a decrease in fish intake over the past decades [21].

Seaweed is a growing industry in the Faroe Islands and an upcoming food gaining popularity. Therefore, we sampled seaweed from the local beach and found a high iodine concentration, in line with the previous reports of seaweed from the North Atlantic [30,31]. Seaweed has an excessive iodine content, but a study

	Meal	lodine content	Meat dinner Meat lunch		Fish dinner Meat lunch		Fish dinner Fish lunch		Whale dinner Fish lunch	
		μg/kg	g	μg	g	μg	g	μg	g	μg
Foods	Cod	307	-		150	46.1	150	46.1	-	
	Salmon	400 ^a	-		-		25	10.0	25	10.0
	Fishcake	84	-		-		20	1.7	20	1.7
	Whale meat	400 ^a	-		-		-		175	70.0
	Whale blubber	400 ^a	-		-		-		50	20.0
	Bread	178 ^b	145	25.8	145	25.8	145	25.8	145	25.8
	Non-sweet biscuit	27	25	0.7	25	0.7	25	0.7	25	0.7
	Butter	28	35	1.0	35	1.0	35	1.0	35	1.0
	Cheese	70	45	3.2	45	3.2	45	3.2	45	3.2
	Liver pate	47	20	0.9	20	0.9	-		-	
	Sausage roll	35	13	0.5	13	0.5	13	0.5	-	
	Wind dried sheep (cold cut)	79	20	1.6	20	1.6	20	1.6	20	1.6
	Fermented sheep (dinner)	50	150	7.5	-		-		-	
	Salt	24	8	0.2	8	0.2	8	0.2	8	0.2
	Kohlrabi (raw)	12	20	0.2	20	0.2	20	0.2	20	0.2
	Kohlrabi (pickled)	8	20	0.2	20	0.2	20	0.2	20	0.2
	Carrot	12	20	0.2	20	0.2	20	0.2	20	0.2
	Potato boiled	23	180	4.1	180	4.1	180	4.1	180	4.1
	Rhubarb jam	5	20	0.1	20	0.1	20	0.1	20	0.1
	Total intake		721	46.2	721	84.8	746	95.6	808	139.0
	Drinks	μg/L	mL	μg	mL	μg	mL	μg	mL	μg
Drinks	Tap water	0	1350	0.0	1350	0.0	1000	0.0	400	0.0
	Sparkling water	0	-		-		200	0.0	-	
	Milk	67	50	3.4	50	3.4	400	26.8	1000	67.0
	Beer	0	200	0.0	200	0.0	-		200	0.0
	Total drink intake		1600	3.4	1600	3.4	1600	26.8	1600	67.0
Estimated	lodine from foods:			46.2		84.8		95.6		139.0
iodine	lodine from drinks			3.4		3.4		26.8		67.0
intake	Total iodine			49.6		88.2		122.4		206.0

Table 2. Estimated iodine intake based on four days of consumption of Faroese-produced food products, each column represents one day. The total intake of food ("g"), and the iodine intake, ("microg").

^alodine content in the food product was >320 μg/kg (see Table 1) and therefore an estimated iodine concentration is provided. ^bBread includes rye- and wheat bread and the iodine content is average one rye bread and one wheat bread sample (see Supplementary Table S1).

showed no short-term adverse effects on thyroid function [32], and seaweed intake may support long-term iodine nutrition when the intake of other marine foods decreases.

Our results indicate that not only marine animals, but also other Faroese foods contribute to the general iodine status, including dairy products and grain. The Faroese have a habit of eating relatively high amounts of bread [21], and our data showed a relatively high median iodine content in grains of 106 µg/kg [IQR 139]. Furthermore, salt contributes to the iodine intake. There is no legislation on salt iodisation in the Faroe Islands, but most salt is imported from Denmark, where iodisation of table and industrial salt is mandatory [33]. Our findings conform to the use of iodised industrial salt for commercial bread production, and the central bakeries confirm the use of iodine-fortified salt. Both iodised and non-iodised salt are available in the stores, albeit all the salt products in our study contained iodine. Hence, our findings corroborate that iodised salt was available in stores to be used for cooking, though the use of iodised salt in our study may have led to a slight overestimation of iodine intake.

Locally produced dairy products had a mean iodine concentration of $67 \mu q/L$ with no apparent time trend. Herds are largely kept indoors all year, and therefore we would not expect a seasonal variation in iodine content of milk. Our findings were lower than those from Norway, reporting a seasonal variation with the higher median of 232 µg/L in winter [7], and iodine fortification of animal fodder since the 1950s [34]. There is only one dairy plant in the Faroe Islands, which receives milk from multiple farms. Thus, dairy products on the market are always pooled, and we do not have data on the individual milk farms. However, our findings show that the iodine content in Faroese milk was 67 µg/kg, lower than the 100 μ g/kg [14] in Denmark and 160 μ g/kg in Norway [35]. This finding suggests an evaluation of the need for iodine fortification of the fodder on Faroese farms.

The Faroese terrestrial diet is mainly lamb. We found that iodine in lamb meat varied widely, comparable to findings from Iceland [19]. We add that this is due to sheep in "Streymoy" having access to the sea and eating seaweed (Figure 2), unlike lamb from "Hestur", a rocky island with steep cliffs and no beaches. This illustrates an importance of proximity to the ocean.



Figure 1. Mean iodine contents of groups of 89 Faroese foods to portray the relative importance of items for iodine intake among people living in isolated islands in the North Atlantic Ocean.



Figure 2. Picture a sheep in the Faroe Islands searching for food on the coast and eating seaweed. It was printed with permission from copyright holder Bjarki G. Dalsgarð.

Tap water iodine is essential for iodine intake in some countries [36,37], but our results show that drinking water does not contribute to iodine intake in the Faroe Islands [17]. Similarly, beverages produced locally all had negligible iodine content and did not contribute to iodine nutrition.

Overall, data on Faroese food intake is sparse, but the major food groups are dairy products, grains, potatoes, meats, vegetables, and fish [21]. Based on our results, a healthy iodine intake relies on daily intake of marine, grain, and dairy products, while the contribution is limited when considering vegetables and beverages other than milk. We used the consumption and iodine content of Faroese foods to calculate a total daily iodine intake. Interestingly, it is possible to meet the recommended iodine intake based on only locally based Faroese foods. Our calculated iodine intake of a local Faroese diet showed that a diet combining terrestrial foods and a low intake of dairy products causes moderate iodine deficiency with an estimated intake of just under 50 µg/day. This is improved by swapping terrestrial with marine foods or adding dairy products and bread to the diet. The combination of seafood, bread and dairy foods is estimated to cause an intake slightly above the recommended range. Hence, Faroese may be iodine replete from local foods, but there is a risk of iodine deficiency when turning away from local marine foods.

The Faroese strive for self-supply, and local food production may support sustainability with limited transport and improve regional knowledge and involvement. Additionally, it may promote food security while supporting iodine intake. These possibilities are made explicit by the remoteness of the Faroe Islands, which highlights the opportunities for others to follow.

It is a strength that the study reports the iodine content of most Faroese foods available and hence portrays the local diet of these islands in the North Atlantic Ocean. More samples could be added to strengthen the validity of our results, but the variation between samples included was limited. Hence, the results provide a valid estimate of the iodine contents. We relied on previous dietary surveys and food frequency questionnaires to determine typical serving sizes for the Faroese population. It is important to note that while the literature on food intake is helpful for estimating dietary intake, it does have limitations and uncertainties. Also, some uncertainty remains on the iodine content of foods with very high iodine content. Still, we demonstrate that Faroese foods may provide both deficient and high iodine intakes depending on the use of a traditional diet.

We could not estimate seasonal variation, which has been reported from other areas with different dietary habits. Thus, this could be an interesting path to pursue in a future study.

Conclusion

The study reported iodine content in food items from the Faroe Island, and calculated iodine intake from traditional Faroe fare. The traditional Faroese diet could provide sufficient iodine since it contains marine foods, rich in iodine. Hence, conscientious food choices may enable the Faroese to be iodine replete without supplementation. Raising awareness may nudge the population towards this ambition, but this requires health promotion initiatives, legislation, and enforcement to promote self-supply. The ongoing dietary changes emphasise the need for such initiatives to improve professional and public knowledge of the possibilities for sustainable and healthy nutrition and the need for continuous monitoring of population iodine status. Presently, dietary changes emphasise a need to use iodised salt even with the local diet to ensure adequate intake. Future research should aim also to promote understanding the relationship between dietary habits and iodine status in the Faroe Islands.

Acknowledgments

Thanks to the hunters for donating samples from their animals. Thanks to Mjólkarvirkið Búnaðarmanna for assisting with the sampling of milk and cream. We acknowledge Eydfinn Lenvig for assessing the size of traditional Faroese dishes used for calculating iodine intakes from Faroese meals. Special thanks to photographer Bjarki G. Dalsgarð for permission to use the photo.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

The Research Council at the Faroese National hospital (H.L.J., grant number 21/00010-36); North Denmark Region (S.A.); Betri Stuðul (H.L.J. grant number 1021).

Authorship

Principal investigators were H.L.J. and A.S.V. The author's contributions were the following: H.L.J., S.A., and A.S. V. sparked the idea, initiated the study, contributed to the design and implementation of the survey, ensured the writing and publishing processes, and contributed to the preparation of the manuscript. S.A. collaborates and assists with chemical analyses. H.L.J. and A.S.V. did the data analyses, and H.L.J., and S.A., were grant holders. All authors contributed to the interpretation of data and read and commented on the final draft.

ORCID

Herborg Líggjasardóttir Johannesen i http://orcid.org/0000-0002-9048-3303

References

- Eastman CJ, Zimmermann MB The lodine deficiency disorders. *Endotext*. Published online February 6, 2018.
 [cited 2022 May 2]. Available from: https://www.ncbi.nlm. nih.gov/books/NBK285556/
- [2] Laurberg P, Cerqueira C, Ovesen L, et al. lodine intake as a determinant of thyroid disorders in populations. Best Pract Res Clin Endocrinol Metab. 2010;24(1):13–27. doi: 10.1016/J.BEEM.2009.08.013
- [3] Andersen S, Waagepetersen R, Laurberg P. Misclassification of iodine intake level from morning spot urine samples with high iodine excretion among Inuit and non-Inuit in Greenland. Br J Nutr. 2015; 113 (9):1433–1440. doi: 10.1017/S0007114515000653
- [4] Carlsen MH, Andersen LF, Dahl L, et al. New lodine food composition database and updated calculations of iodine intake among Norwegians. Nutrients. 2018;10 (7):930. doi: 10.3390/NU10070930
- [5] Andersen S, Hvingel B, Kleinschmidt K, et al. Changes in iodine excretion in 50-69-y-old denizens of an Arctic society in transition and iodine excretion as a biomarker of the frequency of consumption of traditional Inuit foods. Am J Clin Nutr. 2005;81(3):656–663. doi: 10.1093/AJCN/81.3.656
- [6] Andersen S, Hvingel B, Laurberg P. International journal of circumpolar health iodine content of traditional Greenlandic food items and tap water in East and West Greenland. Int J Circumpolar Health. 2002;61(4):332–340. Published online 2002. doi: 10.3402/ijch.v61i4.17491
- [7] Dahl L, Opsahl JA, Meltzer HM, et al. lodine concentration in Norwegian milk and dairy products. Br J Nutr. 2003;90 (3):679–685. doi: 10.1079/BJN2003921
- [8] Andersen S, Pedersen KM, Iversen F, et al. Naturally occurring iodine in humic substances in drinking water in Denmark is bioavailable and determines population

iodine intake. British Journal Of Nutrition. 2008;99 (2):319–325. doi: 10.1017/S0007114507803941

- [9] Visit Faroe Islands. Accessed January. 2. 2023;https:// www.62n.fo/travel/en/faroe-islands/
- [10] Joensen JP. Bót Og Biti (Stock Cube and a Bite). Torshavn, The Faroe Islands: Fróðskapur, Faroese University Press; 2015.
- [11] Tinganes. Whales and whaling in the Faroe Islands. https://www.whaling.fo/en/
- [12] Weihe P, Joensen HD. Dietary recommendations regarding pilot whale meat and blubber in the Faroe Islands. Int J Circumpolar Health. 2012;71(1):1–5. doi: 10.3402/ IJCH.V71I0.18594
- [13] Starvsstovan H. Kosttilmæli Um at Eta Grind (Dietary Recommendation for Consuming Pilot Whale). Recommendation no. 461-200800444-64.Heilsufrøðiliga Starvsstovan. Torshavn, Faroe Islands: Faroese Food and Veterinary Authority; 2011.
- [14] Frida Database med fødevaredata udgivet DTU Fødevareinstituttet. [cited 2022 July 2]. Available from: https://frida.fooddata.dk/
- [15] Vestergaard Z. Føðslukanning 1981-1982 (Dietary Investigation). Fróðskaparrrit (Annales Societatis Scientiarum Færoensis). 1987;33:5–18.
- [16] Johannesen HL, Andersen S, Andersen SL, et al. Iodine nutrition with North Atlantic living: the Faroese adolescents. J Nutr Sci. 2023;12:e17. doi: 10.1017/jns.2022.111
- [17] Johannesen HL, Knudsen GS, Andersen S, et al. lodine nutrition among the adult population of the Faroe Islands: a population-based study. British Journal Of Nutrition. 2022;127(8):1190–1197. doi: 10.1017/ S0007114521001938
- [18] Veyhe AS, Johannesen HL, Weihe P, et al. Urine test strips and iodine contamination: a tricky trick in iodine nutrition surveys. Scand J Clin Lab Invest. 2022;82(3):251–256. doi: 10.1080/00365513.2022.2059699
- [19] Reykdal Ó, Hilmarsson ÓÞ, Auðunsson GA Joð, Selen Og Kvikasilfur í Kjöti, Mjólk Og Eggjum (lodine, Selenium And Mercury In Meat, Milk And Eggs). 2006. [cited 2022 December 8]. Available from: https://www.matis.is/ media/utgafa/matra/Jod_Lokaskyrsla_0804.pdf
- [20] Piernas C, Perez-Cornago A, Gao M, et al. Describing a new food group classification system for UK biobank: analysis of food groups and sources of macro- and micronutrients in 208,200 participants. Eur J Nutr. 2021;60(5):2879–2890. doi: 10.1007/s00394-021-02535-x
- [21] Arctic Monitoring Assessment Program (AMAP). AMAP Assessment 2021: Human Health in the Arctic | AMAP. 2021. [cited 2022 November 25]. Available from: https:// www.amap.no/documents/doc/amap-assessment-2021human-health-in-the-arctic-pre-print/3593
- [22] Veyhe AS. Færøske kvinders kostvaner i tredje trimester. MPH 2006. Dnr U12/3:294. Göteborg, Sweden: Nordic School of Public Health NHV; 2006.
- [23] Wilson B, van Zyl A. The estimation of iodine in thyroidal amino acids by alkaline ashing. S Afr J Med Sci. 1967;32 (3):70–82.
- [24] Laurberg P. Thyroxine and 3,5,3'-triiodothyronine content of thyroglobulin in thyroid needle aspirates in

hyperthyroidism and hypothyroidism. J Clin Endocrinol Metab. 1987;64(5):969–974. doi: 10.1210/JCEM-64-5-969

- [25] Andersen S, Karmisholt J, Pedersen KM, et al. Reliability of studies of iodine intake and recommendations for number of samples in groups and in individuals. British Journal Of Nutrition. 2008;99(4):813–818. doi: 10.1017/ S0007114507842292
- [26] R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [Published 2020. citrd 2023 January 2]. Available from: https://www.R-project.org
- [27] RStudio Team. Rstudio: Integrated Development for R. Boston, MA. Published 2019. [cited 2023 January 2]. Available from: http://www.rstudio.com
- [28] Gunnarsdottir I, Gustavsdottir AG, Steingrimsdottir L, et al. lodine status of pregnant women in a population changing from high to lower fish and milk consumption. Public Health Nutr. 2013;16(2):325–329. doi: 10.1017/ S1368980012001358
- [29] Adalsteinsdottir S, Tryggvadottir EA, Hrolfsdottir L, et al. Insufficient iodine status in pregnant women as a consequence of dietary changes. Food Nutr Res. 2020;64:64. doi: 10.29219/fnr.v64.3653
- [30] Grandorf Bak U. Seaweed cultivation in the Faroe Islands: An investigation of the biochemical composition of selected macroalgal species, optimised seeding technics, and open-ocean cultivation methods from a commercial perspective. Kgs Lyngby, Denmark: Technical University of Denmark; 2019.
- [31] Andersen S, Noahsen P, Rex KF, et al. lodine in edible seaweed, its absorption, dietary use, and relation to lodine Nutrition in Arctic People. J Med Food. 2019;22 (4):421–426. doi: 10.1089/jmf.2018.0187
- [32] Noahsen P, Kleist I, Larsen HM, et al. Intake of seaweed as part of a single sushi meal, iodine excretion and thyroid function in euthyroid subjects: a randomized dinner study. J Endocrinol Invest. 2020;43(4):431–438. doi: 10. 1007/s40618-019-01122-6
- [33] Laurberg P, Jørgensen T, Perrild H, et al. The Danish investigation on iodine intake and thyroid disease, DanThyr: status and perspectives. Eur J Endocrinol. 2006;155(2):219–228. doi: 10.1530/EJE.1.02210
- [34] Madar AA, Heen E, Hopstock LA, et al. lodine Intake in Norwegian Women and Men: The Population-Based Tromsø Study 2015–2016. Nutrients. 2020;12(11):1–13. doi: 10.3390/NU12113246
- [35] The Norwegian Food Safety Authority. Matvaretabellen. [cited 2023 January 11]. Available from: https://matvare tabellen.no/
- [36] Amelsvoort V. Rural water-supply development and the recent appearance of endemic goitre. Trop Geogr Med. 1971;23(3):304–305.
- [37] Manousou S, Stål M, Eggertsen R, et al. Correlations of water iodine concentration to earlier goitre frequency in Sweden - an iodine sufficient country with long-term iodination of table salt. Environ Health Prev Med. 2019;24(1). doi: 10.1186/s12199-019-0821-9