

World human neutrophil antigens investigation survey

Bayat, Behnaz; Lowack, Jonas; Audrain, Marie; Croisille, Laure; Curtis, Brian; Dangerfield, Rebecca; Esmaeili, Behnaz; Grabowski, Claudia; Keller, Margaret; Kim, Hyungsuk; Kroll, Hartmut; Kvanka, Marjeta Macek; Kwok, Janette; Moritz, Elyse; Nathalang, Oytip; Nelson, Derrick; Nielsen, Kaspar René; Pahn, Gail; Poles, Anthony; Porcelijn, Leendert; Sachs, Ulrich J.; Schönbacher, Marlies; Körmöczy, Günther F.; Kupatawintu, Pawinee; Takahashi, Daisuke; Uhrynowska, Malgorzata; Flesch, Brigitte; Fung, Yoke-Lin

Published in:
Vox Sanguinis

DOI (link to publication from Publisher):
[10.1111/vox.13500](https://doi.org/10.1111/vox.13500)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Bayat, B., Lowack, J., Audrain, M., Croisille, L., Curtis, B., Dangerfield, R., Esmaeili, B., Grabowski, C., Keller, M., Kim, H., Kroll, H., Kvanka, M. M., Kwok, J., Moritz, E., Nathalang, O., Nelson, D., Nielsen, K. R., Pahn, G., Poles, A., ... Fung, Y.-L. (2023). World human neutrophil antigens investigation survey. *Vox Sanguinis*, 118(9), 763-774. <https://doi.org/10.1111/vox.13500>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: December 05, 2025

World human neutrophil antigens investigation survey

Behnaz Bayat¹  | Jonas Lowack¹ | Marie Audrain² | Laure Croisille³ | Brian Curtis⁴ | Rebecca Dangerfield⁵ | Behnaz Esmaeili^{6,7} | Claudia Grabowski⁸ | Margaret Keller⁵ | Hyungsuk Kim⁹ | Hartmut Kroll⁸ | Marjeta Macek Kvanka¹⁰ | Janette Kwok¹¹ | Elyse Moritz¹²  | Oytip Nathalang¹³  | Derrick Nelson¹⁴ | Kaspar René Nielsen¹⁵ | Gail Pahn¹⁶ | Anthony Poles¹⁷ | Leendert Porcelijn¹⁸ | Ulrich J. Sachs¹ | Marlies Schönbacher¹⁹ | Günther F. Körmöcz¹⁹ | Pawinee Kupatawintu²⁰ | Daisuke Takahashi²¹ | Malgorzata Uhrynowska²² | Brigitte Flesch²³ | Yoke-Lin Fung²⁴ 

¹Institute for Clinical Immunology, Transfusion Medicine and Haemostasis, Justus-Liebig-University, Gießen, Germany

²Service d'Immunologie, Laboratoire de Biologie, Nantes, France

³Laboratoire HLA-ILP E.F.S Ile-de-France, Créteil, France

⁴Platelet & Neutrophil Immunology Lab, Versiti, Milwaukee, Wisconsin, USA

⁵American Red Cross Neutrophil Immunology Laboratory, St Paul, Minnesota, USA

⁶Department of Basic Medical Sciences, Khoy University of Medical Sciences, Khoy, Iran

⁷Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

⁸Institut für Transfusionsmedizin Dessau, DRK-Blutspendedienst NSTOB, Dessau, Germany

⁹Seoul National University Hospital, Seoul, South Korea

¹⁰Blood Transfusion Centre of Slovenia, Ljubljana, Slovenia

¹¹Division of Transplantation and Immunogenetic, Department of Pathology, Queen Mary Hospital, Pok Fu Lam, Hong Kong

¹²Clinical and Experimental Oncology, Escola Paulista de Medicina, Universidade Federal de Sao Paulo, Sao Paulo, Brazil

¹³Faculty of Allied Health Sciences, Thammasat University, Pathumtani, Thailand

¹⁴Specialized Laboratory Services, South African National Blood Service, Johannesburg, South Africa

¹⁵Klinisk Immunologi, Aalborg Universitetshospital Nord, Aalborg, Denmark

¹⁶Platelet & Granulocyte Reference Laboratory, Australian Red Cross Lifeblood, Brisbane, Australia

¹⁷Department of Histocompatibility & Immunogenetics, NHS Blood & Transplant, Bristol, UK

¹⁸Platelet/Leucocyte Serology Laboratory, Sanquin Diagnostic Services, Amsterdam, The Netherlands

¹⁹Department of Transfusion Medicine and Cell Therapy, Medical University of Vienna, Vienna, Austria

²⁰National Blood Centre, Thai Red Cross Society, Bangkok, Thailand

²¹Japanese Red Cross Society Central Blood Institute, Tokyo, Japan

²²Institute of Hematology and Transfusion Medicine, Warszawa, Poland

²³Laboratory for Immunogenetics/HLA, DRK Blutspendedienst West, Bad Kreuznach, Germany

²⁴School of Health, University of the Sunshine Coast, Sunshine Coast, Queensland, Australia

Behnaz Bayat and Yoke-Lin Fung contributed equally to this study.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 Institute of Clinical Immunology and Transfusion Medicine and The Authors. *Vox Sanguinis* published by John Wiley & Sons Ltd on behalf of International Society of Blood Transfusion.

Correspondence

Behnaz Bayat, Institute for Clinical Immunology, Transfusion Medicine and Haemostasis, Justus-Liebig-University, Langhansstrasse 7, 35392, Giessen, Germany. Email: behnaz.bayat@immunologie.med.uni-giessen.de

Funding information

The authors received no specific funding for this work.

Open Access funding enabled and organized by Projekt DEAL.

Abstract

Background and Objectives: Human neutrophil antigens (HNAs) are categorized into five systems: HNA-1 to HNA-5. Given the importance of neutrophils in immunity, we sought to create awareness of the role of HNA diagnostic services in managing immune neutropenia and transfusion-related acute lung injury. To provide health communities all around the world with access to these services, we conducted a survey to create a directory of these HNA diagnostic services.

Materials and Methods: An Excel table-based survey was created to capture information on the laboratory's location and was emailed to 55 individuals with known or possible HNA investigation activity. The collected data were then summarized and analysed.

Results: Of contacted laboratories, the surveys were returned from 23 (38.2%) laboratories; 17 have already established HNA diagnostic (of them 12 were regular participants of the International Granulocyte Immunobiology Workshop [ISBT-IGIW]), 4 laboratories were in the process of establishing their HNA investigation and the remaining 2 responder laboratories, did not conduct HNA investigations. In established laboratories, investigation for autoimmune neutropenia (infancies and adults) was the most frequently requested, and antibodies against HNA-1a and HNA-1b were the most commonly detected.

Conclusion: The directory of survey respondents provides a resource for health professionals wanting to access HNA diagnostic services. The present study offers a comprehensive picture of HNA diagnostics (typing and serology), identifying weak points and areas for improvement for the first time. Identifying more laboratories involved in HNA diagnostics with limited access to international societies in the field will globally improve HNA diagnostics.

Keywords

alloantibody, autoantibody, autoimmune, HNA, neutropenia, neutrophil, TRALI

Highlights

- Investigation for autoimmune neutropenia (infancies and adults) is the most frequently requested analysis for neutrophil serology worldwide.
- Antibodies against human neutrophil antigen (HNA)-1a and HNA-1b are the most commonly detected antibodies involved in autoimmune neutropenia.
- Although neutrophil serology services are distributed across the globe, the number of services available to African, Middle Eastern and Western Pacific populations is limited.

INTRODUCTION

Neutrophils are the major subtype of granulocytes and have a pivotal role in innate and adaptive immunity. Human neutrophil antigens (HNAs) are distributed on five different glycoproteins on the surface of neutrophils and have been designated as HNA-1 to HNA-5 [1]. Eleven HNA alleles have been described so far [2]. HNA incompatibilities during pregnancy or transfusion may induce the production of HNA antibodies [3]. HNA antibodies have been implicated in the mechanism of neonatal alloimmune neutropenia (NAIN), autoimmune

neutropenia (AIN), transfusion-related acute lung injury (TRALI), immune neutropenia after bone marrow transplant and febrile non-haemolytic transfusion reactions [2, 4].

Neutropenia may arise from a defect in neutrophil production in bone marrow or increased neutrophil destruction for various reasons [5]. Immune neutropenia involves the attachment of neutrophil reactive antibodies to neutrophil antigens, leading to their clearance from circulation [6]. Depending on the cause, treatment for neutropenia varies. It is therefore important to distinguish immune neutropenia from other causes. Therefore, the availability of services to detect

TABLE 1A Directory of participating established laboratories.

	Region	HNA investigating laboratories and contact details	QA (quality assessment)
1	Americas	Dr Elyse Moritz, email: elysemoritz@yahoo.com.br Clinical and Experimental Oncology, Escola Paulista de Medicina, Universidade Federal de Sao Paulo, Brazil.	ISBT-IGIW
2		Rebecca Dangerfield, email: rebecca.dangerfield@redcross.org ; Margaret Keller, email margaret.keller@redcross.org American Red Cross Neutrophil Immunology Laboratory, St Paul, Minnesota, USA.	ISBT-IGIW
3		Dr Brian Curtis, email: brian.curtis@bcw.edu Platelet & Neutrophil Immunology Lab, Versiti, Milwaukee, WI, USA.	ISBT-IGIW
4	Europe	Dr Marlies Schoenbacher, email: marlies.schoenbacher@meduniwien.ac.at Dr Günther Körmöcz, email: guenther.koermoecki@meduniwien.ac.at Department of Transfusion Medicine and Cell Therapy, Medical University of Vienna, Austria	INSTAND EQA
5		Dr Kaspar René Nielsen, email: k.nielsen@rn.dk Klinisk immunologi, Aalborg universitetshospital Nord, Denmark.	ISBT-IGIW
6		Dr Laure Croisille, email: laure.croisille@efs.sante.fr Laboratoire HLA-ILP E.F.S Ile-de-France, Créteil, France.	ISBT-IGIW
7		Dr Marie Audrain, email: marie.audrain@chu-nantes.fr Service d'Immunologie, Laboratoire de Biologie, Nantes, France.	ISBT-IGIW and INSTAND EQA
8		Dr Harmut Kroll, email: hartmut.kroll@bsd-nstob.de Dr Claudia Grabowski, email: claudia.grabowski@bsd-nstob.de Institut für Transfusionsmedizin Dessau, DRK-Blutspendedienst NSTOB, Dessau, Germany.	INSTAND EQA
9		Dr Ulrich Sachs, email: ulrich.sachs@med.uni-giessen.de Dr Behnaz Bayat, email: behaz.bayat@immunologie.med.uni-giessen.de Institute for Clinical Immunology and Transfusion Medicine, Giessen, Germany.	ISBT-IGIW and INSTAND EQA
10		Dr Leendert Porcelijn, email: l.porcelijn@sanquin.nl Platelet/Leucocyte Serology Laboratory, Sanquin Diagnostic Services, Amsterdam, The Netherlands.	ISBT-IGIW
11		Dr Malgorzata Uhrynowska, email: muhrynowska@ihit.waw.pl Institute of Haematology and Transfusion Medicine: Warszawa, Poland.	ISBT-IGIW
12		Dr Anthony Poles, email: anthony.poles@nhsbt.nhs.uk Department of Histocompatibility & Immunogenetics, NHS Blood & Transplant, Bristol, United Kingdom.	ISBT-IGIW
13		Dr Marjeta Macek Kvanka, email: marjeta.macek@ztn.si Blood transfusion centre of Slovenia, Ljubljana, Slovenia	ISBT-IGIW
14		Dr Oytip Nathalang, email: oytipntl@hotmail.com Faculty of Allied Health Sciences, Thammasat University, Pathumtani 12120, Thailand.	
15	Western Pacific	Gail Pahn, email: gpahn@redcrossblood.org.au Platelet & Granulocyte Reference Laboratory, Australian Red Cross Lifeblood, Brisbane, Australia.	ISBT-IGIW
16		Dr Janette Kwok, email: kwoksy@ha.org.hk Queen Mary Hospital, Division of Transplantation and Immunogenetics, Department of Pathology, Hong Kong SAR.	INSTAND EQA
17		Dr Daisuke Takahashi, email: d-takahashi@jrc.or.jp Japanese Red Cross Society Central Blood Institute, Tokyo, Japan.	

anti-neutrophil antibodies enables the physician to confirm immune neutropenia thus avoiding the invasive bone marrow biopsy [7]. This is particularly useful in cases of infant and childhood neutropenia. The implication of HNA antibodies, particularly anti-HNA-3a in clinically severe TRALI has also significantly emphasized the importance of HNA antibody detection.

The International Society of Blood Transfusion (ISBT)-Granulocyte Immunology Working Party (GIWP) organizes an annual quality assessment workshop to assess a laboratory's ability to carry out neutrophil serology and genotyping investigations. There are currently only 18 reference laboratories participating in this quality assessment workshop, but we expect that there are other laboratories performing similar investigations around the world as well. Given the significant role of neutrophils in immunity, we conducted a survey to identify and catalogue laboratories around the globe performing neutrophil serology and genotyping investigations. The goal is to create a worldwide directory of granulocyte investigating laboratories conducting neutrophil serology and genotyping investigations to increase awareness of these diagnostic services and to make these specialized services more accessible to health communities all around the world.

MATERIALS AND METHODS

The survey consisted of an Excel table that captured information on the laboratory's location, the range of neutrophil investigations (e.g., AIN, TRALI, neutrophil reactive antibodies in transplants and convalescent plasma of COVID-19 patients) conducted, list of techniques used for serology and molecular investigations, and whether they participated in a quality assessment programme(s) in the 12-month period from January 2019 to January 2020. The survey was emailed to 55 individuals with known or possible HNA investigation activity, and recipients were encouraged to forward the survey to any other laboratories that may conduct HNA investigations. A literature search (keywords: HNA frequency, human neutrophil antigens, neutrophils, neutrophil antigen 'country') was conducted to gather reports on HNA frequencies from different populations.

RESULTS

A total of 23 surveys were returned equating to a return rate of 41.8%. From the responses received, 17 established laboratories regularly conducted HNA investigations, 3 (17.6%) are in the Americas, 10 (58.8%) in Europe, 1 in South-East Asia (5.9%) and 3 (17.6%) in the Western Pacific area (Table 1A). Of these 17 laboratories, 15 (88.2%) participated in a granulocyte quality assessment program, the most common ($n = 11$) being the International Society of Blood Transfusion-International Granulocyte Immunobiology Workshop (ISBT-IGIW) and the other being the INSTAND External Quality Assessment (EQA) programme (Table 1A). Four institutes (National Blood Centre Thailand, South Africa, Iran and South Korea) are in the process of establishing their HNA investigation services and have

been clustered as 'Laboratories in development' (Table 1B). The remaining two responses did not conduct HNA investigations: the New Zealand Blood Transfusion Service referred their investigations to Australia, and Dr Olnaiyi Olanrewaju from the Irrua Specialist Teaching Hospital, Irrua/Ambrose Alli University, Ekpoma, Edo State Nigeria reported that they had cases of immune neutropenia but did not have a laboratory to refer the samples to.

Serological testing was conducted by 16 of 17 established laboratories and all 4 laboratories in development (Table 2) but the range of techniques varied. All 16 laboratories from first group and 1 from second group conducted granulocyte immunofluorescence tests (GIFT) and used a typed panel of granulocytes. All these laboratories complemented the GIFT with the granulocyte agglutination test (GAT) or another technique, except in Aalborg and Créteil. LabScreen Multi was used by 6 established laboratories and 3 of 4 laboratories in development (Table 2) [8]. Monoclonal antibody immobilization of granulocyte antigen (MAIGA) [9] was conducted by 12 established laboratories, with Sao Paulo and Tehran in the process of optimizing the assay. All laboratories investigating CD16 used two monoclonal antibodies (mAb) except for USA, Versiti. MEM-166 was the most common CD177 mAb and Bear 1 for CD11b. Sixteen established laboratories and two laboratories in development conducted genotyping for HNA-1, HNA-3, HNA-4 and HNA-5, but only five from first group also genotyped for HNA-2 (Table 3).

NAIN investigations were conducted by 15 established laboratories (Table 4). Of the samples tested in the survey period, 88 samples were positive, and the most common antibodies detected were anti-HNA-1a and anti-HNA-1b. Samples were referred by physicians from

TABLE 1B Directory of participating laboratories in development.

1	Africa	Dr Derrick Nelson, email: derrick.nelson@sanbs.org.za Specialized Laboratory Services, South African National Blood Service, Johannesburg, South Africa.
2	Eastern Mediterranean	Dr Esmaeili Behnaz, email: esmaeili.behnaz@yahoo.com Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran.
3	W.Pac	Dr Hyungsuk Kim, email: hyungsuk.kim79@gmail.com Seoul National University Hospital, Seoul, Korea.
4	South East Asia	Dr Pawinee Kupatawintu, email: pawinee.k@redcross.or.th Dr Atthapol Srisuddee, email: atthapol.s@redcross.or.th National Blood Centre, Thai Red Cross Society, Bangkok, Thailand.

Abbreviations: EQA, External Quality Assessment; HNA, human neutrophil antigens; ISBT-IGIW, International Society of Blood Transfusion-International Granulocyte Immunobiology Workshop.

TABLE 2 Human neutrophil antigen (HNA)-antibodies screening assay and monoclonal antibodies (mAb) used in monoclonal antibody immobilization of granulocyte antigen (MAIGA).

Serology	GIFT	GAT	Typed panel	No. typed cells	Laboratory screen	Other techniques	MAIGA	Monoclonal antibodies used in MAIGA					HLA class I
								CD16	CD177	CD11a (LFA-1α)	CD11b	CD18	
1 Brazil, Sao Paulo	Y	Y	Y	3	Y		inP	3G8 LNK16	MEM166		Bear1		
2 USA, ARC	Y	Y	Y	5			Y	3G8 LNK16 DJ130c BCW238.7		7D8		IB4	
3 USA, Versiti	Y	?	Y	2-10			Y	MBC238.7		7D8			
4 Austria, Vienna	Y	Y	Y	3		WIFT	Y	3G8 LNK16 DJ130c	MEM-166	HI111	Bear1	7E4	W6/32
5 Denmark, Aalborg	Y	Y	Y	4		D-GIFT							
6 France, Créteil	Y	Y	Y	3-6			Y	3G8 LNK16	MEM166				
7 France, Nantes	Y	Y	Y	4			Y	3G8 LNK16	MEM-166	HI111	Bear1		
8 Germany, Dessau	Y	Y	Y	4			Y	3G8 LNK16 DJ130c	MEM166	7D8	25.3.1	Bear1	B1G6
9 Germany, Giessen	Y	Y	Y	4			Y	3G8 LNK16	MEM166	7D8		Bear1	W6/32
10 The Netherlands, Sanquin	Y	Y	Y	2			Y	3G8 LNK16 DJ130c	238,7	MEM166	7D8	IB4	
11 Poland, Warsaw	Y	Y	Y	2	Y		Y	3G8 LNK16 DJ130c	MEM166		Bear1	7E4	
12 United Kingdom, NHS	Y	Y	Y	2+	Y	rHNA-3a/3b cell lines, GCLT, LIFT	Y	3G8 LNK16 MBC238.7	MEM166	7D8	Bear1	7E4	
13 Slovenia, Ljubljana	Y	Y	Y	3+	Y		Y						
14 Thailand, Thammasat University	N	N	N	N	N		N						
15 Australia, Lifeblood	Y	Y	Y	3			Y	3G8 LKN16 DJ130c	MEM166	25.3.1	Bear1	W6/32	
16 Hong Kong, Queen Mary	Y	Y	Y	3	Y		N	3G8 LNK16 DJ130	MEM154	3H1029			
17 Japan, JRC	Y	Y	Y	3-5	Y	ICFA	N						
Laboratories in development													
1 South Africa, SANBS					Y		N						

(Continues)

TABLE 2 (Continued)

Serology	GIFT	GAT	Typed panel	No. typed cells	Laboratory screen	Monoclonal antibodies used in MAIGA							HLA class I
						Other techniques	MAIGA	CD16	CD177	CD11a (LFA-1α)	CD11b	CD18	
2	Iran, Tehran	Y	Y	2		LIFT	inP	LNK16	MEM166				
3	Korea, Seoul				Y		N						
4	Thailand, Red Cross				Y		N						

Abbreviations: GAT, granulocyte agglutination test; GCLT, granulocyte chemiluminescence test; GIFT, granulocyte immunofluorescence test; HNA, human neutrophil antigen; ICFA, immunocomplex capture fluorescence analysis; LIFT, lymphocyte immunofluorescence test; MAIGA, monoclonal antibody immobilization of granulocyte antigen; WIFT, white cell immunofluorescence test.

patients with neutropenia (absolute neutrophil count [ANC] < 1500 cells/μL for adults and ANC < 1000 per microliter). The number of investigations for AIN was higher and the positive rates ranged from 9% to 52.2% (Table 5). Requests for AIN investigations were mainly for patients aged 2–36 months and in adults (Table 5). Only 16 established laboratories conducted TRALI investigations, and the number of samples tested in each laboratory varied considerably (Table 6). The majority of TRALI-associated antibodies were in donor samples and only three established laboratories detected antibodies in the patient.

DISCUSSION

The survey results show that the distribution of HNA investigation capability among laboratories is varied. There is a large cluster of laboratories in Europe (58.8%), three in America (17.6%), three in Western Pacific (17.6%) and one in South East Asia (5.8%; Tables 1A and 1B). There are presently no established laboratories in Africa or Eastern Mediterranean, but there is a laboratory in development in Tehran, Iran.

NAIN is a disorder of the foetus that results from maternal alloimmunization against paternal HNA expressed on fetal neutrophils [4, 10]. NAIN is analogous to haemolytic disease of the foetus/newborn (HDFN) which affects red blood cells and foetal/neonatal alloimmune thrombocytopenia (FNAIT) which affects platelets [11]. In NAIN, the mother develops alloantibodies against HNAs expressed on foetal neutrophils that may cause elimination of neutrophils leading to neutropenia in foetus [11]. Of the 17 established laboratories that regularly investigate NAIN, the majority (58.8%) are located in Europe, and there are none in Africa or Easter Mediterranean (Tables 1A and 1B). The glycoproteins carrying epitopes for HNA-1 and HNA-2 are the most frequently involved in the NAIN cases in this study. Seven laboratories identified alloantibodies against HNA-2 as the second most frequent alloantibodies involved in NAIN cases. In the reports from four laboratories (Creteil and Nantes in France, Sanquin in the Netherlands and NHS in UK), alloantibodies against HNA-1c were described in the cases of NAIN. Interestingly, only the UK laboratory reported alloantibodies against HNA-3a in cases of NAIN. There have been rare reports on NAIN associated with high-frequency antigens HNA-4a [12] and HNA-5a [13]. As HNA-4b frequency is very low in many populations, this may explain the rare reports of NAIN cases mediated by alloantibodies against this antigen. In addition, the low frequency of antibodies against HNA-4 and HNA-5 can also be correlated to the non-immunogenic structure of CD11b and CD11a where HNA-4 and HNA-5 epitopes are located.

The survey results show that AIN is the most requested HNA investigation, conducted in 15 established laboratories and the laboratories in development in Iran (Table 5). Specifically, primary AIN occurs mainly in infants and children. The AIN autoantibody is elusive because the antibody concentration can change. Hence, AIN investigation often requires testing of multiple patient sera at different times to detect the antibody [14]. The antibody specificity has

TABLE 3 Human neutrophil antigen (HNA)-genotyping conducted by participants.

Genotyping	Y/N	HNA-1	HNA-2	HNA-3	HNA-4	HNA-5
1. Brazil, Sao Paulo	Y	HNA-1		HNA-3	HNA-4	HNA-5
2. USA, ARC	Y	HNA-1	HNA-2	HNA-3	HNA-4	HNA-5
3. USA, Versiti	Y	HNA-1		HNA-3	HNA-4	HNA-5
4. Austria, Vienna	Y	HNA-1	HNA-2	HNA-3	HNA-4	HNA-5
5. Denmark, Aalborg	Y	HNA-1	HNA-2	HNA-3	HNA-4	HNA-5
6. France, Créteil	Y	HNA-1		HNA-3	HNA-4	HNA-5
7. France, Nantes	Y	HNA-1		HNA-3	HNA-4	HNA-5
8. Germany, Dessau	Y	HNA-1		HNA-3	HNA-4	HNA-5
9. Germany, Giessen	Y	HNA-1		HNA-3	HNA-4	HNA-5
10. The Netherlands, Sanquin	Y	HNA-1		HNA-3	HNA-4	HNA-5
11. Poland, Warsaw	Y	HNA-1	HNA-2	HNA-3	HNA-4	HNA-5
12. United Kingdom, NHS	Y	HNA-1		HNA-3	HNA-4	HNA-5
13. Slovenia, Ljubljana	Y	HNA-1		HNA-3	HNA-4	HNA-5
14. Thailand, Thammasat University	Y	HNA-1		HNA-3	HNA-4	HNA-5
15. Australia, Lifeblood	Y	HNA-1		HNA-3	HNA-4	HNA-5
16. Hong Kong, Queen Mary	Y	HNA-1	HNA-2	HNA-3	HNA-4	HNA-5
17. Japan, JRC	Y	HNA-1		HNA-3		
<i>Laboratories in development</i>						
1. South Africa, SANBS	N					
2. Iran, Tehran	Y	HNA-1		HNA-3	HNA-4	HNA-5
3. Korea, Seoul	Y	HNA-1		HNA-3	HNA-4	HNA-5
4. Thailand, Red Cross	N					

TABLE 4 Human neutrophil antigen (HNA) antibody specificities detected in neonatal alloimmune neutropenia investigations.

Neonatal alloimmune neutropenia investigations		Samples tested (n)	Samples positive (n)	Antibody specificity (%)							HLA-class II	HLA-class I	CD16 Iso Ab
				HNA-1a	HNA-1b	HNA-1c	HNA-2	HNA-3a	HNA-4a				
1	Brazil, Sao Paulo	2	1	0	100	0	0	0	0	0	0	0	
2	USA, ARC	6	2	0	100	0	0	0	0	0	0	0	
3	USA, Versiti	40	3	50	25	0	25	0	0	0	0	0	
4	Austria, Vienna	5	3+	0	33.3	0	33.3	0	0	0	0	33.3	
5	Denmark, Aalborg	7	1	0	100	0	0	0	0	0	0	0	
6	France, Créteil	37	8	62	25	13	0	0	0	0	0	0	
7	France, Nantes	24	6	0	33	17	17	0	0		0	33	
8	Germany, Dessau	3	0	0	0	0	0	0	0	0	0	0	
9	Germany, Giessen	38	8	25	25	0	25	0	0	0	12.5	12.5	
10	The Netherlands, Sanquin	12	5	20	0	20	20	0	0	0	0	40	
11	Warsaw, Poland		380 ^a	42	13	0	0	0	2	6		25	
12	United Kingdom, NHS	55	7	43	0	14	29	14	0	0	0	0	
13	Slovenia, Ljubljana	6	1	50	30	0	20	0	0	0	0	0	
14	Australia, Lifeblood	4	1	0	100	0	0	0	0	0	0	0	
15	Hong Kong, Queen Mary	36	10	100	0	0	0	0	0	0	0	0	

^aTotal number of diagnosed neutropenia (auto- or alloimmune).

also been reported to change with time [15]. Diagnosis of primary AIN is especially useful as it supports exclusion of other causes of neutropenia [16].

Although in many cases of adult AIN no antigen specificity is described, antibodies against HNA-1a and HNA-1b have been detected as the most common target antigens for autoantibodies involved in cases of AIN in adults [17]. In contrast to primary AIN in

infancy, AIN in adults is not a self-limiting condition and manifests a higher frequency in females (up to 70% of cases) [6].

Transfusion of blood products containing HNA alloantibodies to recipients with the cognate antigen has been known to induce reactions such as TRALI [18]. The role of neutrophils in TRALI is well established [19]. However, some alloantibodies such as alloantibodies against HNA-3a have been implicated in severe and some fatal TRALI

TABLE 5 Patient age distribution and positivity rate in neonatal autoimmune neutropenia investigations.

Autoimmune neutropenia investigations		Sample tested (n)	Positive results		Patient age distribution (%)			
			n	%	Newborn	2–36 months	Primary	Adult
1	Brazil, Sao Paulo	33	11	33.3	0.0	36.4	18.2	45.5
2	USA, ARC	95	32	33.7	0.0	43.2	0.0	56.8
3	USA, Versiti	1981	526	26.6	0.1	47.0	18.2	34.8
4	Austria, Vienna	75	15	20.0	21.4	64.3	0.0	14.3
5	Denmark, Aalborg	90	20	22.2	0.0	54.1	18.9	27.0
6	France, Créteil	1169	105	9.0	0.0	30.1	0.0	69.9
7	France, Nantes	774	87	11.2	1.2	26.2	0.0	72.6
8	Germany, Dessau	63	9	14.3	0.0	36.4	36.4	27.3
9	Germany, Giessen	990	255	25.8	0.0	17.9	13.5	68.6
10	The Netherlands, Sanquin	270	79	29.3	57.0			43.0
11	Warsaw, Poland	380			0.0	91.5	0.0	8.5
12	United Kingdom, NHS	2004	488	24.4	0.6	58.5	30.4	10.5
13	Slovenia, Ljubljana	30	8	26.7	0.0	43.2	18.9	37.8
14	Australia, Lifeblood	141	58	41.1	0.0	2.1	1.4	96.5
15	Hong Kong, Queen Mary	11	0	0.0	0.0	18.2	36.4	45.5
LiD	Iran, Tehran	23	12	52.2	0.0	40.0	40.0	20.0

TABLE 6 Transfusion-related acute lung injury (TRALI) investigation findings.

TRALI investigations		Samples tested (n)	Patient antibody		Donor antibody	
			n	%	n	%
1	USA, ARC	60	0	0	6	10.0
2	USA, Versiti	73			1	1.4
3	Austria, Vienna	3	0	0	2	66.7
4	Denmark, Aalborg	9			1	11.1
5	Germany (Dessau)	276	1	0.3	1	0.3
6	Germany (Giessen)	37	0	0	3	8.1
7	France, Nantes	29	0	0	0	0.0
8	The Netherlands, Sanquin	2	0	0	0	0.0
9	Poland, Warsaw	22			1	4.5
10	United Kingdom, NHS	78	Not tested		4	20.0
11	Slovenia, Ljubljana	3			1	33.3
12	Thailand Red Cross	7	1	14.3	1	14.3
13	Thailand, Thammasat University	2	0	0	2	100.0
14	Australia, Lifeblood	2	0	0	1	50.0
15	Hong Kong	0	0	0	0	0
16	Japan, JRC	72	9	12.5	28	38.9

TABLE 7 The human neutrophil antigens (HNAs)-allele frequencies in different populations.

Population	HNA-1			HNA-2		HNA-3		HNA-4		HNA-5		References
	1a	1b	1c	Positive	Negative	3a	3b	4a	4b	5a	5b	
Asian												
Southern Thai	0.619	0.365	0.012			0.808	0.192	0.973	0.027	0.656	0.344	Intharanut et al., 2019 [42]
Central Thai	0.548	0.452	0.004			0.718	0.282	0.975	0.025	0.771	0.229	Intharanut et al., 2019 [42]
Northern Thai	0.677	0.323	0			0.775	0.225	0.965	0.035	0.748	0.252	Intharanut et al., 2019 [42]
Northeastern Thai	0.696	0.301	0			0.785	0.215	0.972	0.028	0.676	0.324	Intharanut et al., 2019 [42]
Thai				0.995	0.005							Nathalang et al., 2018 [41]
Chinese (Han, Guangzhou)	0.667	0.333	0	1	0	0.738	0.262	0.996	0.004	0.854	0.146	Xia et al., 2011 [33]
Chinese (Hong Kong)	0.678	0.315	0	0.983	0.017	0.71	0.29	0.995	0.005	0.852	0.148	Tam et al., 2018 [35]
Chinese (Han, Zhejiang)	0.613	0.387	0			0.654	0.346	1	0	0.896	0.104	He and Zhang, 2014 [34]
Korean						0.695	0.305	0.986	0.014	0.959	0.041	Han and Han, 2015 [38]; Han and Han, 2006 [39]
Indian	0.433	0.444	0.086	0.9927	0.012	0.812	0.188	0.955	0.045	0.237	0.763	Gogri et al., 2022 [36]
Japanese	0.623	0.377	0	0.987	0.013	0.654	0.346	1	0	0.84	0.16	Matsuhashi et al., 2012 [37]
Burmese	0.605	0.395	0.031			0.747	0.253	0.971	0.029	0.559	0.441	Simtonget al., 2018 [32]
Karen	0.725	0.275	0			0.845	0.155	0.956	0.044	0.693	0.307	Simtong et al., 2018 [32]
Malays (Total)	0.706	0.294	0.037			0.758	0.242	0.977	0.023	0.708	0.292	Manaf et al., 2015 [40]
European												
German	0.36	0.631	0.019			0.801	0.199	0.889	0.111	0.665	0.335	Grabowski et al., 2019 [44]
Danish	0.348	0.623	0.029			0.814	0.186	0.881	0.119	0.724	0.276	Nielsen et al., 2012 [26]
English (Caucasoid)	0.318	0.668	0.014			0.768	0.232	0.882	0.118	0.736	0.264	Cardoso et al., 2013 [43]
Russian (St. Petersburg)	0.384	0.584	0.032			0.804	0.196	0.898	0.102	0.708	0.292	Krobinets et al., 2020 [45]
Turkish	0.42	0.564	0.03			0.737	0.263	0.881	0.119	0.754	0.246	Hauck et al., 2011 [46]
African												
Zambian	0.395	0.345	0.25			0.975	0.025	0.895	0.105	0.5	0.5	Nielsen et al., 2012 [26]
American												
United States (Black population)	0.59	0.77	0.23			0.929	0.071					Kissel et al., 2000 [31]
United States (African American)						0.826	0.174					Bowens et al., 2012 [30]
United States (Hispanic/Latino)						0.946	0.054					Bowens et al., 2012 [30]
United States (Native)						0.81	0.19	0.822	0.178	0.711	0.289	Bowens et al., 2012 [30]
Brazilian	0.315	0.637	0.048									Santos et al., 2016 [27]; Lopes et al., 2014 [28]; Cardone et al., 2006 [29]

(Continues)

TABLE 7 (Continued)

Population	HNA-1			HNA-2		HNA-3		HNA-4		HNA-5		References
	1a	1b	1c	Positive	Negative	3a	3b	4a	4b	5a	5b	
Middle Eastern												
Iranian	0.34	0.63	0.03			0.63	0.37	0.85	0.15	0.72	0.28	Esmaili, 2022 [3]
Syrian	0.375	0.58	0.04			0.742	0.258	0.86	0.14	0.66	0.34	Hauck-Dilmi et al., 2018 [47]

cases. This may be due to the fact that HNA-3a is expressed not only on neutrophils but also on other cells such as endothelial cells, monocytes and platelets [18, 20]. Conscious of the role of HNA antibodies in TRALI, the ISBT Working Party on Granulocyte Immunobiology published recommendations for leukocyte antibody investigations in TRALI [21]. Data from this survey confirm that in the majority of cases the antibodies are in the transfused blood product (Table 6). As two-thirds of TRALI cases are associated with alloantibodies; both HLA Class I and II, and HNA [22], these investigations provide a vital way to identify and manage donors with the culprit antibodies.

During COVID-19 pandemic, transfusion of plasma obtained from COVID-19-immunized healthy donors containing neutralizing antibodies against COVID-19 antigen to COVID-19 patients (known as COVID-19 convalescent plasma [CCP]) was considered as promising therapy. It is worth noting that previous reports indicate the development of TRALI in COVID-19-infected patients after transfusion with CCPs [23, 24]. Therefore, analysis of convalescent plasma for detection of anti-HNAs and HLAs antibodies was conducted. Among participants of this current survey, four laboratories have tested CCPs. However, the frequency of anti-HNAs positive samples among investigated CCPs in these laboratories is not yet reported.

Investigation for the presence of HNA antibodies usually involves screening test, most commonly GIFT and GAT [25, 26]. Use of HNA-typed panel cells at the screening stage may facilitate identification of antibody specificity. The MAIGA provides another way to confirm antibody specificity but is liable to false negatives. Among this survey's participants, 16 established laboratories and 1 laboratory in development conducted GIFT, 11 of 17 established laboratories and one laboratory in development conduct GAT and 12 established laboratories conduct MAIGA with two laboratories optimizing the MAIGA (Table 2). Six established laboratories and three laboratories in development use the LabScreen Multi test kit [21]. The combination of GIFT, GAT and MAIGA assays is considered as the 'gold standard' for HNA testing.

It is interesting to note that although there are only five established laboratories in the Asia Pacific, since 2012, there have been at least 12 publications of HNA frequencies from that region (Table 7). Genotyping has made this feasible. There is a gap in our knowledge on HNA frequencies in African, the Middle Eastern and Western Pacific populations (Table 7). HNAs allele frequency analysis among different populations indicated a different pattern for HNA-1 alleles between Asian and Western populations; in many Asian populations, no HNA-1c allele has been detected. Currently, we do not know if the deviation in allele frequency is due to genotyping techniques applied in the region or if there is a real absence of HNA-1c in some Asian populations.

To evaluate the accuracy and effectiveness of a laboratory's ability to detect HNA antibodies, the ISBT-GIWP conducts the International Granulocyte Immunobiology Working Party (IGIWP) every year. In 2022, each of the 18 participating laboratories received four blinded samples (four DNA and four serum samples) to analyse. Of the 18 participating laboratories, only 13 responded to this survey. It is hoped that this survey will encourage other laboratories to

participate in IGIWP workshop. In addition to IGIWP, there is the INSTAND EQA for Granulocyte Immunobiology. This is an interdisciplinary non-profit, scientific-medical association designated by the German Medical Association to promote quality assurance. For each evaluation, INSTAND distributes blinded samples for HNAs genotyping and detection of anti-HNA-alloantibodies.

This is the first study to provide a comprehensive synopsis of HNA typing and serology conducted in diagnostic laboratories across the whole world. The number of samples received and analysed provides a picture of HNA-related diseases such as adults and infants AIN, immunization frequencies against HNA and is indication of the population-related HNA involved in alloimmune neutropenia. The data provide a summary of investigation techniques employed that may help interested health services develop HNA investigation capability and improve the activities of established laboratories. Importantly, the list of participants (Tables 1A and 1B) provides a useful contact list for physicians and also regional hospitals wanting to contact HNA-diagnostic services.

ACKNOWLEDGEMENTS

B.B. and L.C. designed the research study, prepared the questionnaire and mailing list, collected data and wrote the manuscript, J.L. prepared the synopsis on HNA allele frequencies in different populations, M.A., L.C., B.C., R.D., B.E., C.G., M.K., H.Ki., H.Kr., M.M.K., J.K., E.M., O.N., D.N., K.R.N., G.P., A.P., L.P., U.J.S., M.S., G.F.K., P.K., D.T. and M. U. provided the data. B.F. provided valuable input into the drafting and finalization of the manuscript. Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST STATEMENT

There are no conflicts identified.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Behnaz Bayat  <https://orcid.org/0000-0002-5608-4317>

Elyse Moritz  <https://orcid.org/0000-0002-0636-4556>

Oytip Nathalang  <https://orcid.org/0000-0002-2858-3063>

Yoke-Lin Fung  <https://orcid.org/0000-0002-7777-6179>

REFERENCES

- Flesch BK, Reil A. Molecular genetics of the human neutrophil antigens. *Transfus Med Hemother*. 2018;45:300–9.
- Browne T, Dearman RJ, Poles A. Human neutrophil antigens: nature, clinical significance and detection. *Int J Immunogenet*. 2021;48:145–56.
- Esmaili B, Bayat B, Alirezaee A, Delkhah M, Mehdizadeh MR, Pourpak Z. Human neutrophil antigen genotype and allele frequencies in Iranian blood donors. *J Immunol Res*. 2022;2022:e4387555.
- Porcelijn L, de Haas M. Neonatal alloimmune neutropenia. *Transfus Med Hemother*. 2018;45:311–6.
- Justiz Vaillant AA, Zito PM. Neutropenia. *StatPearls*. Treasure Island, FL: StatPearls Publishing; 2022.
- Capsoni F, Sarzi-Puttini P, Zanella A. Primary and secondary autoimmune neutropenia. *Arthritis Res Ther*. 2005;7:208–14.
- Solomou EE, Salamaliki C, Lagadinou M. How to make the right diagnosis in neutropenia. *Clin Hematol Int*. 2021;3:41–6.
- Martins JO, Moritz E, Abbas SA, Lopes LB, Barros MMO, Chiba AK, et al. Antibodies against human neutrophil antigens in non-transfused women with red blood cell alloimmunisation induced by pregnancy. *Blood Transfus*. 2021;19:479–86.
- Bux J, Kober B, Kiefel V, Mueller-Eckhardt C. Analysis of granulocyte-reactive antibodies using an immunoassay based upon monoclonal-antibody-specific immobilization of granulocyte antigens. *Transfus Med*. 1993;3:157–62.
- Prasath A, Grafius A, Bonanno M, Ambrusco S, Nair J. Alloimmune neutropenia in a neonate: case report and review of literature. *Antibodies*. 2022;11:63.
- Arneth B. Neonatal immune incompatibilities between newborn and mother. *J Clin Med*. 2020;9:1470.
- Fung YL, Pitcher LA, Willett JE, Reed C, Mison L, Bux J, et al. Alloimmune neonatal neutropenia linked to anti-HNA-4a. *Transfus Med*. 2003;13:49–52.
- Porcelijn L, Abbink F, Terraneo L, Hoogen LO, Huiskes E, de Haas M. Neonatal alloimmune neutropenia due to immunoglobulin G antibodies against human neutrophil antigen-5a. *Transfusion*. 2011;51:574–7.
- Bux J, Behrens G, Jaeger G, Welte K. Diagnosis and clinical course of autoimmune neutropenia in infancy: analysis of 240 cases. *Blood*. 1998;91:181–6.
- Bruin M, Dassen A, Pakrt D, Buddelmeyer L, Kuijpers T, de Haas M. Primary autoimmune neutropenia in children: a study of neutrophil antibodies and clinical course. *Vox Sang*. 2005;88:52–9.
- Audrain M, Martin J, Fromont P, Prié N, Thomas C, Muller et J-Y. Autoimmune neutropenia in children: analysis of 116 cases: autoimmune neutropenia in children. *Pediatr Allergy Immunol*. 2011;22:494–6.
- Fujita M, Kawabata H, Oka T, Hishizawa M, Kitano T, Kondo T, et al. A rare case of adult autoimmune neutropenia successfully treated with prednisolone. *Intern Med*. 2017;56:1415–9.
- Reil A, Keller-Stanislawski B, Günay S, Bux J. Specificities of leucocyte alloantibodies in transfusion-related acute lung injury and results of leucocyte antibody screening of blood donors. *Vox Sang*. 2008;95:313–7.
- Fung YL, Silliman CC. The role of neutrophils in the pathogenesis of transfusion-related acute lung injury. *Transfus Med Rev*. 2009;23:266–83.
- Bayat B, Tjahjono Y, Sydykov A, Werth S, Hippenstiel S, Weissmann N, et al. Anti-human neutrophil antigen-3a induced transfusion-related acute lung injury in mice by direct disturbance of lung endothelial cells. *Arterioscler Thromb Vasc Biol*. 2013;33:2538–48.
- ISBT Working Party on Granulocyte Immunobiology, Bierling P, Bux J, Curtis B, Flesch B, Fung L, et al. Recommendations of the ISBT Working Party on Granulocyte Immunobiology for leucocyte antibody screening in the investigation and prevention of antibody-mediated transfusion-related acute lung injury. *Vox Sang*. 2009;96:266–9.
- Muller MCA, Porcelijn L, Vlaar APJ. Prevention of immune-mediated transfusion-related acute lung injury; from bloodbank to patient. *Curr Pharm Des*. 2012;18:3241–8.
- Amrutiya V, Patel R, Baghal M, Patel B, Waykole T, Patel H, et al. Transfusion-related acute lung injury in a COVID-19-positive convalescent plasma recipient: a case report. *J Int Med Res*. 2021;49:1–5.
- Juskewitch JE, Stubbs JR, Gandhi MJ. Elevated rate of HLA antibodies in male COVID-19 convalescent plasma donors: a risk factor for transfusion-related acute lung injury. *Mayo Clin Proc*. 2021;96:500–2.

25. Fung YL, Minchinton RM, Fraser JF. Neutrophil antibody diagnostics and screening: review of the classical versus the emerging. *Vox Sang*. 2011;101:282–90.
26. Nielsen KR, Koelbaek MD, Varming K, Baech J, Steffensen R. Frequencies of HNA-1, HNA-3, HNA-4, and HNA-5 in the Danish and Zambian populations determined using a novel TaqMan real time polymerase chain reaction method. *Tissue Antigens*. 2012;80:249–53.
27. Santos VC, Grecco M, Pereira KMC, Terzian CCN, Andrade LEC, Silva NP. Fc gamma receptor IIIb polymorphism and systemic lupus erythematosus: association with disease susceptibility and identification of a novel FCGR3B*01 variant. *Lupus*. 2016;25:1237–43.
28. Lopes LB, Baleotti W Jr, Suzuki RB, Fabron A Jr, Chiba AK, Vieira-Filho JPB, et al. HNA-3 gene frequencies in Brazilians and a new polymerase chain reaction–restriction fragment length polymorphism method for HNA-3a/3b genotyping. *Transfusion*. 2014;54:1619–21.
29. Cardone JDB, Bordin JO, Chiba AK, Norcia AMMI, Vieira-Filho JPB. Gene frequencies of the HNA-4a and -5a neutrophil antigens in Brazilian persons and a new polymerase chain reaction–restriction fragment length polymorphism method for HNA-5a genotyping. *Transfusion*. 2006;46:1515–20.
30. Bowens KL, Sullivan MJ, Curtis BR. Determination of neutrophil antigen HNA-3a and HNA-3b genotype frequencies in six racial groups by high-throughput 5' exonuclease assay. *Transfusion*. 2012; 52:2368–74.
31. Kissel K, Hofmann C, Gittinger FS, Daniels G, Bux J. HNA-1a, HNA-1b, and HNA-1c (NA1, NA2, SH) frequencies in African and American Blacks and in Chinese. *Tissue Antigens* 2000;56:143–8.
32. Simtong P, Puapairoj C, Leelayuwat C, Santoso S, Romphruk AV. Assessment of HNA alloimmunisation risk in Northeastern Thais, Burmese and Karen. *Transfus Med*. 2018;28:47–55.
33. Xia W, Bayat B, Sachs U, Chen Y, Shao Y, Xu X, et al. The frequencies of human neutrophil alloantigens in the Chinese Han population of Guangzhou. *Transfusion*. 2011;51:1271–7.
34. He J, Zhang W. Genotyping of human neutrophil antigens by polymerase chain reaction sequence-based typing. *Blood Transfus*. 2014; 12:s292–s298.
35. Tam K, Tang I, Ho J, Yeung W, Lee CK, Ip P, et al. A study of human neutrophil antigen genotype frequencies in Hong Kong. *Transfus Med*. 2018;28:310–8.
36. Gogri H, Parihar M, Kulkarni S, Madkaikar M, Sharma J, Gorakshakar A. Phenotyping and genotyping of HNA: prevalence, risk of alloimmunization, and HNA incompatibilities in Indians. *Transfus Med Hemother*. 2022;50:1–9.
37. Matsuhashi M, Tsuno NH, Kawabata M, Mishima Y, Okochi N, Santoso S, et al. The frequencies of human neutrophil alloantigens among the Japanese population. *Tissue Antigens*. 2012;80:336–40.
38. Han TH, Han KS. Gene frequency of human neutrophil alloantigen-3 in the Korean population. *Korean J Blood Transfus*. 2015;26:185–92.
39. Han TH, Han KS. Gene frequencies of human neutrophil antigens 4a and 5a in the Korean population. *Korean J Lab Med*. 2006;26:114–8.
40. Manaf SM, NurWaliyuddin HZA, Panneerchelvam S, Zafarina Z, Norazmi MN, Chambers GK, et al. Human neutrophil antigen profiles in Banjar, Bugis, Champa, Jawa and Kelantan Malays in Peninsular Malaysia. *Blood Transfus*. 2015;13:610–5.
41. Nathalang O, Siripanthong K, Petvises S, Jeumjanya N. Flow-cytometric analysis of HNA-2 expression and phenotypes among Thai blood donors. *Ann Lab Med*. 2018;38:362–6.
42. Intharanut K, Sasikarn W, Mitundee S, Nathalang O. HNA-1, -3, -4, and -5 genotyping using multiplex PCR among southern Thais: developing continual HNA-1 null detection. *J Clin Lab Anal*. 2019;33: e22651.
43. Cardoso SP, Chong W, Lucas G, Green A, Navarrete C. Determination of human neutrophil antigen-1, -3, -4 and -5 allele frequencies in English Caucasoid blood donors using a multiplex fluorescent DNA-based assay. *Vox Sang*. 2013;105:65–72.
44. Grabowski C, Jorks S, Kroll H. Genotyping of human neutrophil antigens 1, 3, 4 and 5 using a novel multiplex polymerase chain reaction. *Transfus Med*. 2019;29:110–5.
45. Krobinets II, Mineeva NV, Bogdanova IO, Chechetkin AV. Human neutrophil antigen allele frequencies and assessment of HNA alloimmunisation risk in donors and hematological patients. *Bull Sib Med*. 2020;19:48–54.
46. Hauck B, Philipp A, Eckstein R, Ott S, Zimmermann R, Dengler T, et al. Human neutrophil alloantigen genotype frequencies among blood donors with Turkish and German descent. *Tissue Antigens*. 2011;78:416–20.
47. Hauck-Dlimi B, Damrah M, Achenbach S, Ott S, Zimmermann R, Zingsem J, et al. Human neutrophil alloantigen genotype frequencies among Syrian population. *Clin Lab*. 2018;64:597–601.

How to cite this article: Bayat B, Lowack J, Audrain M, Croisille L, Curtis B, Dangerfield R, et al. World human neutrophil antigens investigation survey. *Vox Sang*. 2023;118: 763–74.