

**Intermediate-dose versus low-dose low-molecular-weight heparin in pregnant and post-partum women with a history of venous thromboembolism (Highlow study)**

*an open-label, multicentre, randomised, controlled trial*

Bistervels, Ingrid M; Buchmüller, Andrea; Wiegers, Hanke M G; Ní Áinle, Fionnuala; Tardy, Bernard; Donnelly, Jennifer; Verhamme, Peter; Jacobsen, Anne F; Hansen, Anette T; Rodger, Marc A; DeSancho, Maria T; Shmakov, Roman G; van Es, Nick; Prins, Martin H; Chauleur, Céline; Middeldorp, Saskia; Highlow Block writing committee; Highlow Investigators

*Published in:*  
Lancet

*DOI (link to publication from Publisher):*  
[10.1016/S0140-6736\(22\)02128-6](https://doi.org/10.1016/S0140-6736(22)02128-6)

*Creative Commons License*  
CC BY-NC-ND 4.0

*Publication date:*  
2022

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Bistervels, I. M., Buchmüller, A., Wiegers, H. M. G., Ní Áinle, F., Tardy, B., Donnelly, J., Verhamme, P., Jacobsen, A. F., Hansen, A. T., Rodger, M. A., DeSancho, M. T., Shmakov, R. G., van Es, N., Prins, M. H., Chauleur, C., Middeldorp, S., Highlow Block writing committee, & Highlow Investigators (2022). Intermediate-dose versus low-dose low-molecular-weight heparin in pregnant and post-partum women with a history of venous thromboembolism (Highlow study): an open-label, multicentre, randomised, controlled trial. *Lancet*, 400(10365), 1777-1787. [https://doi.org/10.1016/S0140-6736\(22\)02128-6](https://doi.org/10.1016/S0140-6736(22)02128-6)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

**Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from [vbn.aau.dk](http://vbn.aau.dk) on: December 05, 2025

Intermediate-dose versus low-dose low-molecular-weight heparin in pregnant and postpartum women with a history of venous thromboembolism (Highlow Study): an open-label, multicentre, randomised, controlled trial

Authors:

Executive writing committee: Ingrid M. Bistervels\*, M.D., Andrea Buchmüller\*, M.D., Hanke M.G. Wiegers\*, M.D., Prof Fionnuala Ní Áinle, M.D., Prof Bernard Tardy, M.D., Jennifer Donnelly, M.D., Peter Verhamme, M.D., Prof Anne F. Jacobsen, M.D., Anette T. Hansen, M.D., Prof Marc A. Rodger, M.D., Prof Maria T. DeSancho, M.D., Prof Roman G. Shmakov, M.D., Nick van Es, M.D., Prof Martin H. Prins, M.D., Prof Céline Chauleur, M.D., Prof Saskia Middeldorp, M.D., *for the Highlow Investigators*\*\*

*\*Ingrid M. Bistervels, Andrea Buchmüller and Hanke M.G. Wiegers contributed equally to this article.*

\*\*The members of the executive writing committee and the block writing committee assume responsibility for the overall content and integrity of this article. The full names, academic degrees, and affiliations of the members of the executive writing committee and the block writing committee are listed in the **Supplementary Appendix**. A complete list of the Highlow Investigators is provided in the **Supplementary Appendix**.

Address reprint requests to Prof. dr. Saskia Middeldorp at Department of Internal Medicine & Radboud Institute of Health Sciences (RIHS), Radboud university medical center, Nijmegen, the Netherlands, or at [saskia.middeldorp@radboudumc.nl](mailto:saskia.middeldorp@radboudumc.nl).

Corresponding author:

Prof. dr. Saskia Middeldorp, Department of Internal Medicine, Radboud university medical center, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, the Netherlands  
saskia.middeldorp@radboudumc.nl, telephone: +31 24 3614763; mobile number +31610630300

Word count abstract: 313

Word count manuscript: 3701

Number of references: 24

Number of tables/figures/panels: 3/3/1

## Abstract

**Background:** Pregnancy-related venous thromboembolism (VTE) is a leading cause of maternal morbidity and mortality, and thromboprophylaxis is indicated in pregnant and postpartum women with a history of VTE. The optimal dose of low-molecular-weight heparin to prevent recurrent venous thromboembolism in pregnancy and the postpartum period is uncertain.

**Methods:** In this international, open-label, randomised trial, pregnant women with a history of venous thromboembolism were randomised before 14 weeks of gestational age to weight-adjusted intermediate-dose or fixed low-dose low-molecular-weight heparin until 6 weeks postpartum. The primary efficacy outcome was objectively confirmed venous thromboembolism. The primary safety outcome was major bleeding.

**Findings:** A total of 1,110 pregnant women were randomised and included in the intention-to-treat population. Venous thromboembolism occurred in 11 of 555 (2.0%) women assigned to weight-adjusted intermediate-dose low-molecular-weight heparin and in 16 of 555 (2.9%) assigned to fixed low-dose low-molecular-weight heparin (relative risk, 0.69; 95% confidence interval [CI], 0.32-1.47;  $P=0.33$ ). Venous thromboembolism occurred antepartum in five (0.9%) and five (0.9%) women, and postpartum in six (1.1%) and 11 women (2.0%) in the intermediate-dose and low-dose groups, respectively. On-treatment venous thromboembolism in the per-protocol population ( $N=972$ ) occurred in 1.0% and 2.4% (relative risk, 0.43; 95% CI, 0.15-1.20). On-treatment major bleeding in the safety population ( $N=1,045$ ) occurred in 4.4% and 3.8% receiving intermediate-dose or low-dose low-molecular-weight heparin, respectively (relative risk, 1.16; 95% CI, 0.65-2.09).

**Interpretation:** In women with a history of venous thromboembolism, weight-adjusted intermediate-dose low-molecular-weight heparin during the combined antepartum and postpartum periods was not associated with a lower risk of recurrence than fixed low-dose low-molecular-weight heparin. These results indicate that low-dose low-molecular-weight heparin for thromboprophylaxis during pregnancy is the

appropriate dose. Postpartum, intermediate-dose low-molecular-weight heparin may be more effective than low dose low-molecular-weight heparin. (ClinicalTrials.gov number, NCT01828697)

**Funding:** This investigator-initiated study was financially supported by grants from the French Ministry of Health, Health Research Board Ireland, and unrestricted grants from GSK/Aspen and Pfizer.

**Key words (MeSH terms)**

venous thromboembolism; venous thrombosis; pulmonary embolism; pregnancy; postpartum period; postpartum haemorrhage; heparin; heparin, low-molecular-weight

## 1    **Research in context**

### 2    *Evidence before this study*

3    We searched PubMed for studies published between the inception of the database and June 29, 2022, using  
4    the search terms (“venous thrombosis” OR “pulmonary embolism” OR “venous thromboembolism”) AND  
5    (“heparin” OR “low-molecular-weight heparin” OR “thromboprophylaxis”) AND “pregnancy” to find  
6    randomised trials and meta-analyses of randomised trials, published in English, that evaluated the  
7    effectiveness of heparin or low-molecular-weight heparin in pregnant women with a history of venous  
8    thromboembolism. We found two small randomised, controlled trials that evaluated the use of heparin in  
9    pregnant women with a history of venous thromboembolism. One was a randomised trial focusing on the  
10    safety of heparin during pregnancy, that included 40 women with a history of venous thromboembolism.  
11    Women were randomised between unfractionated heparin 10,000 international units (IU) twice daily  
12    antepartum followed by unfractionated heparin 8,000 IU twice daily for 6 weeks postpartum, or to  
13    unfractionated heparin 8,000 IU twice daily for 6 weeks postpartum alone. One woman in the control group  
14    developed deep-vein thrombosis at a gestational age of 28 weeks. The other trial was a placebo-controlled  
15    randomised pilot trial of enoxaparin 40 mg that included women with a history of previous thromboembolic  
16    events, women with a known congenital thrombophilia, and women with other accepted risk factors for  
17    which clinicians would consider the use of antenatal heparin. The primary outcome was the number of  
18    recruited women. The recruitment period was 22 months in which 16 women were recruited in 11 centres;  
19    one woman in the placebo group had pulmonary embolism 29 days after delivery. We also identified one  
20    randomised, controlled trial in 292 women with thrombophilia and various clinical manifestations that  
21    investigated antepartum prophylactic dose dalteparin of 5,000 IU once daily, doubled at 20 weeks  
22    gestational age, versus no antenatal dalteparin; all women received dalteparin 5,000 IU once daily  
23    postpartum for 6 weeks. The trial included 36 women with a history of venous thromboembolism. Of 21  
24    women allocated to antenatal dalteparin, one patient had antenatal venous thromboembolism and two

postpartum venous thromboembolism, whereas none of 9 women in the control group had venous thromboembolism. The abovementioned randomised trials have also been summarised in a systematic review and an evidence-based guideline.

#### *Added value of this study*

The international Highlow study is the first large, randomised, controlled thromboprophylaxis trial in pregnant and postpartum women with a history of venous thromboembolism, comparing two doses of low-molecular-weight heparin to prevent recurrence. There was no statistically significant difference between antepartum and postpartum weight-adjusted intermediate-dose low-molecular-weight heparin and fixed low-dose low-molecular-weight heparin on the risk of venous thromboembolism during the combined antepartum and postpartum periods. Our study showed that despite thromboprophylaxis, the absolute risk of venous thromboembolism (deep-vein thrombosis and/or pulmonary embolism) during pregnancy or up to 6 weeks postpartum was 2·0% in women receiving intermediate-dose low-molecular-weight heparin and 2·9% in those receiving low-dose low-molecular-weight heparin (RR 0·69, 95% CI 0·32 to 1·47). Postpartum, intermediate-dose low-molecular-weight heparin appeared to be more effective than low dose low-molecular-weight heparin, with risks of venous thromboembolism of 1·1% and 2·0% respectively.

#### *Implications of all the available evidence*

In women with a history of venous thromboembolism, low-dose low-molecular-weight heparin for thromboprophylaxis during pregnancy is the appropriate dose. The suggestion of a higher efficacy of intermediate-dose low-molecular-weight heparin than low-dose low-molecular weight heparin during the postpartum period is an important finding that calls for confirmation in a future randomised, controlled trial.

## 1 Introduction

2 Deep-vein thrombosis and pulmonary embolism during pregnancy or the postpartum period are the leading  
3 cause of maternal morbidity and mortality.<sup>1,2</sup> Conversely, thromboprophylaxis can contribute to major  
4 bleeding<sup>3</sup>, which also is a major cause of maternal morbidity and mortality. Despite these high stakes,  
5 evidence-based thromboprophylaxis strategies are lacking. Without thromboprophylaxis, women with a  
6 history of venous thromboembolism have a 2 to 10% risk of developing pregnancy-associated recurrent  
7 venous thromboembolism.<sup>4-8</sup> Hence, for pregnant women with a history of venous thromboembolism who  
8 are not using long-term anticoagulation, guidelines recommend postpartum thromboprophylaxis with  
9 subcutaneous low-molecular-weight heparin in all, and antepartum thromboprophylaxis in those who have  
10 a moderate or high risk of recurrent venous thromboembolism.<sup>9-12</sup>

11 The optimal dose of low-molecular-weight heparin for antepartum and postpartum thromboprophylaxis in  
12 women with a history of venous thromboembolism is uncertain. Due to a lack of randomised studies in  
13 pregnancy, dosing is extrapolated from non-pregnant populations.<sup>13-15</sup> However, physiological changes  
14 during pregnancy including weight gain, increase in glomerular filtration rate, and plasma volume expansion  
15 may influence low-molecular-weight heparin pharmacokinetics and reduce efficacy.<sup>16,17</sup> Indeed, pregnancy-  
16 related recurrent venous thromboembolism despite prophylaxis was high in some observational studies.<sup>18-</sup>  
17 <sup>20</sup> Guidelines from several professional societies indicate that there is no evidence to base the suggested  
18 thromboprophylactic dose on <sup>12</sup>, and provide no clear guidance: the American College of Chest Physicians  
19 (ACCP, 2012) suggests the use of either a prophylactic low or intermediate-dose of low-molecular-weight  
20 heparin antepartum and postpartum;<sup>9</sup> the American Society of Hematology (ASH, 2018) suggests  
21 prophylactic low-dose over intermediate-dose low-molecular-weight heparin antepartum, and either dose  
22 postpartum;<sup>11</sup> and the American Society of Obstetrics and Gynecologists (ACOG, 2018) states that  
23 intermediate-dose low-molecular-weight heparin may be considered at extremes of body weight or as  
24 pregnancy progresses.<sup>10</sup>



We performed a randomised, controlled trial comparing the efficacy and safety of intermediate versus low-dose low-molecular-weight heparin in pregnant women with a history of venous thromboembolism.

## **Methods**

### *Study design and participants*

The Highlow study was an investigator-initiated, multicentre, international, open-label, randomised, controlled trial (ClinicalTrials.gov number, NCT01828697), of which the rationale and design have been reported previously.<sup>21</sup> The study was conducted at 70 hospitals in The Netherlands, France, Ireland, Belgium, Norway, Denmark, Canada, United States, and Russia. The protocol was approved by the institutional review board or ethics committee of all participating centres. Written informed consent was obtained from all patients prior to randomisation.

Pregnant women aged 18 years or older with a history of objectively confirmed venous thromboembolism, either unprovoked or provoked by hormonal or minor risk factors, and gestational age of 14 weeks or less, were eligible. Exclusion criteria were a previous venous thromboembolism related to a major risk factor only (i.e. surgery, major trauma or plaster cast immobilization in the absence of concomitant use of hormones), an indication for therapeutic-dose anticoagulants, or a contraindication to low-molecular-weight heparin. Use of low-molecular-weight heparin according to local protocol prior to randomisation was allowed and recorded. Women were allowed to participate more than once and were randomised per pregnancy.

### *Randomisation and masking*

Women were randomly assigned to weight-adjusted intermediate-dose low-molecular-weight heparin or fixed low-dose low-molecular-weight heparin once-daily using a web-based system, balanced in permuted blocks of six and stratified by centre. Hence, physicians and patients could not foresee the outcome of

1 randomisation (i.e. concealment of allocation). After randomisation, there was no masking to assigned study  
2 group for physicians or patients, as medication was supplied by pharmacies in usual patient care settings or  
3 as study drug in accordance with national regulatory requirements. A central independent adjudication  
4 committee, whose members were unaware of the treatment allocation, adjudicated all suspected episodes  
5 of venous thromboembolism, superficial thrombophlebitis, major bleeding, clinically relevant non-major  
6 bleeding, suspected type 1 allergy to low-molecular-weight heparin, and suspected heparin-induced  
7 thrombocytopenia, using pre-specified criteria.

## 8 9 *Procedures*

10 Participants were instructed to administer the allocated dose of low-molecular-weight heparin from pre-  
11 filled syringes subcutaneously. The intermediate-dose low-molecular-weight heparin regimen was  
12 approximately half of a therapeutic dose, categorised by actual body weight and adjusted if needed during  
13 pregnancy or postpartum, with cut-offs of <50 kilograms (kg), 50 to 70 kg, 70 to 100 kg, and  $\geq 100$  kg. Once  
14 daily doses ranged from 3,800 to 9,500 international units (IU) for nadroparin, 6,000 to 12,000 IU for  
15 enoxaparin, 7,500 to 15,000 IU for dalteparin, or 4,500 to 12,000 IU for tinzaparin (**Panel**). The fixed low-  
16 dose regimen was based on weight at randomisation (<100 kg or  $\geq 100$  kg) as per clinical practice in many  
17 centres and suggested by the Royal College of Obstetricians and Gynaecologists' Green-top Guideline <sup>12</sup>,  
18 and consisted of nadroparin 2,850 or 3,800 IU, enoxaparin 4,000 or 6,000 IU, dalteparin 5,000 or 7,500 IU,  
19 or tinzaparin 3,500 or 4,500 IU, and was not changed throughout pregnancy or postpartum. The preferred  
20 type of low-molecular-weight heparin varied per centre. Women were instructed to stop low-molecular-  
21 weight heparin at first signs of labour. If delivery was planned, the last dose was given at least 24 hours prior  
22 to delivery. Required time intervals between last dose of low-molecular-weight heparin and neuraxial  
23 anaesthesia were according to local guidelines, i.e. 24 hours for the intermediate-dose, and 10 to 12 hours

for the low-dose. Low-molecular-weight heparin was continued until 6 weeks postpartum, also if a pregnancy ended in miscarriage, abortion, or stillbirth.

In-person or telephone contacts were scheduled 2 weeks after randomisation, at 20 and 30 weeks of gestation, and 1 week, 6 weeks and 3 months postpartum. At each contact, suspected outcome events, adverse events, compliance with low-molecular-weight heparin use, and concomitant medications were recorded. In the intermediate-dose group, dose adjustments of low-molecular-weight heparin were made if required based on change in body weight. Women were instructed to contact the study team in case of signs or symptoms of venous thromboembolism or bleeding, upon which clinical assessment and diagnostic imaging were performed.

### *Outcomes*

The primary efficacy outcome was symptomatic objectively confirmed venous thromboembolism up to 6 weeks postpartum, including deep-vein thrombosis, pulmonary embolism, or unusual site venous thrombosis, e.g. splanchnic vein or cerebral sinus thrombosis. Secondary efficacy outcomes were the three components of the primary outcome, objectively confirmed superficial thrombophlebitis, and a composite of venous thromboembolism or superficial thrombophlebitis.

The primary safety outcome was major bleeding, which included antepartum, early postpartum (within 24 hours after delivery), and late postpartum major bleeding (after 24 hours of delivery until 6 weeks postpartum) based on population-specific definitions proposed by a subcommittee of the International Society on Thrombosis and Haemostasis (ISTH).<sup>22</sup> Antepartum and late postpartum major bleeding included placenta praevia requiring delivery, placental abruption, fetal or neonatal death due to bleeding, or acute clinically overt maternal bleeding associated with one or more of the following: occurring in a critical organ, associated with a fall in haemoglobin level of 2 g/dL or more, requiring transfusion of two or more units of whole blood or red cells to maintain a haemoglobin level of more than 7-9 g/dL, or leading to maternal

1 death. Early postpartum major bleeding was defined as bleeding within 24 hours after delivery requiring  
2 transfusion of two or more units of whole blood or red cells, or an estimated blood loss of  $\geq 1,000$  mL  
3 necessitating a second line of uterotonics, a uterine intervention for haemostasis, balloon tamponade,  
4 embolisation, conservative surgery, hysterectomy, or maternal death. Secondary safety outcomes were  
5 clinically relevant non-major bleeding and minor bleeding using population-specific definitions.<sup>22</sup>

6 The list of outcomes, including the primary safety outcome, was revised during the trial. In the first version  
7 of the protocol, we defined major bleeding according to the standard ISTH definitions for the evaluation of  
8 anticoagulants,<sup>23</sup> and definitions of postpartum haemorrhage with a cut-off for  $>500$  mL within 24 hours of  
9 delivery. However, after start of the central adjudication by the multidisciplinary committee, its members  
10 judged that these definitions did not reflect clinical relevance of peripartum haemorrhage, ultimately  
11 leading to a pregnancy-specific classification of major bleeding that was endorsed by the ISTH Scientific and  
12 Standardization Committee. The final revision took place in June 2017 before the data were unblinded.

### 14 *Statistical Analysis*

15 The study hypothesis was that weight-adjusted intermediate-dose low-molecular-weight heparin would be  
16 superior to fixed low-dose low-molecular-weight heparin for the prevention of pregnancy-associated  
17 venous thromboembolism. The sample size was event-driven, with a targeted number of 29 primary  
18 outcome events, assuming a 65% relative risk reduction with intermediate-dose as compared with low-dose  
19 low-molecular-weight heparin, a power of 80%, and a two-sided significance level of 0.05. Based on an  
20 expected incidence of venous thromboembolism of 4 to 5% in the low-dose group,<sup>18</sup> the expected sample  
21 size was determined to be 859 to 1,074.

22 The primary efficacy analysis was performed in the intention-to-treat population and included all data and  
23 adjudicated outcomes from randomisation up to 6 weeks postpartum in all randomised women. A  
24 secondary analysis of the efficacy outcomes was performed with outcomes from randomisation up to 3

months postpartum. The primary efficacy outcome up to 6 weeks postpartum was also evaluated in the per-protocol population and included outcomes occurring during the on-treatment period (defined as the time from randomisation to the last day of allocated low-molecular-weight heparin dose plus 2 days). Women were considered off-treatment in case of deviation from the allocated dose for more than 2 consecutive weeks. The per-protocol population included women without major protocol deviations, a self-reported adherence of 80% or more, and who received at least one dose of allocated study treatment. The analysis of safety outcomes was performed in the safety population and included all data and adjudicated outcomes from randomisation up to 6 weeks postpartum during the on-treatment period. The safety population included all women who received at least one dose of allocated study treatment. Predefined subgroup analyses were performed according to maternal age, location of previous venous thromboembolism, provoking factors of previous venous thromboembolism, body mass index (BMI), thrombophilia, parity, low-molecular-weight heparin use before randomisation, and use of acetylsalicylic acid during pregnancy. Sensitivity analyses were performed including the first pregnancy in this study only and excluding women who experienced miscarriage before 14 weeks. For all outcomes, a two-sided Chi-squared test (or a Fisher's exact test if fewer than 5 observations) was performed to compare the intermediate-dose with the low-dose group. Relative risks (RR) with 95% confidence intervals (CI) based on normal approximation were calculated. A time-to-event analysis using Cox proportional hazards models was performed to obtain hazard ratios (HR) with 95% CI, censoring patients at loss to follow-up, withdrawal of informed consent, or end of study period. The proportionality assumption was checked by visual inspection of the log-minus-log plots and assessment of Schoenfeld residuals. Missing data were not imputed and only observed data were analysed. Baseline data that were missing but were collected on follow-up visits were used to complete the baseline table. For baseline variables such as weight, the available weight closest to the randomisation date was used. For primary and secondary outcomes, data from the visits following the missed visit were used.

All analyses were conducted using R, version 4.0.3 (R Project for Statistical Computing, Vienna, Austria), particularly using the package ‘survival’ (version 3.2-7) and ‘epitools’ (version 0.5-10.1).

#### *Role of the funding source*

The Highlow study was partially supported by various grants; the French Ministry of Health (2014, PHRC national, number 1408211), the Health Research Board Ireland (Definitive Interventions and Feasibility Awards (DIFA) 2017, number DIFA-2017-040), GSK (2012), which was taken over by Aspen (2016), and Pfizer. None of the funders had a role in the design, data collection, data analysis, data interpretation or writing of the report.

## **Results**

Between April 24, 2013 and October 31, 2020, 1,110 women were enrolled and randomised (**Figure 1**). The number of included women per country were as follows: The Netherlands (n=516), France (n=388), Ireland (n=99), Belgium (n=42), Norway (n=28), Denmark (n=15), Canada (n=12), United States (n=7), and Russia (n=3). Baseline characteristics are shown in **Table 1**. The mean age was 32.0 years (standard deviation [SD], 4.8), median BMI 25.0 kg/m<sup>2</sup> (interquartile range [IQR], 22.0 to 30.0), and 903 women (81.4%) had a history of venous thromboembolism related to hormone use, pregnancy, or postpartum period. Women were randomised at a median gestational age of 9 weeks and 3 days (IQR, 7+2 to 11+6). Median follow-up duration was 247 days (IQR, 228 to 266) during which 1,018 women (91.7%) had a live birth at a median gestational age of 38 weeks and 5 days (IQR, 37+2 to 39+5).

The primary efficacy outcome of symptomatic venous thromboembolism during pregnancy or up to 6 weeks postpartum occurred in 11 of 555 women (2.0%) in the weight-adjusted intermediate-dose low-molecular-weight heparin group and in 16 of 555 (2.9%) in the fixed low-dose low-molecular-weight heparin group (RR, 0.69; 95% CI, 0.32 to 1.47; P=0.33). The time to occurrence of the primary efficacy outcome is

1 shown in **Figure 2A**. During pregnancy, five of 555 women (0.9%) had venous thromboembolism in each  
2 treatment group. Postpartum, six of 555 women (1.1%) in the intermediate-dose, and 11 of 555 women  
3 (2.0%) in the low-dose group had venous thromboembolism. Pulmonary embolism, a component of the  
4 primary efficacy outcome, occurred in one patient in the intermediate-dose group and in nine patients in  
5 the low-dose group (RR, 0.11; 95% CI, 0.01 to 0.87). None of the thrombotic events were fatal. The  
6 composite of venous thromboembolism or superficial thrombophlebitis occurred in 13 patients (2.3%) in  
7 the intermediate-dose group and in 29 (5.2%) in the low-dose group (RR, 0.45; 95% CI, 0.24 to 0.85).  
8 Findings were consistent in the secondary analyses up to 3 months postpartum (Table 2). Efficacy outcomes  
9 in the intention-to-treat population are shown in **Table 2**, for subgroups in **Table S2**, and for the sensitivity  
10 analyses in **Table S3**. As shown in Table S2, there were no specific subgroups of women who experienced  
11 venous thromboembolism.

12 In the on-treatment analysis of the per-protocol population, the primary efficacy outcome of symptomatic  
13 venous thromboembolism during pregnancy or up to 6 weeks postpartum occurred in five of 481 women  
14 (1.0%) in the intermediate-dose group and in 12 of 491 (2.4%) in the low-dose group (RR, 0.43; 95% CI, 0.15  
15 to 1.20; **Table S4 and Figure S1**). Other secondary efficacy outcomes in the per-protocol population during  
16 the on-treatment period are shown in **Table S4**.

17 Major bleeding during pregnancy or up to 6 weeks postpartum occurred in 23 of 520 women (4.4%)  
18 receiving intermediate-dose low-molecular-weight heparin and in 20 of 525 (3.8%) receiving low-dose low-  
19 molecular-weight heparin (RR, 1.16; 95% CI, 0.65-2.09;  $P=0.63$ ). Antepartum major bleeding occurred in  
20 two of 520 women (0.4%) in the intermediate-dose, and in two of 525 women (0.4%) in the low-dose group.  
21 Early postpartum major bleeding occurred in 19 of 520 women (3.7%) in the intermediate-dose, and in 18  
22 of 525 women (3.4%) in the low-dose group. Late postpartum major bleeding occurred in two of 520 (0.4%)  
23 in the intermediate-dose group and in none of 525 women (0.0%) in the low-dose group. The time to  
24 occurrence of on-treatment major bleeding is shown in **Figure 2B**. There were no maternal deaths during

the study. All safety outcomes are shown in **Table 3** and **Figures S2** and **S3**. The primary safety outcome for subgroups is shown in **Table S5** and for the sensitivity analyses in **Table S3**.

## Discussion

The Highlow study, which involved 1,110 pregnant women with a history of venous thromboembolism, showed that antepartum and postpartum weight-adjusted intermediate-dose low-molecular-weight heparin did not reduce the risk of venous thromboembolism compared to fixed low-dose low-molecular-weight heparin. Despite thromboprophylaxis, in the intention-to-treat analysis including all randomised women (i.e. also those with protocol deviations), we observed an absolute risk of venous thromboembolism during pregnancy or up to 6 weeks postpartum of 2.0% in women receiving intermediate-dose low-molecular-weight heparin and 2.9% in those receiving low-dose low-molecular-weight heparin (RR 0.69, 95% CI 0.32 to 1.47). In the on-treatment analysis in the per-protocol population, the risk difference between the treatment groups appeared larger, but this difference also was not statistically significant (1.0% versus 2.4%; RR, 0.43; 95%CI, 0.15 to 1.20). There was no difference in on-treatment major bleeding (4.4% versus 3.8%, RR 1.16, 95% CI 0.65 to 2.09).

Some observed differences between the treatment groups are noteworthy. First, the risk of pulmonary embolism, a component of the primary efficacy outcome, was markedly lower with intermediate-dose low-molecular-weight heparin than with fixed low-dose low-molecular weight heparin (RR, 0.11; 95% CI, 0.01 to 0.87). Second, venous thromboembolism or superficial thrombophlebitis up to 6 weeks postpartum, a pre-specified secondary efficacy outcome, occurred in 2.3% in the intermediate-dose group and in 5.2% in the low-dose group (RR, 0.45; 95% CI, 0.24 to 0.85). This outcome is clinically relevant, as superficial thrombophlebitis occurring while using thromboprophylaxis often leads to increasing the dose of low-molecular-weight heparin.<sup>24</sup> Third, there appeared to be a differential effect of the interventions in the antepartum versus the postpartum period. Women who were allocated to receive intermediate-dose low-



1 molecular-weight heparin had a lower incidence of postpartum pulmonary embolism (1 versus 7) and  
2 superficial thrombophlebitis (0 versus 11) than women allocated to low-dose low-molecular-weight  
3 heparin. Interestingly, we did not observe subgroups with a differential treatment effect, such as history of  
4 provoked or unprovoked VTE or based on body weight (Table S2). Although the absolute risk of venous  
5 thromboembolism may be higher in women with increased body weight or age, our study was not designed  
6 to draw conclusions about such differences between subgroups.

7 What are the implications of our findings? The results of the Highlow study provide an evidence base for  
8 guidelines and show that low-dose low-molecular-weight heparin for thromboprophylaxis during pregnancy  
9 is the appropriate dose.<sup>9-12</sup> Higher doses of low-molecular-weight heparin complicate peripartum  
10 management due to a longer required interval for neuraxial anaesthesia and are associated with increased  
11 costs and a potential for more side effects such as bruising and bleeding. The suggestion of greater efficacy  
12 of intermediate-dose low-molecular-weight heparin versus low-dose low-molecular weight heparin during  
13 the postpartum period is to be regarded as hypothesis generating and would ideally be confirmed in a future  
14 randomised, controlled trial. In addition, finding ways to increase adherence to LMWH use during pregnancy  
15 and postpartum and assessment of its effect on VTE risk would be extremely valuable.

16 To date, only two small randomised, controlled trials (N=16 and N=40) evaluated the efficacy of  
17 thromboprophylaxis in pregnant women with a history of venous thromboembolism.<sup>14,15</sup> This is likely the  
18 result of major funding, regulatory, ethical and structural barriers challenging the conduct of randomised  
19 trials in pregnancy. In the Highlow study, a large number of women was prospectively followed with careful  
20 documentation of outcomes and adverse events. Loss to follow-up was very low, as was the rate of  
21 withdrawal of consent. As the trial was conducted in nine countries with use of different types of low-  
22 molecular-weight heparin, its findings are generalisable.

23 Our study has limitations. The Highlow trial did not include a placebo arm, as the standard of care according  
24 to various guidelines is to provide pharmacologic thromboprophylaxis with LMWH to women with history

1 of VTE. For pragmatic reasons, we used an open-label design which may increase the risk of diagnostic  
2 suspicion bias. It was also judged unethical to have clinicians blinded to the assigned low-molecular-weight  
3 heparin dose due to the requirement for different peripartum management strategies required for each  
4 group. However, the main efficacy and safety outcomes were adjudicated by a central committee unaware  
5 of treatment allocation. For the intermediate-dose LMWH, we chose to increase the dose with increasing  
6 body weight. We did not increase the low-dose during the course of pregnancy as is suggested by the ACOG  
7 and may be rational based on pharmacokinetic studies.<sup>10,16,17</sup> There was a considerable number of protocol  
8 deviations (n=146), such as nonadherence to required weight adjustments in the intermediate-dose group,  
9 differences in peripartum low-molecular-weight heparin management due to concerns about postpartum  
10 bleeding or inaccessibility to neuraxial anaesthesia, and premature discontinuation of low-molecular-weight  
11 heparin during the postpartum period. The impact of these deviations may be reflected in the greater  
12 observed efficacy of intermediate-dose low-molecular-weight heparin in the on-treatment analysis in the  
13 per-protocol population. We relied on self-reported adherence rather than on a syringe count during each  
14 visit. Selection bias may have been introduced by allowing use of low-molecular-weight heparin prior to  
15 randomisation, multiple enrolments for the same women, and not restricting inclusion to the first  
16 pregnancy after the previous venous thromboembolism, potentially resulting in a population with a lower  
17 risk of recurrence. We allowed inclusion until 14 weeks of gestational age of women who had started  
18 thromboprophylaxis prior to inclusion in the study, since a history of VTE sometimes only becomes apparent  
19 at the first prenatal visit. In the subgroup analyses based on LMWH use prior to randomisation (Table S2)  
20 the relative effect of the interventions is similar between the groups, with absolute risks that vary between  
21 1.5% (with prior LMWH use) and 2.6% (without prior LMWH use) in the intermediate-dose group, and 2.0  
22 to 3.1% respectively in the low-dose group. Although this suggests risk modification by prior use of LMWH,  
23 our data do not allow to draw this conclusion firmly. Sensitivity analyses with exclusion of 67 women who  
24 participated more than once did not materially alter the results. In absence of data, we assumed a relative

1 risk reduction of 65% in an event-driven sample size calculation, but the observed relative risk reduction  
2 was smaller. Because of multiple reports<sup>18-20</sup> of “breakthrough recurrences” on low-dose LMWH, we  
3 hypothesized that the low-dose would be as ineffective as placebo and we assumed a relative risk reduction  
4 of the intermediate-dose versus the low-dose of 65% in the sample size calculation, but the observed  
5 absolute risk as well as the relative risk reduction was smaller. Finally, we did not meet the targeted number  
6 of centrally adjudicated and confirmed venous thromboembolism.

7 The knowledge gap about optimal low-molecular-weight heparin dosing in pregnant and postpartum  
8 women has been explicitly identified by major guidelines.<sup>2,11</sup> Hence, we are confident that the results of  
9 our study will impact international and national guidelines and that recommendations will be rapidly taken  
10 up by clinicians, knowledge users and policymakers throughout the world. For individual clinicians,  
11 counselling of women facing a pregnancy challenged by a history of venous thromboembolism will, for the  
12 first time, be supported by high-quality data.

13  
14 In conclusion, among women with a history of venous thromboembolism, weight-adjusted intermediate-  
15 dose low-molecular-weight heparin during the combined antepartum and postpartum periods was not  
16 associated with a lower risk of recurrence than fixed low-dose low-molecular-weight heparin. Secondary  
17 data analysis suggests intermediate dose low-molecular-weight heparin may be more effective than low  
18 dose low-molecular-weight heparin in the postpartum period.

## Authorship Contributions

SM designed the trial. SM, AB, FNA, PV, AFJ, ATH, MAR, MTdS, and RGS were national coordinators. SMB, IMB and HMGW were international coordinators. IMB, AB, HMGW, FNA, JD, PV, AFJ, ATH, MAR, MTdS, RGS, CC, and all investigators listed in the supplementary appendix recruited patients. SM, IMB, HMGW, NvE and MHP designed the statistical analysis plan, HMGW and NvE performed the analyses and data were interpreted by all authors. All authors had full access to the data. IMB, HMGW, NvE and SM wrote the first draft of the manuscript, which was critically reviewed and revised by all authors. The executive writing committee (IMB, AB, HMGW, FNA, BT, JD, PV, AFJ, ATH, MAR, MTdS, RGS, NvE, MHP, CC, SM) was responsible for the decision to submit the manuscript. All authors approved the final version of the manuscript.

## Declaration of interests

Executive writing committee; the authors report the following unrelated support: FNA reports grants as principal investigator from the Irish Health Research Board during the conduct of the study, and grants as PI from Daiichi-Sankyo, Bayer and Sanofi (IIS paid to university) outside the submitted work; AB reports grants from French Ministry of Health, during the conduct of the study; JD reports grants as principal investigator from the Irish Health Research Board during the conduct of the study; NvE reports other fees from Bristol-Meyers Squibb, LEO Pharma, and Bayer, outside the submitted work; AFJ reports personal fees from Sanofi-Aventis, outside the submitted work; SM reports grants from GSK, Aspen, and Pfizer during the conduct of the study, grants and personal fees from Daiichi-Sankyo, Bayer, Pfizer, and Boehringer-Ingelheim, personal fees from Portola/Alexion, Abbvie, Pfizer/ Bristol-Meyers Squibb, Norgine, Viatrix, and Sanofi, outside the submitted work; MTdS reports personal fees from Sanofi Genzyme and Bioproducts Laboratory, outside the submitted work; PV reports grants and personal fees from LEO Pharma, Boehringer-

1 Ingelheim, Daiichi -Sankyo, Bayer, Pfizer / Bristol-Meyers Squibb, and personal fees from Anthos,  
2 Portola/Alexion, outside the submitted work.

3 Block writing committee; the authors report the following unrelated support: LB reports grants from French  
4 Ministry of Health, during the conduct of the study; BB reports grants from the Irish Health Research Board  
5 during the conduct of the study; FC reports grants and personal fees from Bayer, Bristol-Meyers Squibb,  
6 Pfizer, and other fees from Sanofi and LEO Pharma, outside the submitted work; JF reports personal fees  
7 from CSL Behring, outside the submitted work.

8 Other authors declare no competing financial interests.

## 10 **Data sharing**

11 Requests for data sharing can be sent to [saskia.middeldorp@radboudumc.nl](mailto:saskia.middeldorp@radboudumc.nl). Only deidentified data will  
12 be made available to academic researchers upon reasonable request. Data will be available once all planned  
13 analyses have been completed and published or presented. Non-Highlow investigators will be asked to sign  
14 a Data Sharing Agreement prior to data sharing. The study protocol, statistical analysis plan, Central  
15 Adjudication Committee Charter, and informed consent form have been available without restrictions to  
16 others from the start of the study on.

## 17 **Acknowledgments**

18 The Highlow study was partially supported by a grant from the French Ministry of Health (2014, PHRC  
19 national, number 1408211), the Health Research Board Ireland (Definitive Interventions and Feasibility  
20 Awards (DIFA) 2017, number DIFA-2017-040), an unrestricted grant from GSK (2012), which was taken over  
21 by Aspen (2016), and Pfizer. Neither the grant agencies nor GSK, Aspen, or Pfizer had any role in the design,

1 data collection, analysis, or interpretation of the study results. No funding for this specific analysis was  
2 obtained and authors declare no competing financial interests.

3 We would like to acknowledge all women who participated in the study as well as all Highlow investigators  
4 (listed in Supplementary Appendix) and all clinicians who referred patients to us. We would like to thank  
5 Drs. Jan Beyer-Westendorf, Andra James, Michael Greaves, Ida Martinelli, Ingrid Pabinger and Stavros  
6 Konstantinides for their support during the initial design of the trial. The Highlow study was endorsed and  
7 infrastructurally supported by the Dutch Thrombosis Network (DTN), Dutch Consortium for Healthcare  
8 Evaluation and Research in Obstetrics and Gynecology (NVOG Consortium 2.0), by two French  
9 gynaecological networks (GO-CIC and CROG), by the French network on thrombosis (INNOVTE), by the Irish  
10 network for VTE research (INViTE), by the Irish HRB Mother and Baby Clinical Trials Network  
11 (<http://www.hrb-mbctni.ie/>) and by the international network of venous thrombosis networks INVENT  
12 ([www.INVENT-VTE.com](http://www.INVENT-VTE.com)). We would like to thank these networks for the collaboration and their support.

13

## References

1. Greer IA. Pregnancy Complicated by Venous Thrombosis. *N Engl J Med* 2015; **373**(6): 540-7.
2. Knight M, Bunch K, Tuffnell D, et al. Saving Lives, Improving Mothers' Care - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2017-19.: Oxford: National Perinatal Epidemiology Unit, University of Oxford 2021; 2021.
3. Hull RD, Schellong SM, Tapson VF, et al. Extended-duration venous thromboembolism prophylaxis in acutely ill medical patients with recently reduced mobility: a randomized trial. *Ann Intern Med* 2010; **153**(1): 8-18.
4. James AH, Jamison MG, Brancazio LR, Myers ER. Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *Am J Obstet Gynecol* 2006; **194**(5): 1311-5.
5. Brill-Edwards P, Ginsberg JS, Gent M, et al. Safety of withholding heparin in pregnant women with a history of venous thromboembolism. Recurrence of Clot in This Pregnancy Study Group. *N Engl J Med* 2000; **343**(20): 1439-44.
6. Pabinger I, Grafenhofer H, Kaider A, et al. Risk of pregnancy-associated recurrent venous thromboembolism in women with a history of venous thrombosis. *J Thromb Haemost* 2005; **3**(5): 949-54.
7. De Stefano V, Martinelli I, Rossi E, et al. The risk of recurrent venous thromboembolism in pregnancy and puerperium without antithrombotic prophylaxis. *Br J Haematol* 2006; **135**(3): 386-91.
8. Jacobsen AF, Sandset PM. Venous thromboembolism associated with pregnancy and hormonal therapy. *Best Pract Res Clin Haematol* 2012; **25**(3): 319-32.
9. Bates SM, Greer IA, Middeldorp S, Veenstra DL, Prabulos AM, Vandvik PO. VTE, thrombophilia, antithrombotic therapy, and pregnancy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012; **141**(2 Suppl): e691S-e736S.
10. American College of Obstetricians Gynecologists' Committee on Practice B-O. ACOG Practice Bulletin No. 196: Thromboembolism in Pregnancy. *Obstet Gynecol* 2018; **132**(1): e1-e17.
11. Bates SM, Rajasekhar A, Middeldorp S, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: venous thromboembolism in the context of pregnancy. *Blood Adv* 2018; **2**(22): 3317-59.
12. Royal College of Obstetricians and Gynaecologists. Reducing the Risk of Venous Thromboembolism during Pregnancy and the Puerperium. Green-top Guideline No37a. London: RCOG; 2015.
13. Middleton P, Shepherd E, Gomersall JC. Venous thromboembolism prophylaxis for women at risk during pregnancy and the early postnatal period. *Cochrane Database Syst Rev* 2021; **3**: CD001689.
14. Gates S, Brocklehurst P, Ayers S, Bowler U. Thromboprophylaxis and pregnancy: two randomized controlled pilot trials that used low-molecular-weight heparin. *Am J Obstet Gynecol* 2004; **191**(4): 1296-303.
15. Howell R, Fidler J, Letsky E, de Swiet M. The risks of antenatal subcutaneous heparin prophylaxis: a controlled trial. *British journal of obstetrics and gynaecology* 1983; **90**(12): 1124-8.
16. Sanghavi M, Rutherford JD. Cardiovascular physiology of pregnancy. *Circulation* 2014; **130**(12): 1003-8.
17. Patel JP, Green B, Patel RK, Marsh MS, Davies JG, Arya R. Population pharmacokinetics of enoxaparin during the antenatal period. *Circulation* 2013; **128**(13): 1462-9.
18. Roeters van Lennep JE, Meijer E, Klumper FJ, Middeldorp JM, Bloemenkamp KW, Middeldorp S. Prophylaxis with low-dose low-molecular-weight heparin during pregnancy and postpartum: is it effective? *J Thromb Haemost* 2011; **9**(3): 473-80.
19. Lepercq J, Conard J, Borel-Derlon A, et al. Venous thromboembolism during pregnancy: a retrospective study of enoxaparin safety in 624 pregnancies. *BJOG* 2001; **108**(11): 1134-40.
20. de Moreuil C, Tromeur C, Daoudal A, et al. Risk factors for recurrence during a pregnancy following a first venous thromboembolism: A French observational study. *J Thromb Haemost* 2022; **20**(4): 909-18.

- 1 21. Bleker SM, Buchmuller A, Chauleur C, et al. Low-molecular-weight heparin to prevent recurrent  
2 venous thromboembolism in pregnancy: Rationale and design of the Highlow study, a randomised trial of  
3 two doses. *Thromb Res* 2016; **144**: 62-8.
- 4 22. Tardy B, Chalayer E, Kamphuisen PW, et al. Definition of bleeding events in studies evaluating  
5 prophylactic antithrombotic therapy in pregnant women: A systematic review and a proposal from the ISTH  
6 SSC. *J Thromb Haemost* 2019; **17**(11): 1979-88.
- 7 23. Schulman S, Kearon C, Subcommittee on Control of Anticoagulation of the S, Standardization  
8 Committee of the International Society on T, Haemostasis. Definition of major bleeding in clinical  
9 investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost* 2005; **3**(4):  
10 692-4.
- 11 24. Di Nisio M, Wichers IM, Middeldorp S. Treatment for superficial thrombophlebitis of the leg.  
12 *Cochrane Database Syst Rev* 2018; **2**: CD004982.

13


14



# 1 **Figure 1. CONSORT Flow Diagram**

2 *Abbreviations: LMWH, low-molecular-weight heparin; VTE, venous thromboembolism*

3 *The number of women excluded from the per protocol population is less than the sum of the reasons, since*  
 4 *some patients met more than one major protocol deviation.*



1 Panel: Dosing schemes for all low-molecular-weight heparin types in the Highlow study

2

Weight-adjusted intermediate-dose						Fixed low-dose					
Weight		nadroparin	enoxaparin	dalteparin	tinzaparin	Weight		nadroparin	enoxaparin	dalteparin	tinzaparin
In kg	In lbs					In kg	In lbs				
< 50	< 110	3,800 IU	6,000 IU	7,500 IU	4,500 IU	< 100	< 220	2,850 IU	4,000 IU	5,000 IU	3,500 IU
50 to < 70	110 to < 154	5,700 IU	8,000 IU	10,000 IU	7,000 IU						
70 to < 100	154 to < 220	7,600 IU	10,000 IU	12,500 IU	10,000 IU						
≥100	≥ 220	9,500 IU	12,000 IU	15,000 IU	12,000 IU	≥100	≥ 220	3,800 IU	6,000 IU	7,500 IU	4,500 IU

3 All doses are administered once daily

4 Abbreviations: kg, kilograms; lbs, pounds; IU, International Units; mg, milligram

1 **Table 1. Patient characteristics of the intention-to-treat population**

2

	Intermediate-dose low-molecular-weight heparin (N=555)	Low-dose low-molecular-weight heparin (N=555)
Mean age - years (SD)	32·0 (4·8)	32·0 (4·8)
Median body mass index - kg/m <sup>2</sup> (IQR)	25·0 (22·0-30·0)	25·0 (22·0-29·0)
Weight at randomisation - no. (%)		
<50 kg	9 (1·6)	16 (2·9)
50 to <70 kg	241 (43·4)	237 (42·7)
70 to <100 kg	250 (45·0)	242 (43·6)
≥ 100 kg	55 (9·9)	58 (10·5)
Primigravidity - no. (%)	141 (25·4)	153 (27·6)
Nulliparity - no. (%)	203 (36·6)	214 (38·6)
Median gestational age at randomisation –weeks and days (IQR)	9 and 4 (7 and 3 to 11 and 6)	9 and 3 (7 and 1 to 12 and 0)
Use of low-molecular-weight heparin prior to randomisation - no. (%)	267 (48·1)	248 (44·7)
Median time since previous venous thromboembolism † - years (IQR)	5·5 (2·6-8·9)	5·1 (2·2-9·0)
History of ≥2 episodes of venous thromboembolism - no. (%)	41 (7·4)	46 (8·3)
Location of previous venous thromboembolism ‡ - no. (%)		
Pulmonary embolism with or without DVT ¶	250 (45·0)	222 (40·0)
Upper or lower-extremity DVT only ¶	253 (45·6)	283 (51·0)
Unusual site venous thrombosis §	48 (8·6)	45 (8·1)
No confirmed venous thromboembolism ¥¥	4 (0·7)	4 (0·7)
Provoking factors of previous venous thromboembolism ** - no. (%)		

Hormone therapy for contraception or assisted reproduction	320 (57.7)	326 (58.7)
During pregnancy	100 (18.0)	80 (14.4)
Unprovoked	73 (13.2)	88 (15.9)
Postpartum period	71 (12.8)	69 (12.4)
Air travel	30 (5.4)	38 (6.8)
Minor trauma	16 (2.9)	14 (2.5)
Major transient risk factor only	5 (0.9)	3 (0.5)
Known thrombophilia - no./no. previously tested. (%)	142/310 (25.6)	149/315 (26.8)
History of caesarean section - no. (%)	100 (18.0)	85 (15.3)
History of postpartum haemorrhage $\geq$ - no. (%)	33 (5.9)	30 (5.4)
Known allergic skin reactions to low-molecular-weight heparin - no. (%)	37 (6.7)	44 (7.9)
Type of low-molecular-weight heparin after randomisation - no. (%)		
Enoxaparin	198 (35.7)	215 (38.7)
Nadroparin	205 (36.9)	203 (36.6)
Dalteparin	69 (12.4)	76 (13.7)
Tinzaparin	82 (15.0)	58 (10.5)
Acetylsalicylic acid use during pregnancy - no. (%)	38 (6.8)	33 (5.9)
Country of inclusion - no. (%)		
The Netherlands	259 (46.7)	257 (46.3)
France	194 (35.0)	194 (35.0)
Ireland	49 (8.8)	50 (9.0)
Belgium	21 (3.8)	21 (3.8)
Norway	14 (2.5)	14 (2.5)
Denmark	8 (1.4)	7 (1.3)
Canada	6 (1.1)	6 (1.1)
United States of America	3 (0.5)	4 (0.7)
Russia	1 (0.2)	2 (0.4)

- 1 Abbreviations: DVT, deep-vein thrombosis; IQR, interquartile range; SD, standard deviation; kg/m<sup>2</sup>, kilograms per square meter
- 2

1  
2    ¥ Defined as  $\geq 500\text{mL}$  according to the criteria of the World Health Organization  
3    † In case of multiple episodes of venous thromboembolism: the most recent event  
4    ‡ Some patients had venous thromboembolism at multiple sites at the same time  
5    ¶ Including: extensive thrombophlebitis that was treated as deep-vein thrombosis, muscle vein thrombosis  
6    close to popliteal vein treated as deep-vein thrombosis, isolated calf vein thrombosis, isolated pelvic vein  
7    thrombosis,  
8    § Including cerebral thrombosis, jugular vein thrombosis, abdominal vein thrombosis, and ovarian vein  
9    thrombosis.  
10    \*\* Some patients had more than one risk factor  
11    ¥¥ Includes patients without previous venous thromboembolism (one in intermediate-dose group and one in  
12    low-dose group), patient with arterial thrombosis (one in low-dose group), superficial thrombophlebitis not  
13    treated as deep-vein thrombosis (one in intermediate-dose group and one in low-dose group), and patients  
14    with retinal vein thrombosis (two in intermediate-dose group and one in low-dose group)  
15  
16    Missing for 'Use of low-molecular-weight heparin prior to randomisation': 16 in intermediate-dose group  
17    and 19 in low-dose group.  
18

1 Table 2. Efficacy outcomes

	Intermediate-dose low-molecular- weight heparin	Low-dose low- molecular-weight heparin	Relative risk (95% CI)	Hazard ratio (95% CI)
<b>Efficacy outcomes during pregnancy or until 6 weeks postpartum</b>	no. of patients/total no. (%)			
<i>Intention-to-treat population</i>	555	555		
Primary efficacy outcome: venous thromboembolism	11 (2·0)	16 (2·9)	0·69 (0·32-1·47)	0·68 (0·32-1·47)
Antepartum	5 (0·9)	5 (0·9)		
Postpartum	6 (1·1)	11 (2·0)		
Pulmonary embolism	1(0·2)	9 (1·6)	0·11 (0·01-0·87)	**
Antepartum	0 (0·0)	2 (0·4)		
Postpartum	1 (0·2)	7 (1·3)		
Deep-vein thrombosis	8 (1·4)	6 (1·1)	1·33 (0·47-3·82)	1·32 (0·46-3·81)
Antepartum	4 (0·7)	3 (0·5)		
Postpartum	4 (0·7)	3 (0·5)		
Unusual site venous thrombosis*	2 (0·4)	1 (0·2)	2·00 (0·18- 22·00)	1·99 (0·18-21·96)
Antepartum	1 (0·2)	0 (0·0)		
Postpartum	1 (0·2)	1 (0·2)		
Superficial thrombophlebitis ‡	3 (0·5)	13 (2·3)	0·23 (0·07-0·81)	0·22 (0·06-0·79)
Antepartum	3 (0·5)	2 (0·4)		
Postpartum	0 (0·0)	11 (2·0)		
Venous thromboembolism or superficial thrombophlebitis	13 (2·3)	29 (5·2)	0·45 (0·24-0·85)	0·44 (0·23-0·85)

Antepartum	8 (1·4)	7 (1·3)		
Postpartum	5 (0·9)	22 (4·0)		
<b>Efficacy outcomes during pregnancy or until 3 months postpartum</b>	no. of patients/total no. (%)			
<i>Intention-to-treat population</i>	555	555		
Venous thromboembolism	13 (2·3)	18 (3·2)	0·72 (0·36-1·46)	0·71 (0·35-1·45)
Pulmonary embolism	3 (0·5)	9 (1·6)	0·33 (0·09-1·22)	**
Deep-vein thrombosis	8 (1·4)	7 (1·3)	1·14 (0·42-3·13)	1·14 (0·41-3·13)
Unusual site venous thrombosis*	2 (0·4)	2 (0·4)	1·00 (0·14 -7·07)	0·99 (0·14 -7·05)
Superficial thrombophlebitis	4 (0·7)	13 (2·3)	0·31 (0·10-0·94)	0·30 (0·10-0·93)
Venous thromboembolism or superficial thrombophlebitis	16 (2·9)	31 (5·6)	0·52 (0·29-0·93)	0·51 (0·28-0·92)

Abbreviations: 95% CI, 95% confidence interval.

\* Including: up to 6 weeks: one cerebral venous thrombosis antepartum and one cerebral venous thrombosis postpartum in the intermediate-dose group; one abdominal venous thrombosis in the low-dose group; up to 3 months: one additional cerebral venous thrombosis postpartum in the low-dose group.

\*\* Hazard ratio was not estimated due to violation of the proportionality assumption.

¥ Superficial thrombophlebitis was centrally adjudicated. After diagnosis, two patients in the intermediate-dose group and seven in the low-dose group were treated with therapeutic anticoagulant therapy; one patient in the intermediate-dose group continued with intermediate dose low-molecular-weight heparin, three patients in the low-dose group were treated with intermediate-dose low-molecular-weight heparin; three patients in the low-dose group continued with low-dose low-molecular-weight heparin.

1 Table 3. Safety outcomes

	Intermediate-dose low-molecular- weight heparin	Low-dose low- molecular-weight heparin	Relative risk (95% CI)	Hazard ratio (95% CI)
<b>Safety outcomes during pregnancy or until 6 weeks postpartum</b>	no. of patients/total no. (%)			
<i>Safety population</i>	520	525		
Primary safety outcome: major bleeding	23 (4.4)	20 (3.8)	1.16 (0.65- 2.09)	1.25 (0.69-2.28)
Antepartum	2 (0.4)	2 (0.4)		
Early postpartum	19 (3.7)	18 (3.4)		
Late postpartum	2 (0.4)	0 (0.0)		
Secondary safety outcomes				
Major or clinically relevant non-major bleeding	50 (9.6)	45 (8.6)	1.12 (0.76-1.65)	1.21 (0.81-1.81)
Antepartum	23 (4.4)	10 (1.9)		
Early postpartum	25 (4.8)	35 (7.4)		
Late postpartum	2 (0.4)	0 (0.0)		
Clinically relevant non-major bleeding	27 (5.2)	25 (4.8)	1.09 (0.64-1.85)	**
Antepartum	21 (4.0)	8 (1.5)		
Early postpartum	6 (1.2)	17 (3.2)		
Late postpartum	0 (0.0)	0 (0.0)		
Minor bleeding	76 (14.6)	66 (12.6)	1.16 (0.86-1.58)	1.27 (0.91-1.77)
Antepartum	17 (3.3)	18 (3.4)		
Early postpartum	55 (10.6)	46 (8.7)		



Late postpartum	4 (0·8)	2 (0·4)		
Any bleeding	123 (23·7)	110 (21·0)	1·13 (0·90-1·42)	1·23 (0·95-1·59)
Antepartum	39 (7·5)	28 (5·3)		
Early postpartum	78 (15·0)	80 (15·2)		
Late postpartum	6 (1·2)	2 (0·04)		
<b>Other AE</b>				
Heparin-induced thrombocytopenia	0 (0·0)	0 (0·0)		
Type 1 allergy	8 (1·5)	2 (0·4)		
Congenital abnormality or birth defect	9 (1·7)	5 (1·0)		
Bruising	248 (47·7)	184 (35·0)		
Type IV allergic skin reaction	180 (34·6)	115 (21·9)		

1 Abbreviations: 95% CI, 95% confidence interval.

2 \*\* Hazard ratio was not estimated due to violation of the proportionality assumption.

Figure 2A. Kaplan-Meier cumulative event rates for venous thromboembolism during pregnancy or up to 6 weeks postpartum in the intention-to-treat population





Figure 2B. Kaplan-Meier cumulative event rates for major bleeding during the on-treatment period in the safety population



1 **Appendix (main manuscript)**

2 The members of the executive writing committee are as follows:

3 Ingrid M. Bistervels, M.D., Andrea Buchmüller, M.D., Hanke M.G. Wiegers, M.D., Prof Fionnuala Ní Áinle,  
4 M.D., Prof Bernard Tardy, M.D., Jennifer Donnelly, M.D., Peter Verhamme, M.D., Prof Anne F. Jacobsen,  
5 M.D., Anette T. Hansen, M.D., Prof Marc A. Rodger, M.D., Prof Maria T. DeSancho, M.D., Prof Roman G.  
6 Shmakov, M.D., Nick van Es, M.D., Prof Martin H. Prins, M.D., Prof Céline Chauleur, M.D., Prof Saskia  
7 Middeldorp, M.D.

8

9 The members of the block writing committee are as follows:

10 Eline S. van den Akker, M.D., Prof Mireille N. Bekker, M.D., Thomas van Bemmelen, M.D., Prof Laurent  
11 Bertoletti, M.D., Julie Blanc, M.D., Suzanne M. Bleker, M.D., Aude Bourtembourg-Matras, M.D., Prof  
12 Florence Bretelle, M.D., Bridgette Byrne, M.D., Prof Francis Couturaud, M.D., Pierre Delorme, M.D., Elise S.  
13 Eerenberg, M.D., Maureen T.M. Franssen, M.D., Jens Fuglsang, M.D., Wessel Ganzevoort, M.D., Prof  
14 François Goffinet, M.D., Jiska M. de Haan-Jebbink, M.D., Wieteke Heidema, M.D., Monique A. Hertzberg,  
15 MSc., Marcel M.C. Hovens, M.D., Prof Menno V. Huisman, M.D., Leonie de Jong-Speksnijder, M.D., Prof  
16 Pieter-Willem Kamphuisen, M.D., Denis J. O’Keeffe, M.D., Karine Lacut (*deceased*), M.D., Josje Langenveld,  
17 M.D., M. Simone Lunshof, M.D., Caroline P. Martens, M.D., Adel Merah, M.D., Emmanuelle Le Moigne, M.D.,  
18 Dimitri N.M. Papatsonis, M.D., Prof Gilles Pernod, M.D., Prof Franck Perrotin, M.D., Edith Peynaud-Debayle,  
19 M.D., Prof Fabrice Pierre, M.D., Prof Geneviève Plu Bureau, M.D., Tiphaine Raia-Barjat, M.D., Robbert J.P.  
20 Rijnders, M.D., Roger Rosario, M.D., Marc Ruivard, M.D., Prof Jeannot Schmidt, M.D., Marieke Sueters, M.D.,  
21 Thomas Vanassche, M.D., Marie-Noëlle Varlet, M.D., Alexandre J. Vivanti, M.D., Matthieu Y. van der Vlist,  
22 M.D., Lucet F. van der Voet, M.D., Karlijn C. Vollebregt, M.D., Prof Johanna I.P. de Vries, M.D., Sabina de

1 Weerd, M.D., Peter E. Westerweel, M.D., Lia D.E. Wijnberger, M.D., Marije ten Wolde, M.D., Paula F. Ypma,  
2 M.D., Catherine Zuily-Lamy, M.D., Joost J. Zwart, M.D.

3

4 The authors' affiliations are as follows:

5 Amsterdam University Medical Center, University of Amsterdam, Amsterdam (I.M.B., H.M.G.W., S.M.B.,  
6 N.v.E., W.G., S.M.), Albert Schweitzer Hospital, Dordrecht (P.E.W., S.d.W., M.Y.v.d.V.), Amphia Hospital,  
7 Breda (D.P., L.d.J.-S., M.S.L.), Deventer Hospital, Deventer (J.J.Z., L.F.v.d.V.), Leiden University Medical  
8 Center, Leiden (M.S., M.V.H.), Rijnstate Hospital, Arnhem (M.M.C.H., L.W.), OLVG Hospital, Amsterdam  
9 (E.S.v.d.A., J.M.d.H.-J.), Gelre Ziekenhuizen, Apeldoorn (T.v.B., E.S.E.), Jeroen Bosch Hospital, 's  
10 Hertogenbosch (R.J.P.R.), Haga Hospital, The Hagues (P.F.Y.), University Medical Center Utrecht, Utrecht  
11 (M.N.B.), Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam (J.I.P.d.V.), Spaarne Gasthuis,  
12 Haarlem (K.C.V.), Radboud university medical center, Nijmegen (W.H., S.M.), Flevoziekenhuis, Almere  
13 (M.t.W.), Zuyderland Medical Center, Heerlen (J.L.), and University Medical Center Groningen, Groningen  
14 (M.T.M.F.) — all in the Netherlands;

15 CIC 1408 Centre Hospitalier Universitaire de Saint-Etienne, Saint-Etienne (A.B., C.C., T.R.-B.), Centre  
16 Hospitalier Universitaire de Brest, Brest (E.L.M., F.C., K.L.), Assistance Publique Hôpitaux de Paris Port Royal,  
17 Paris Cité (P.D., F.G., G.P.B.), Hôpital Nord, Marseille (F.B., J.B.), Hôpital Saint Joseph, Marseille (R.R.), Centre  
18 Hospitalier Universitaire de Clermont-Ferrand, Clermont-Ferrand (J.S., M.R.), Assistance Publique Hôpitaux  
19 de Paris Antoine Béclère, Clamart (A.J.V.), Assistance Publique Hôpitaux de Paris Louis-Mourier, Colombes  
20 (E.P.-D.), Centre Hospitalier Universitaire de Grenoble, La Tronche (G.P.), Centre Hospitalier Universitaire  
21 de Besançon, Besançon (A.B.M.), Centre Hospitalier Universitaire de Nancy, Nancy (C.Z.-L.), Centre

1     Hospitalier Universitaire de Poitiers, Poitiers (F. Pierre), Centre Hospitalier Universitaire de Tours, Tours (F.  
2     Perrotin), — all in France;

3     Rotunda Hospital, Dublin (F.N.A., J.D.), School of Medicine, University College Dublin, Dublin (F.N.A.) The  
4     Coombe Women & Infants University Hospital, Dublin (B.B.), University Hospital Limerick, Limerick (D.J.O’K.)  
5     — all in Ireland; University of Leuven, Leuven, Belgium (P.V., T.V., C.P.M.); Oslo University Hospital, Oslo  
6     (A.F.J., M.A.H.), and Faculty of Medicine, University of Oslo, Oslo, - both in Norway; Aalborg University  
7     Hospital, Aalborg (A.T.H.), and Aarhus University Hospital, Aarhus, (A.T.H. and J.F.) — both in Denmark; The  
8     Ottawa Hospital, Ottawa, Canada (M.A.R.); New York Presbyterian Hospital, Weill Cornell Medicine, New  
9     York, United States of America (M.T.d.S.); and Institute of Obstetrics, National Medical Research Center for  
10    Obstetrics, Gynecology and Perinatology, Ministry of Healthcare of the Russian Federation, Moscow, Russia  
11    (R.G.S.).

12    P.W.K., L.B., B.T., M.-N.V., and A.M. were members of the Independent Central Adjudication Committee.  
13    M.H.P. was the study statistician.

14

15    A complete list of the Highlow Investigators is provided in the **Supplementary Appendix**.

16

17