



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Higher physical activity is associated with lower activity limitation

*Cross-sectional analyses among the Spanish working population*

López-Bueno, R.; López-Sánchez, G. F.; Smith, L.; Sundstrup, E.; Andersen, L. L.; Casajús, J. A.

*Published in:*  
Science and Sports

*DOI (link to publication from Publisher):*

[10.1016/j.scispo.2021.12.010](https://doi.org/10.1016/j.scispo.2021.12.010)

*Creative Commons License*  
CC BY-NC-ND 4.0

*Publication date:*  
2023

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

### *Citation for published version (APA):*

López-Bueno, R., López-Sánchez, G. F., Smith, L., Sundstrup, E., Andersen, L. L., & Casajús, J. A. (2023). Higher physical activity is associated with lower activity limitation: Cross-sectional analyses among the Spanish working population. *Science and Sports*, 38(3), 247-254. <https://doi.org/10.1016/j.scispo.2021.12.010>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



Disponible en ligne sur  
**ScienceDirect**  
[www.sciencedirect.com](http://www.sciencedirect.com)

Elsevier Masson France  
**EM|consulte**  
[www.em-consulte.com](http://www.em-consulte.com)



## ORIGINAL ARTICLE

# Higher physical activity is associated with lower activity limitation: Cross-sectional analyses among the Spanish working population

*Une activité physique plus intense est associée à une limitation d'activité plus faible : analyses transversales parmi la population active espagnole*

R. López-Bueno<sup>a,b,\*</sup>, G.F. López-Sánchez<sup>c</sup>, L. Smith<sup>d</sup>,  
E. Sundstrup<sup>b</sup>, L.L. Andersen<sup>b,e</sup>, J.A. Casajús<sup>f</sup>

<sup>a</sup> Department of Physical Medicine and Nursing, University of Zaragoza, Zaragoza, Spain

<sup>b</sup> National Research Centre for the Working Environment, Copenhagen, Denmark

<sup>c</sup> Division of Preventive Medicine and Public Health, Department of Public Health Sciences, School of Medicine, University of Murcia, Murcia, Spain

<sup>d</sup> Centre for Health, Performance, and Wellbeing, Anglia Ruskin University, Cambridge, UK

<sup>e</sup> Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

<sup>f</sup> Faculty of Health Sciences, University of Zaragoza, Zaragoza, Spain

Reçu le 9 juin 2020 ; accepté le 1<sup>er</sup> décembre 2021

Disponible sur Internet le 7 January 2023

## KEYWORDS

Usual activity ;  
Workplace ;  
Physical exercise ;  
Disability ;  
Sick leave

## Summary

**Objectives.** — Prevention of disability in all its forms is a major public health challenge and promoting a healthy lifestyle with sufficient physical activity (PA) may be a way forward. In contrast to long-term deprivation of usual activities, usually linked to critical injuries or conditions, little is known about shorter periods of those limitations in working populations.

**Equipment and methods.** — We used data from the Spanish National Health Survey 2017 ( $n = 9,885 \geq 17$  years; 47.4% women). Workers self-reported usual activity limitation due to health problems (AL) ( $\leq 14$  days) whereas the International Physical Activity Questionnaire (IPAQ) short form was used to measure PA. After calculating MET-minutes/week, workers

\* Corresponding author.

Adresses e-mail : [rlopezbu@unizar.es](mailto:rlopezbu@unizar.es) (R. López-Bueno), [gfls@um.es](mailto:gfls@um.es) (G.F. López-Sánchez), [lee.smith@aru.ac.uk](mailto:lee.smith@aru.ac.uk) (L. Smith), [esu@nfa.dk](mailto:esu@nfa.dk) (E. Sundstrup), [lla@nfa.dk](mailto:lla@nfa.dk) (L.L. Andersen), [joseant@unizar.es](mailto:joseant@unizar.es) (J.A. Casajús).

**MOTS CLÉS**

Activité habituelle ;  
Lieu de travail ;  
Exercice physique ;  
Handicap ;  
Arrêt de travail

were divided into two categories: 1) Less than 600 MET-minutes/week. 2). At least 600 MET-minutes/week. We conducted a multivariable logistic regression to assess associations. Covariates were age, sex, education, occupation, smoking habits, BMI, diabetes, hypertension, neck pain, low back pain, chronic depression, and anxiety.

**Results.** – The overall prevalence of AL was 10.7%, whereas the overall prevalence of workers performing less than 600 MET-minutes/week of PA was 29.6%. In final adjusted models, those workers performing less than 600 MET-minutes/week of PA subgroup were associated with significantly higher odds for AL (OR 1.33, 95%CI 1.15–1.54). This association was strongest in workers aged 37–43 (OR 1.75, 95%CI 1.27–2.41), and 53–69 years (OR 1.67, 95%CI 1.22–2.28).

**Conclusions.** – The results suggest that reaching  $\geq 600$  MET-minutes/week of PA is associated with lower odds of AL among a general working population, especially among specific age ranges.

© 2022 L'Auteur(s). Publié par Elsevier Masson SAS. Cet article est publié en Open Access sous licence CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Résumé**

**Objectif.** – La prévention du handicap sous toutes ses formes est un défi majeur de santé publique et la promotion d'un mode de vie sain avec une activité physique suffisante peut être une voie à suivre. Contrairement à la privation à long terme des activités habituelles, généralement liée à des blessures ou des conditions critiques, on sait peu de choses sur les périodes plus courtes de ces limitations dans les populations actives. Équipement et méthodes Nous avons utilisé les données de l'Enquête nationale espagnole sur la santé 2017 ( $n = 9885 \geq 17$  ans; 47,4 % de femmes). Les travailleurs ont déclaré une limitation de leurs activités habituelles en raison de problèmes de santé (LA) ( $\leq 14$  jours), tandis que le questionnaire abrégé du Questionnaire international sur l'activité physique (IPAQ) a été utilisé pour mesurer l'activité physique (AP). Après avoir calculé les MET-minutes/semaine, les travailleurs ont été divisés en deux catégories: 1) Moins de 600 MET-minutes/semaine. 2). Au moins 600 MET-minutes/semaine. Nous avons effectué une régression logistique multivariable pour évaluer les associations. Les covariables étaient l'âge, le sexe, l'éducation, la profession, les habitudes tabagiques, l'IMC, le diabète, l'hypertension, les cervicalgies, les lombalgies, la dépression chronique et l'anxiété.

**Résultats.** – La prévalence globale de l'LA était de 10,7 %, tandis que la prévalence globale des travailleurs effectuant moins de 600 MET-minutes/semaine de APPL était de 29,6 %. Dans les modèles ajustés finaux, les travailleurs effectuant moins de 600 MET-minutes/semaine de sous-groupe AP étaient associés à des cotes significativement plus élevées pour LA (OR 1,33, IC à 95 % 1,15–1,54). Cette association était la plus forte chez les travailleurs âgés de 37 à 43 ans (OR 1,75, IC à 95 % 1,27–2,41) et de 53 à 69 ans (OR 1,67, IC à 95 % 1,22–2,28).

**Conclusions.** – Les résultats suggèrent que l'atteinte  $\geq 600$  MET-minutes/semaine de AP est associée à une plus faible probabilité de LA parmi une population active générale, en particulier dans des tranches d'âge spécifiques.

© 2022 L'Auteur(s). Publié par Elsevier Masson SAS. Cet article est publié en Open Access sous licence CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The boundaries between impairment, disability, and functional and activity limitation (AL) are not clear. From a holistic view, the frontiers between health and illness are fragile and represent a continuum and reversible process across life. According to the International Classification of Functioning, Disability, and Health (ICF), “an activity limitation is a difficulty encountered by an individual in executing a task or action” [1]. That could range from slight to a severe deviation regarding quantity or quality in the manner or the extent that is expected of people without the health condition [1]. The notion of AL is usually considered under the umbrella of disability, involving daily activity restrictions (e.g. showering or dressing), and, in some cases, assistance for health care [2]. The severity of that disability depends on a wide variety of factors [3] (i.e. type of disease/disorder, symptoms, and health service support), and comprises a wide range of status from total dependence up to timely

assistance [4]. Further, that disability status preventing from performing daily and usual activities could even lead to a general practitioner to certificate a sickness absence or, in the worst-case scenario, a disability pension [3].

Normally, research about AL has focused more on the types of usual task restrictions, using a long-term reference for those (i.e. 6 months) [5–8]. However, we have no knowledge about shorter-term values of AL ( $\leq 14$  days), which are also important to estimate since these might precede more frequent or more extended periods of AL. It is a well-known fact that either illnesses or critical injuries lead to a reduction or limitation of the usual activity (i.e. working, housework, leisure-time activities, and studying) [9,10]. Thus, work ability and sickness absence might result in a temporary loss of physical or mental capabilities. Because either moderate or severe AL also contribute extensively to higher health care expenditures in the absence of chronic conditions [11], those assuming high burden (i.e. governments and companies) want to reduce it. However,

not only it is a question of financial costs but also a health and well-being issue as global AL (i.e. health disorders and conditions impacting people's usual activities) has also been associated with mortality [5].

While AL has been previously studied with older populations or particular conditions during long periods [12–16], little is known about the AL prevalence among other specific populations such as workers. An onset of AL might involve restrictions related to work, although it does not necessarily lead to sickness absence, because it could solely occur during leisure or non-working days (e.g. affecting housework, education, and leisure-time activities). Further, suffering from an illness or condition does not strictly mean to remain in bed, since prior research has reported a third of workers had gone to work two or more times a year when feeling sick instead of taking sick leave [17]. Thus, to investigate both prevalence and potential ways of preventing AL among a general working population covers a research gap between sickness absence and sickness presenteeism, two of the highest-ranked costs for both governments and companies [18].

On the other hand, living a healthy lifestyle could contribute to reducing AL since physical activity (PA), normal BMI, and smoking deprivation have shown to associate with better-perceived health [19]. Particularly, PA has been found to significantly reduce all-cause mortality [20], long-term sickness absence [21], and presenteeism [22] risks; therefore, we hypothesised that higher PA might associate with a lower prevalence of AL. Also, vigorous PA has been estimated to achieve higher all-cause mortality risk reductions [23], thus, the intensity of PA might play a critical role in this association. Regarding other possible variables influencing AL, we assumed that those behind the highest risk for sickness absence [24], such as cardiovascular diseases (i.e. hypertension), and musculoskeletal (i.e. back pain) or mental disorders (i.e. chronic depression and anxiety) [24] might also influence AL.

The primary aim of this research was to examine the association of different levels of PA with AL ( $\leq 14$  days) among the general Spanish working population.

## 2. Methods

This study consists of a survey assessing general health among a Spanish population of citizens, in which the 2017 round was analyzed to investigate potential associations between PA and short AL. In order to ensure comprehensive reporting of the data for the present cross-sectional study, the STROBE guidelines were followed [25].

### 2.1. The survey

Data from the Spanish National Health Survey 2017 was employed for this analysis. Data were collected through a survey carried out in Spain from October 2016 to October 2017 [26]. A stratified three-stage sampling considering census sections, family dwellings, and adults (15 years or more) was respectively implemented. The dwellings were chosen using systematic sampling, whereas the random Kish method was used to select the participants who were going to complete the Adult Questionnaire. A computer-assisted

personal interview (CAPI) was conducted in the homes of the selected participants, who were assisted by trained interviewers. A total of 30.1% ( $n=11,287$ ) of the chosen dwellings did not reply to the survey due to different reasons (i.e. empty dwelling, absence, refusal, or inability to answer). As a result, a representative sample of the adult population resident in Spain was collected and consisted of 23,089 adults aged 15–103 years. All the participants signed an informed consent form before responding to the survey.

Because the age group of adults  $\geq 70$  years did not fulfill the IPAQ (International Physical Activity Questionnaire) short form (i.e. the tool used for assessing PA) embedded in the Adult Questionnaire, we excluded them for the present analyses ( $n=5,310$ ). Of the remaining population, those participants not working at data collection were also excluded ( $n=7,894$ ). Thus, a total of 9,885 workers, comprising a wide range of job roles (i.e. from white and blue-collar workers to self-employed) were included in the present study.

### 2.2. Activity limitation (Outcome)

Measures on AL were obtained through the following question "Have you had to reduce or limit your usual activity during the last two weeks, for at least half a day, due to either one or several pains or symptoms?". Those participants responding "yes" to this question were considered to have had an AL onset during that term. According to the survey methodology, that usual activity comprised working activities, housework, attendance to learning centres or leisure-time events usually shared with family and friends.

### 2.3. Physical activity (Exposure)

The short version of the IPAQ served to assess PA [27]. The IPAQ has proved to have good validity ( $\rho=0.30$ , 95%CI 0.23–0.36) and reliability (Spearman's  $\rho=0.81$ , 95%CI 0.79–0.82) when used in different adult populations worldwide [27]. Total PA MET (Metabolic Equivalent of Task, i.e. a caloric expenditure unit) minutes/week were calculated adding the METs from three types of PA with the formula: *sum of Vigorous + Moderate + Walking MET-minutes/week scores* [28]. To further check the PA association with AL, we also categorised PA into quartiles regarding METS.

In consonance with the data processing and analysis protocol of the IPAQ, the respondents were divided into two categories: "(1) fewer than 600 MET-minutes/week" and "(2) at least 600 MET-minutes/week" [28], which is equivalent to achieve the current WHO guidelines.

### 2.4. Covariates

According to research in the field of occupational health, the selected covariates included sociodemographic (age, sex, education, and job role), lifestyle (body mass index and smoking), and health variables (diabetes, hypertension, neck pain, low back pain, chronic depression, and anxiety) [29,30].

Measurements on BMI (Body Mass Index, i.e. weight in kilograms divided by height in squared meters), in which obesity was set at  $BMI \geq 30 \text{ kg/m}^2$ , derived from the

self-reported height and weight. The smoking status was categorised as a current smoker, former smoker, and never. Education was divided into three categories regarding the highest achieved level, corresponding the highest rank to university studies ( $\leq$  primary, secondary, and  $\geq$  tertiary). The categorisation of job role (168 different groups) was made in accordance with the Spanish national list of occupations [31].

Regarding the health-related covariates, those who answered "yes" to the question "Have you suffered from neck pain within the last twelve months?" were considered to have neck pain. The same question and procedure were used for assessing the prevalence of the remaining health variables.

## 2.5. Statistical analysis

We conducted a statistical analysis using SPSS 22.0. Differences in the prevalence of AL by sample characteristics were assessed by Chi<sup>2</sup> tests. The association between PA (exposure) and AL (outcome) was estimated through multi-variable logistic regression analysis conducted for the whole sample as well as by age subgroups (15–36, 37–43, 44–52, and 53–69 years). Age-segmented analyses were adjusted for sex, BMI, education, smoking, job role, diabetes, hypertension, neck pain, low back pain, depression, and anxiety, while the analysis for the overall sample was also adjusted for age. Except for age, included as a continuous variable, all variables were included in the models as categorical variables. Participants with missing data ( $n=190$ ) were not considered for the analyses. Cut-off points for age-segmentation and METS were estimated through equal percentile based on explored data option from the SPSS visual grouping procedure. As a result of the logistic regression analyses, we calculated odds ratios (ORs) with 95% confidence intervals (CIs). The level of statistical significance was set at  $P<0.05$ .

## 3. Results

**Table 1** shows the baseline characteristics of the study population. The cohort includes a population of Spanish general workers (47.4% women) on average 44.4 (SD 10.4) years old. Concerning lifestyle, 56.8% of the cohort is linked to smoking habits, and 51.1% show overweight or obesity. The overall prevalence of AL is 10.7%, whereas the overall prevalence of people doing less than 600 MET-minutes/week of PA is 29.6%. Regarding PA guidelines subgroups, workers doing less and more than 600 MET-minutes/week of PA present a 13.2% and a 9.7% prevalence of AL, respectively. In estimates without adjustment, women, former smokers, less PA, diabetes, hypertension, neck pain, low back pain, depression, and anxiety were associated with a significantly higher prevalence of AL.

**Fig. 1** shows an upward trend in the prevalence of AL, from 10.9% to 17.4%, when age increasing with those participants in the first PA quartile. The fourth PA quartile observes the less prevalent values for AL for the three older subgroups.

When performing adjusted overall analysis, the less than 600 MET-minutes/week of PA subgroup was associated with

significantly higher odds (OR 1.33, 95%CI 1.15–1.54) for AL (**Table 2**). Advanced age, former smoker, hypertension, neck pain, low back pain, depression, and anxiety were also associated with significantly higher odds for AL (**Table 2**). The full adjusted association between PA and AL remains significant among those subgroups formed by workers aged 37 to 43, and 53 to 69 when age-stratified (**Table 3**).

## 4. Discussion

In the present sample of Spanish general workers, those workers performing less than 600 MET-minutes/week of PA had significantly higher odds of AL. In the age-stratified analyses, AL remained only significant for both 1) 37–43 years, and 2) 53–69 year's subgroups achieving PA guidelines. The prevalence of AL observed a clear trend to diminish with the ageing process when achieving very high levels of PA, whereas those workers performing low levels of PA presented the highest odds of AL in the three subgroups with older workers (**Fig. 1**). Therefore, in order to reduce AL values, PA strategies aiming at the older population of workers would be desirable.

Prevalence of several of the most common cardiovascular, metabolic, physical, and mental disorders among the workforce observed significant values regarding AL for the present sample. Except for diabetes, all the diseases showed significantly increased odds for AL in the adjusted model, although both low back pain and depression presented the highest significant association with AL.

Owing to the use of the IPAQ short-form tool, the present study focus PA on the quantitative (i.e. total amount of METS) instead of the qualitative element (i.e. intensity) of the exposure. Thus, achieving a meaningful value of METS such as those recommended in the WHO guidelines [32], regardless of the way those have been achieved (i.e. different possible combinations of high, moderate PA and walking to get the same value), could result in a significant reduction of odds for AL. Although we have not found specific research investigating the association between PA and AL, several studies have suggested the inverse relationship of both quantitative and qualitative aspects of PA with short ( $\leq$  14 days) and long-term sickness absence ( $>$  14 days) [33,34], even with a different population of workers [21,35].

When adjusted for occupational class (i.e. job role), our proposed model did not show significant modifications in the association between PA and AL; this finding contrasts with those observed in other studies carried out with a general population of Danish workers, in which high levels of occupational PA were observed to significantly increase the risk of worse global health [21].


Regarding age, we found a clear trend to increase odds for AL when adjusted by the rest of the variables. When unadjusted and categorised into quartiles (**Fig. 1**), the odds of AL remained higher for the older workers performing low PA and lower for those older workers achieving very high levels of PA. Thus, the age factor seems to take its part when explaining AL, and older but not younger workers could be more sensitive to PA. Similar findings were observed among female eldercare workers, in which physical exertion increased the disability pension risk in older but not in younger workers [36].

**Table 1** Characteristics of the study population.

	N	%	Mean	SD	Activity limitation <sup>a</sup>		P-value <sup>b</sup>
					N	%	
Age (years)	9885		44.4	10.4			
Sex							<0.001
Men	5199	52.6			483	9.3	
Women	4686	47.4			580	12.4	
Education							0.620
≤ Primary	970	9.8			100	10.3	
Secondary	6001	60.7			660	11.0	
≥ Tertiary	2914	29.5			303	10.4	
Smoking							0.010
Current	2951	29.9			333	11.3	
Former	2656	26.9			315	11.9	
Never	4269	43.2			414	9.7	
Missing	9	0.0					
BMI ( $\text{kg} \cdot \text{m}^{-2}$ )							0.241
Low	180	1.9			20	11.1	
Normal	4556	47.0			500	11.0	
Overweight	3600	37.1			364	10.1	
Obese	1359	14.0			164	12.1	
Missing	190	1.9					
Physical activity (MET-minutes/week) WHO guidelines							<0.001
< 600	2930	29.6			386	13.2	
≥ 600	6862	69.5			665	9.7	
Missing	93	0.9					
Physical activity (MET-minutes/week)							<0.001
Low (0–480)	2514	25.5			349	13.9	
Moderate (481–1386)	2914	29.5			299	10.3	
High (1387–2826)	1932	19.5			190	9.8	
Very high (2827 or more)	2432	24.6			213	8.8	
Missing	93	0.9					
Diabetes							0.026
No	9585	97.0			1019	10.6	
Yes	300	3.0			44	14.7	
Hypertension							<0.001
No	8761	88.6			895	10.2	
Yes	1122	11.4			167	14.9	
Missing	2	0.0					
Neck pain							<0.001
No	8702	88.0			768	8.8	
Yes	1183	12.0			295	24.9	
Low back pain							<0.001
No	8304	84.0			689	8.3	
Yes	1581	16.0			374	23.7	
Depression							<0.001
No	9514	96.3			926	9.7	
Yes	368	3.7			136	37.0	
Missing	3	0.0					
Anxiety							<0.001
No	9362	94.7			914	9.8	
Yes	523	5.3			149	28.5	

BMI: Body Mass Index.

<sup>a</sup> Number (and percentage) of individuals with that sample characteristic who have had an activity limitation spell<sup>b</sup> P-value was calculated with Chi<sup>2</sup> tests.

**Figure 1** Prevalence of activity limitation (AL) by level of physical activity (PA) and by age groups.**Table 2** Association of physical activity and other covariates with activity limitation (outcome) estimated by multivariable logistic regression.

Characteristic		Activity limitation	
Physical activity (MET-minutes/week)	< 600	1.33 <sup>c</sup>	[1.15, 1.54]
	≥ 600	ref	
Age (years)	Per unit increase	0.98 <sup>b</sup>	[0.98, 0.99]
Sex	Men	0.91	[0.76, 1.09]
	Women	ref	
Education	≤ Primary	0.91	[0.65, 1.27]
	Secondary	1.04	[0.83, 1.31]
	≥ Tertiary	ref	
Smoking	Never	ref	
	Former	1.22 <sup>a</sup>	[1.03, 1.45]
	Current	1.11	[0.94, 1.32]
BMI ( $\text{kg m}^{-2}$ )	Low	ref	
	Normal	1.12	[0.67, 1.87]
	Overweight	0.95	[0.56, 1.61]
	Obese	1.06	[0.61, 1.82]
Diabetes	No	ref	
	Yes	1.19	[0.82, 1.73]
Hypertension	No	ref	
	Yes	1.32 <sup>a</sup>	[1.06, 1.63]
Neck pain	No	ref	
	Yes	1.88 <sup>c</sup>	[1.55, 2.28]
Low back pain	No	ref	
	Yes	2.31 <sup>c</sup>	[1.94, 2.76]
Depression	No	ref	
	Yes	2.99 <sup>c</sup>	[2.23, 4.01]
Anxiety	No	ref	
	Yes	1.46 <sup>b</sup>	[1.11, 1.91]

Estimates are odds ratio [95% confidence interval]. Models are adjusted for all variables in the Table and job role.

<sup>a</sup>  $P < 0.05$ .

<sup>b</sup>  $P < 0.01$ .

<sup>c</sup>  $P < 0.001$ .

**Table 3** Association between different levels of physical activity and activity limitation (outcome) estimated by multivariable logistic regression.

Age (years)	Association between physical activity (< 600 MET-minutes/week) and activity limitation	
17–36	1.09	[0.78, 1.51]
37–43	1.75**	[1.27, 2.41]
44–52	1.10	[0.82, 1.48]
53–69	1.67**	[1.22, 2.28]

Estimates are odds ratio [95% confidence interval]. Models are adjusted for sex, education, BMI, smoking, job role, diabetes, hypertension, neck pain, low back pain, depression and anxiety. \*  $P < 0.05$ . \*\*  $P < 0.01$ . \*\*\*  $P < 0.001$ .

Several of the health confounders used in our study have been ranked as one of the most prevalent among the workforce [37,38]. However, despite the high influence of hypertension, back pain, and mental disorders in the proposed model, the association between PA and AL remained significant for this study. Furthermore, the significantly increased AL odds for five out of six of the health variables suggests that those highlighted health risk factors for long-term sickness absence could also be substantial for AL. Except for the category of former smokers, current smokers or other confounders such as BMI or education did not find a substantial increase of odds for AL when adjusted for the remaining variables. These results differ from the findings of other studies outlining the association of both current smoking habits and obesity with a higher risk of general sickness absence in specific populations of workers (i.e. local employees and health workers) [39,40].

Overall, achieving a minimum of 600 weekly METS of PA may contribute to reducing AL among a general population of workers. Health strategies aiming at older workers performing low levels of PA or those suffering from anxiety, depression, or back pain might reduce AL. Future research on AL might investigate its use as a potential marker for the two of them, although longitudinal research would be first required to confirm the suggested association.

The strengths of the current study comprise using a large random sample of Spanish workers, as well as measuring the exposure variable with a reliable tool. To further avoid a possible recall bias in the outcome variable, only onsets, instead of days, of AL were contemplated. In addition, a representative set of confounders covering a wide range of possible risk factors for AL were considered for the implemented logistic model. However, several limitations make this research to be carefully considered. First, the observational design of the study does not permit the inspection of a causal relationship between PA and AL. Thus, low levels of PA may lead to AL, but the reverse association may also be present, i.e. those with activity limitations are more prone to not being sufficiently physically active. Second, the subjective nature of the measured variables, which represents an inherent risk of information bias. Particularly, the question used to estimate the main outcome (AL) was not validated, which can exacerbate the aforementioned

information bias. Also, self-reported data on PA usually lead to overestimating overall PA. Last, according to the IPAQ protocol, the final values of that tool do not discriminate among vigorous or moderate PA, which limits the scope of the study.

## 5. Conclusions

All things considered, the results of the present study suggest a determined level of PA ( $\geq 600$  METS/week) to associate with lower AL, particularly among older workers. Health strategies aiming at this specific population might reduce levels of AL.

## Disclosure of interest

The authors declare that they have no competing interest.

## Funding

The survey was funded by the Spanish Ministry of Health, Social Services and Equality, and the National Institute of Statistics as a part of an assessment program on general health.

## Références

- [1] World Health Organization (WHO). International classification of functioning, disability and health world. Geneva; 2001. Disponible sur: <https://apps.who.int/iris/bitstream/handle/10665/42407/9241545429.pdf> [cité le 16 avr 2019].
- [2] Van Oyen H, Heyden J, Perenboom R, Jagger C. Monitoring population disability: evaluation of a new Global Activity Limitation Indicator (GALI). Soz Praventivmed 2006;51:153–61.
- [3] Morgell R, Backlund LG, Arrelöv B, Strenger LE, Nilsson GH. Health problems and disability in long-term sickness absence: ICF coding of medical certificates. BMC Public Health 2011;11.
- [4] Tennant A, Penta M, Tesio L, Grimby G, Thonnard JL, Slade A, et al. Assessing and adjusting for cross-cultural validity of impairment and activity limitation scales through differential item functioning within the framework of the Rasch model: the PRO-ESOR project. Med Care 2004;42:37–48.
- [5] Berger N, Van der Heyden J, Van Oyen H. The global activity limitation indicator and self-rated health: two complementary predictors of mortality. Arch Public Heal 2015;73:1–7.
- [6] Jagger C, Gillies C, Cambois E, Van Oyen H, Nusselder W, Robine JM. The Global Activity Limitation Index measured function and disability similarly across European countries. J Clin Epidemiol 2010;63:892–9.
- [7] Berger N, Van Oyen H, Cambois E, Fouweather T, Jagger C, Nusselder W, et al. Assessing the validity of the global activity limitation indicator in fourteen European countries. BMC Med Res Methodol 2015;15.
- [8] von der Lippe E, Fehr A, Lange C. Limitations to usual activities due to health problems in Germany. J Heal Monit 2017;2:84–90.
- [9] Ohtake PJ, Coffey Scott J, Hinman RS, Lee AC, Smith JM. Impairments, activity limitations and participation restrictions experienced in the first year following a critical illness: protocol for a systematic review. BMJ Open 2017;7:1–5.

- [10] Xiang H, Leff M, Stallones L. Non-fatal injuries among adults with activity limitations and participation restrictions. *Inj Prev* 2005;11:157–62.
- [11] Van Der Heyden J, Van Oyen H, Berger N, De Bacquer D, Van Herck K. Activity limitations predict health care expenditures in the general population in Belgium. *BMC Public Health* 2015;15:1–12.
- [12] Abraido-Lanza AF, White K, Armbrister AN, Link BG. Health status, activity limitations, and disability in work and housework among Latinos and non-Latinos with arthritis: an analysis of national data. *Arthritis Care Res* 2006;55:442–50.
- [13] Thyberg I, Hass UAM, Nordenskiöld U, Gerde B, Skogh T. Activity limitation in rheumatoid arthritis correlates with reduced grip force regardless of sex: The Swedish TIRA project. *Arthritis Care Res* 2005;53:886–96.
- [14] Goverover Y, Strober L, Chiaravalloti N, DeLuca J. Factors that moderate activity limitation and participation restriction in people with multiple sclerosis. *Am J Occup Ther* 2015;69:1–9.
- [15] Ibikunle lanrewaj PO, Oladipo kunday SE, Chukwu goz JN, Odole hristian AC, Okeke foem AI. Establishing the reliability and construct validity of the Igbo version of Screening Activity Limitation and Safety Awareness scale in persons with Hansen disease. *Lepr Rev* 2015;86:220–8.
- [16] Moon JH. Factors affecting activity limitation in the elderly: data processed from the Korea National Health and Nutrition Examination Survey, 2016. *Osong Public Heal Res Perspect* 2019;10:117–22.
- [17] Aronsson G, Gustafsson K, Dallner M. Sick but Yet at Work. An empirical study of sickness presenteeism sick but yet at work. An sickness presenteeism empirical study of 2014;54:502–9.
- [18] Nagata T, Mori K, Ohtani M, Nagata M, Kajiki S, Fujino Y, et al. Total health-related costs due to absenteeism, presenteeism, and medical and pharmaceutical expenses in Japanese employers. *J Occup Environ Med* 2018;60:e273–80.
- [19] Molarius A, Berglund K, Eriksson C, Lambe M, Nordström E, Eriksson HG, et al. Socioeconomic conditions, lifestyle factors, and self-rated health among men and women in Sweden. *Eur J Public Health* 2007;17:125–33.
- [20] Saint-Maurice PF, Coughlan D, Kelly SP, Keadle SK, Cook MB, Carlson SA, et al. Association of leisure-time physical activity across the adult life course with all-cause and cause-specific mortality. *JAMA Netw Open* 2019;2:e190355.
- [21] Holtermann A, Hansen JV, Burr H, Søgaard K, Sjøgaard G. The health paradox of occupational and leisure-time physical activity. *Br J Sports Med* 2012;46:291–5.
- [22] Walker TJ. The longitudinal relation between self-reported physical activity and presenteeism. *J Med Virol* 2010;82:175–85.
- [23] Gebel K, Ding D, Chey T, Stamatakis E, Brown WJ, Bauman AE. Effect of moderate to vigorous physical activity on all-cause mortality in middle-aged and older Australians. *JAMA Intern Med* 2015;175:970–7.
- [24] Sundstrup E, Due Jakobsen M, Mortensen OS, Andersen LL. Joint association of multimorbidity and work ability with risk of long-term sickness absence: a prospective cohort study with register follow-up. *Scand J Work Environ Heal* 2017;43:146–54.
- [25] von Elm E, Altman DG, Egger M, Pocock SJ, Götzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Epidemiology* 2007;18:800–4.
- [26] Ministerio de Sanidad Servicios Sociales e Igualdad Instituto Nacional de Estadística. Spanish National Health Survey 2017: Methodology; 2017. Disponible sur: [https://www.ine.es/en/metodologia/t15/t153041917\\_en.pdf](https://www.ine.es/en/metodologia/t15/t153041917_en.pdf) [cité le 25 avr 2019].
- [27] Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-Country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381–95.
- [28] Ipaq. Guidelines for data processing and analysis of the international physical activity questionnaire (IPAQ)—short and long forms. Ipaq; 2005. p. 1–15. Disponible sur: <https://sites.google.com/site/theipaq/scoring-protocol> [cité le 15 avr 2019].
- [29] López-Bueno R, Sundstrup E, Vinstrup J, Casajús JA, Andersen LL. High leisure-time physical activity reduces the risk of long-term sickness absence. *Scand J Med Sci Sport* 2020;30:939–46.
- [30] López-Bueno R, Bláfoss R, Calatayud J, López-Sánchez GF, Smith L, Andersen LL, et al. Association between physical activity and odds of chronic conditions among workers in Spain. *Prev Chronic Dis* 2020;17:200105.
- [31] Instituto Nacional de Estadística. Clasificación Nacional de Ocupaciones (CNO). Inst Nac Estadística; 2011. Disponible sur: [https://www.ine.es/dyngs/INEbase/es/operacion.htm?c=Estadistica\\_C&cid=1254736177033&menu=ultiDatos&idp=1254735976614](https://www.ine.es/dyngs/INEbase/es/operacion.htm?c=Estadistica_C&cid=1254736177033&menu=ultiDatos&idp=1254735976614) [cité le 9 avr 2019].
- [32] World Health Organization (WHO). Global recommendations on physical activity for health. Geneva; 2014. Disponible sur: <https://www.who.int/publications/i/item/9789241599979> [cité le 11 avr 2019].
- [33] Losina E, Yang HY, Deshpande BR, Katz JN, Collins JE. Physical activity and unplanned illness-related work absenteeism: data from an employee wellness program. *PLoS One* 2017;12:1–13.
- [34] Lahti J, Laaksonen M, Lahelma E, Rahkonen O. The impact of physical activity on sickness absence. *Scand J Med Sci Sport* 2010;20:191–9.
- [35] Tolonen A, Rahkonen O, Lahti J. Leisure-time physical activity and direct cost of short-term sickness absence among Finnish municipal employees. *Arch Environ Occup Heal* 2017;72:93–8.
- [36] Andersen L, Villadsen E, Clausen T. Influence of physical and psychosocial working conditions for the risk of disability pension among healthy female eldercare workers: Prospective cohort. *Scand J Public Health* 2020;48:460–7.
- [37] de Vroome EMM, Uegaki K, van der Ploeg CPB, Treutlein DB, Steenbeek R, de Weerd M, et al. Burden of sickness absence due to chronic disease in the Dutch workforce from 2007 to 2011. *J Occup Rehabil* 2015;25:675–84.
- [38] Hassard J, Teoh K, Cox T, Dewe P, Cosmar M, Gründler R, et al. Calculating the cost of work-related stress and psychosocial risks. Luxembourg; 2014.
- [39] Laaksonen M, Piha K, Martikainen P, Rahkonen O, Lahelma E. Health-related behaviours and sickness absence from work. *Occup Environ Med* 2009;66:840–7.
- [40] Torres Lana A, Cabrera de León A, Marco García MT, Aguirre Jaime A. Smoking and sickness absence among public health workers. *Public Health* 2005;119:144–9.