



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Psychophysical evaluation of the effect of electrode location on sensations during electrocutaneous stimulation

Geng, Bo; Yoshida, Ken; Jensen, Winnie

Published in:

Abstracts of the XVIII Congress of the International Society of Electrophysiology and Kinesiology, ISEK 2010, 16-19 June 2010, Aalborg, Denmark [CD-ROM]

Publication date:

2010

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Geng, B., Yoshida, K., & Jensen, W. (2010). Psychophysical evaluation of the effect of electrode location on sensations during electrocutaneous stimulation. In D. Falla, & D. Farina (Eds.), *Abstracts of the XVIII Congress of the International Society of Electrophysiology and Kinesiology, ISEK 2010, 16-19 June 2010, Aalborg, Denmark [CD-ROM]* (pp. No. P390). Department of Health Science and Technology. Aalborg University.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

PSYCHOPHYSICAL EVALUATION OF THE EFFECT OF ELECTRODE LOCATION ON SENSATIONS DURING ELECTROCUTANEOUS STIMULATION

Geng B¹, Yoshida K^{1,2}, Jensen W¹

¹ Aalborg University, Aalborg, Denmark

² Indiana University-Purdue University Indianapolis, Indianapolis, U.S.A.

E-mail: bogeng@hst.aau.dk, yoshidak@iupui.edu, wj@hst.aau.dk

AIM: Electrical stimulation can potentially be used to create sensations for use as feedback in advanced prosthetics. This study aims to evaluate the psychophysical effect of electrode location on sensations during electrocutaneous stimulation.

METHODS: 16 healthy subjects were included in the study (Aged 22-36). Five electrodes were placed around the left forearm 5 cm distant to the antecubital crease, as follows: 1) Electrode 1 (E1) over the median nerve, 2) Electrode 2 (E2) laterally adjacent to E1, 3) Electrode 3 (E3), 4 (E4), and 5 (E5) equally spaced between E2 and E1. Fixed-amplitude biphasic square pulse stimuli at three times of sensory threshold were applied through the five electrodes. Immediately following each stimulus, subjects were presented a questionnaire to score the perceived sensation types and quality. A visual analog scale (VAS) was used to measure the sensation strength.

RESULTS: All the 16 subjects reported tactile sensations (press or pulsing) at the E1 site, while fewer subjects reported tactile sensations at the E3 or the E4 site. The lowest sensation strength reported by the subjects appeared at site E4. Table 1 describes the responses of tactile sensation in detail. Based on the Friedman test, the five electrode sites show statistically significant differences in the perceived strength of tactile sensation ($p=0.003$). Moreover, clustering of sensation types indicated that E3 and E4 tend to evoke similar sensation types, while E1, E2 and E5 fall into another cluster (figure 1 (a)). Clustering of sensation quality gives the same cluster result (figure 1(b)).

CONCLUSION: Electrical stimulation to the different sides of the forearm evoked different responses. Stimulation of the ventral side was more likely to induce tactile sensations. Moreover, the magnitude of the perceived sensations was stronger on the ventral side compared to the dorsal side at the same level of stimulation. It seems to be more appropriate to induce sensation feedback on the ventral than the dorsal side of the forearm.

Table 1: The number of subjects who reported ‘tactile’ sensation, the mean of the tactile sensation strength (VAS) at the five electrode sites.

Electrode site	E1	E2	E3	E4	E5
Number of subjects	16	13	8	9	14
Mean of VAS	2.54	2.95	2.80	1.63	2.10




Figure 1: Clustering of the five electrode sites (16 subjects): (a) Sensation type. (b) Sensation quality.