



A study of the applicability of xenon clathrates for the preservation of human stem cells and skin fragments

Ponomarev, Alexander; Gurevich, Leonid; Makeev, Oleg; Zvereva, Anastasia

Published in:

International Summer School on Application of Scanning Probe Microscopy in Life Sciences, Soft Matter and Nanofabrication

Publication date:

2013

Document Version

Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Ponomarev, A., Gurevich, L., Makeev, O., & Zvereva, A. (2013). A study of the applicability of xenon clathrates for the preservation of human stem cells and skin fragments. In *International Summer School on Application of Scanning Probe Microscopy in Life Sciences, Soft Matter and Nanofabrication* River Publishers. http://same-nano.com/Abstracts2013/School%202013_Abtract_Ponomarev_v3.pdf

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

A study of the applicability of xenon clathrates for the preservation of human stem cells and skin fragments

Ponomarev A.^{1}, Gurevich L.², Makeev O.¹, Zvereva A.¹*

¹⁾ Ural State Medical Academy, Ekaterinburg, Russia

²⁾ Aalborg University, Aalborg, Denmark

*) E-mail: biopreservation@gmail.ru

Rapid development of regenerative medicine calls for new approaches to the problem of long-term preservation of cells, tissues and organs. This is one of the key issues in, e.g., transplantology and tissue engineering. The existing cryopreservation techniques are largely limited by the toxic effects that commonly used cryoprotector components such as DMSO [5] and glycerol [4] exhibit on the stored bio-objects [1] and eventual recipients [2]. Xenon gas can be a viable alternative to DMSO due to its ability to form clathrates in aqueous solutions, which effectively prevent the formation of ice crystals and therefore limit the damage to the cells [3]. Due to its low toxicity, xenon gas is widely used nowadays in anesthesia as well as a medicine for treatment of various diseases [6]. Several earlier studies have demonstrated the possibility to use xenon for cryopreservation of cell suspensions [7]. In another work, a mixture of xenon and oxygen at a pressure of 6 to 7 atm has been used for short-term cryopreservation of mammalian tissues [8]. However, to the authors' knowledge, all previous works have studied either cell suspensions or animal tissues, while very limited information has been collected on human tissues. Moreover, no quantitative data on the properties of clathrates used for conservation and characteristics of the preserved cells (e.g. viability) have been provided. The purpose of this study is to explore the possibility of using xenon clathrates as a cryopreservation agent for biological objects of human origin, such as human stem cells and skin tissues.

A custom-designed sealed chamber equipped with pressure and temperature sensors to monitor the parameters of clathrate formation and a window to visually register the clathrate point in the system was built for the experiments. Human stem cells and skin samples were preserved in clathrates for 1 and 2 days, respectively. The damaging effect of xenon clathrates on the stages of clathrate formation and storage was evaluated using an environmental scanning electron microscope (Zeiss Evo) equipped with a cryostage. The viability of the skin samples after cryopreservation in xenon clathrates was evaluated by expansion of fibroblasts in DMEM with 10% fetal bovine serum [9]. Histology of the clathrates-preserved and reference (intact) skin samples was performed using standard techniques [10]

The studies showed no visible damaging effect of xenon clathrates on stem cells during clathrate preservation and storage (Fig. 1). However, the cells were almost completely destroyed during the re-activation stage. On the hand, the skin samples preserved in xenon clathrates did not exhibit any histological difference from the reference specimens. The skin samples stored in xenon clathrates exhibited a steady output of fibroblasts with positive dynamics and did not differ from the reference specimens after 25 days of culture (Fig. 2).

In conclusion, our preliminary studies demonstrated that xenon clathrates can be used as a preservation agent for storage of human skin. Preservation, storage and re-activation stages exhibited virtually no impact on the skin samples. At the same time, while the stem cells were seemingly intact at the time of preservation and storage, they were almost completely destroyed during re-activation. This might indicate that the suspension of stem cells is less resistant to decompression as compared to skin samples and requires some adjustment in our procedure. In the future work we plane to address this issue, evaluate the percentage of damaged cells in the

current and future cryopreservation protocols and make a detailed description of the histological skin fragments preserved in xenon clathrates.




Fig. 1. Stem cells preserved in xenon clathrates. Note the intact cell membrane, no ice crystals present.




Fig. 2. The expansion of the fibroblasts of the skin sample after preservation in Xe clathrates. Tendency to form a monolayer on the 25th day of cultivation was observed

References.

1. Hunt C.J., Armitage S.E., Pegg D.E. Cryopreservation of umbilical cord blood: 2. Tolerance of CD34+ cells to multimolar dimethyl sulphoxide and effect of cooling rate on recovery after freezing and thawing // *Cryobiology*.-2003.-Vol.46.-P.76-87.
2. Higman MA, Port JD, Beauchamp NJ Jr, Chen AR. Reversible leukoencephalopathy associated with re-infusion of DMSO preserved stem cells. *Bone Marrow Transplant*.2000. 26, 797–800.
3. Rodin V.V., Isangalin F.S., Volkov V.J. Structure aqueous protein solutions in the presence of xenon clathrates. *Cryobiology and cryomedicine*. 1984; 14, 3-7. [In Russian]
4. Janz Fde L, Debes Ade A, Cavaglieri Rde C, Duarte SA, Romão CM, Morón AF, Zugaib M, Bydlowski SP. Evaluation of distinct freezing methods and cryoprotectants for human amniotic fluid stem cells cryopreservation. *J Biomed Biotechnol*. 2012;2012:649353.
5. Abbruzzese L, Agostini F, Durante C, Toffola RT, Rupolo M. Long term cryopreservation in 5% DMSO maintains unchanged CD34(+) cells viability and allows satisfactory hematological engraftment after peripheral blood stem cell transplantation. *Vox Sang*. 2013;105(1):77-80.
6. Stoppe C, Rimek A, Rossaint R, Rex S, Stevanovic A, Schälte G, Fahlenkamp A, Czaplik M, Bruells CS, Daviet C, Coburn M. Xenon consumption during general surgery: a retrospective observational study.// *Med Gas Res*. 2013 Jun 11;3(1):12.
7. Rodin V.V. Volkov V.Y., Isangalin F.S. "Freezing" of the bacterial suspensions at positive temperatures xenon clathrates. NMR data. Cryodamage and cryoprotection mechanisms of biological objects: Proc. 2nd Vsesoyuz. konf. , Kharkov, 1984, Vol.1, p.199. [In Russian]
8. Sheleg S, Hixon H, Cohen B, Lowry D, Nedzved M. Cardiac mitochondrial membrane stability after deep hypothermia using a xenon clathrate cryostasis protocol - an electron microscopy study. *Int J Clin Exp Pathol*. 2008 Jan 1;1(5):440-7.
9. Tanaka Y, Matsuo K, Yuzuriha S. Long-term histological comparison between near-infrared irradiated skin and scar tissues. *Clin Cosmet Investig Dermatol*. 2010 Nov 25;3:143-9.
10. Pollard JW. Basic cell culture. *Methods Mol Biol*.1990;5:1-12.