



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Topological Model of the Dissolution Kinetics of Silicate Glasses

Wang, Mengyi; Pignatelli, Isabella; Oey, Tandr ; Smedskj r, Morten Mattrup; Mauro, John C.; Sant, Gaurav; Bauchy, Mathieu

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Wang, M., Pignatelli, I., Oey, T., Smedskj r, M. M., Mauro, J. C., Sant, G., & Bauchy, M. (2017). *Topological Model of the Dissolution Kinetics of Silicate Glasses*. Abstract from 12th Pacific Rim Conference on Ceramic and Glass Technology, Waikoloa, Hawaii, United States.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Topological Model of the Dissolution Kinetics of Silicate Glasses

Mengyi Wang<sup>1</sup>, Isabella Pignatelli<sup>1</sup>, Tandr  Oey<sup>1</sup>, Morten M. Smedskjaer<sup>2</sup>, John C. Mauro<sup>3</sup>, Gaurav Sant<sup>1</sup>,  
Mathieu Bauchy<sup>1</sup>

<sup>1</sup>University of California, Los Angeles, Los Angeles, CA, United States.

<sup>2</sup>Aalborg University, Aalborg, Denmark.

<sup>3</sup>Corning Incorporated, Corning, NY, United States.

Understanding and predicting the dissolution rate of silicate glasses is of primary importance for various applications, including bioactive glasses and borosilicate wastefoms. However, the mechanism of silicates' dissolution – and its rate-limiting step – remains poorly understood. In particular, present models linking the composition and structure of silicate glasses to their dissolution rate in a given solvent have remained largely empirical thus far. Here, based on vertical scanning interferometry (VSI), we study the dissolution of a family of borosilicate glasses under several values of pH. In addition, we parametrized a new transferable inter-atomic potential to assess the structure of these glasses from molecular dynamics (MD) simulations. From a detailed analysis of the simulated structures, we demonstrate that the kinetics of the dissolution is controlled by the topology of the atomic network. Finally, we propose a new topological model of the dissolution, which is shown to offer realistic predictions of the activation energy of dissolution for a wide selection range of silicate glasses and crystals.

**KEYWORDS:** Dissolution, Borosilicate glasses, Topological constraint theory, Vertical scanning Interferometry, Molecular Dynamics.