



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Incidence of Urinary Tract Infection after surgery for fractured hip

Rønfeldt, Ingerlise; Larsen, Lis Kjær; Pedersen, Preben Ulrich

*Publication date:*  
2019

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Rønfeldt, I., Larsen, L. K., & Pedersen, P. U. (2019). *Incidence of Urinary Tract Infection after surgery for fractured hip*. Poster presented at ICON 2019, Kolding , Denmark.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Incidence of urinary tract infection after surgery for fractured hip

Ingerlise Rønfeldt, RN, Cand. Scient KVT, Aalborg University Hospital, Orthopedic Division, Clinic Farsoe, Denmark.  
Lis Kjær Larsen, RN, MKS, Aalborg University Hospital, Orthopedic Division, Clinic Hjoerring, Denmark.  
Preben Ulrich Pedersen, RN, Professor, Ph.D., Centre of Clinical Guidelines – Clearinghouse, University of Aalborg, Denmark.  
Clinical Nursing research Unit, Aalborg University Hospital, Denmark

## Introduction

To prevent urinary tract infections (UTI) The National Clinical Guidelines for patients with Hip Fracture - recommends e.g. early and systematic mobilization and no use of indwelling catheters (1). Since 2003, these recommendations have existed and been implemented in the ward. Besides serious individual implications, hip fracture is also associated with substantial health costs (2).

In order to achieve the best possible result for the hip patient, two areas of nursing interventions seem particularly important. UTI is one of the most frequent complications related to hip fracture (3) and early mobilization accelerates functional refunds and contributes to a shortening of hospitalization (4).

## Aim

To describe the number of patients mobilized within 24 hours after surgery and number of patients developing UTI during admission.

## Methods

There were used a descriptive prospective design. This study was conducted from October 2015 to december 2016 at two departments of orthopedic surgery in Farsoe and Hjoerring, Aalborg University Hospital in Denmark.

The criteria for inclusions were adult patients who had surgery for fractured hip. Exclusions criteria were patients discharged to other departments than orthopedics, diagnosed dementia, not being able to talk and understand the Danish language, patients with catheter a demure, usually catheterization, or patients in antibiotic treatment on admission.

In this study all patients had urine samples collected on admission and at discharge, using sterile intermittent catheterisation. The urine sample should be collected within eight hours after admission to the emergency room. The definition of Centers for Disease Control (5) modified for Danish conditions ( $10^4$ ) were used to define urinary tract infection (1). Both urine samples were send to microbiological analysis, to diagnose the presence of UTI. Demographic data: age, gender, type of fracture, comorbidity, form of anaesthesia and waiting time for surgery. Care process data: time for early mobilization after surgery and which and for how long urinary catheter were used. Data are processed in the statistical program SPSS, version 23.

## Discussion

In our study, 29.2 % had a positive urine culture on admission. Other studies showed that 8- 52 % of patients had UTI during admission (7-9). It is therefore within the normal area of patients with UTI.

In this study, none of the patients with an indwelling catheter, was found to affect the incidence of UTI.

Furthermore, the study shows that the nurses had followed the National Clinical Guidelines, prevention UTI. There seem not to be problems in relation to adhere to the quality of hygiene by applying intermittent catheterization or catheter a demure.

The National Clinical Guideline emphasizes the importance of early mobilization, for patients with a fractured hip (1). The goal for mobilization is 90 %, within 24 hours postoperatively. The mobilization rate in this study were 52.3 %. This is lower than expected, when the National Clinical Guideline were implemented several years ago. Moreover, it was expected that the nurses knew the national Clinical Guideline.

## Results

The findings showed that 29.2% of patients with a fractures hip had a positive urine culture on admission to hospital. 6.2% contracted nosocomial urinary tract infection during admission. None of these patients had catheter a demure at hospitalization. All the patients with a diagnosed UTI received antibiotics for the infection, during their hospital stay. At discharge, 20% of the patients had a positive urine sample, but no symptoms. 13 patients were given a catheter a demure under hospitalization. 4 patients had Catheter a Demure beneath 24 hours (mean 14.75 hours). 9 patients had Catheter a Demure in more than 24 hours (mean 93.11 hours). None of those had urinary tract infection. The chi- square test used, testing whether patients with UTI and indwelling catheter, more often had an UTI than those, who not have been catheterized. Significance level was 0.898 and thus not significant.

Urine cultivation was performed on 96.9% of patients after hospitalization within 8 hours of arrival at the emergency room. 83 % got it examined at discharge. Four patients did not wanted to use intermittent catheterization upon discharge, because of

no symptoms and therefore didn't want the intervention. Five patients were treated for UVI during hospitalization.

Patient with UTI 9.2 % were intermitted catheterized. These patients were catheterized from 0- 23 times, except the two times, which were part of the project.

55.5 % of patients were not intermitted catheterized. Out of the 27 patients who were intermittent catheterized more than once, 5.4 % received a nosocomial urinary tract infection.

At first mobilization, nurse noted data and time. Mobilized patients within 24 hours postoperatively, were 52.3% of the patients. One patient didn't want mobilization. The rest of not mobilized patients, reason was not possible finding

Patients mobilized within 24 hours and who received an UTI, were 36.9 %.

Table 1. Descriptive statistics	N= 65
Female	87.7 %
Male	12.3 %
Age (mean, SD)	80.82 (SD 9.1)
(range)	55- 96
<b>Anesthetic:</b>	
General anesthesia	43.1 %
Spinal anesthesia	55.4 %
<b>Comorbidities:</b>	
Malignancy	6.2 %
Moderate to severe Chronic Kidney disease (CKD)	6.2 %
Congestive heart failure (CHF)	49.2 %
Chronic Obstructive Pulmonary Disease (COPD)	13.8 %
Cerebrovascular accident	7.7 %
Peptic ulcer disease	1.5 %
Connective tissue disease	12.3 %
Peripheral vascular disease	4.6 %
Diabetes mellitus	4.6 %
Other diseases	35.4 %
<b>Fracture type:</b>	
Colli femoris	44.6 %
Pertrocanteric	43.1 %
Interchrocanteric	3.1 %
Subtrocanteric	9.2 %
Waiting time for surgery, hours (mean, SD)	20.14 (SD 13.7)
(range hours)	0 - 72

Table 2 Percentage of results	N= 65
Urine culture at admission	96.9 %
Urine culture at discharge	83.0 %
<b>Number of patients with:</b>	
UTI on admission	29.2 %
UTI during admission	6.2 %
UTI at discharge	20.0 %
Patients with UTI, where sterile intermittent catheterization were used	9.2 %
Mobilization < 24 hours postoperatively	52.3 %
Patients mobilized < 24 hours postoperatively, also having UTI	36.9 %
Patients with Catheter a Demure	18.5 %
Mean (hours)	69.07
(range hours)	9 - 168

## Conclusions

The two departments who took part in this study do not have a higher number of UTI than other departments. Furthermore, there is a difference in the result because the sample results are number of bacteria, which is  $10^4$ , whereas the results compared to, are  $10^5$ .

In this study, all patients who had UTI were mobilized within 24 hours. Even if evidences indicate that mobilization can help prevent UTI, it does not appears as a valid factor in this study. However, it should be mentioned that this study was carried out on a basis of 65 patients, of which only 24 had UTI.

## Referencer

1. DrHOFTBRUD Dansk Tvaerfagligt Register for Hofteenaere Lærbensbrud. 2015;(december 2015). Available from: [https://www.sundhed.dk/content/cms/62/4662\\_hofte-fraktur-årsrapport\\_2017.pdf](https://www.sundhed.dk/content/cms/62/4662_hofte-fraktur-årsrapport_2017.pdf)
2. Hussey PS, Huckfeldt P, Hirshman S, Mehrotra A. Hospital and regional variation in medicare payment for inpatient episodes of care. JAMA Intern Med. 2015;175(6):1056-7.
3. Rose S, Maffuli N. Hip fractures. An epidemiological review.pdf. 1999;197-201, Volume 58, number 4.
4. Kristensen PK, Thillemann TM, Soballe K, Johnsen SP H. Evidensbaserede retningslinjer redder liv. 2017;(3):38-48.
5. Tambyah PA, Oon J. Catheter-Associated Urinary Tract Infection. Curr Opin Infect Dis. 2012;25(4):365-70.
6. Merchant RA, Lui KL, Ismail NH, Wong HP, Sitoh YY. The relationship between postoperative complications and outcomes after hip fracture surgery. Ann Acad Med Singapore. 2005;34(2):163-8.
7. Garcia-Alvarez F, Al-Ghanem R, Garcia-Alvarez I, López-Baïsson A, Bernal M. Risk factors for postoperative infections in patients with hip fracture treated by means of Thompson arthroplasty. Arch Gerontol Geriatr. 2010;50(1):51-5.
8. Johansson I, Athlin E, Frykholm L, Bolinder H, Larsson G. Intermittent versus indwelling catheters for older patients with hip fractures. J Clin Nurs [Internet]. 2002;11(5):651-656 6p. Available from: <http://search.ebscohost.com/login.aspx?direct=true&db=cin20&AN=106986051&lang=es&site=ehost-live&scope=site>