

## Computational screening of electrochemical properties of biological quinones for use in RFB technology

Kristensen, Sebastian Birkedal; van Mourik, Tanja; Pedersen, Tobias Bruun; Sørensen, Jens Laurids; Muff, Jens

*Publication date:*  
2019

*Document Version*  
Other version

[Link to publication from Aalborg University](#)

### *Citation for published version (APA):*

Kristensen, S. B., van Mourik, T., Pedersen, T. B., Sørensen, J. L., & Muff, J. (2019). *Computational screening of electrochemical properties of biological quinones for use in RFB technology*. Poster presented at The International Flow Battery Forum, Lyon, France.

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# Computational screening of electrochemical properties of biological quinones for use in RFB technology



Sebastian Birkedal Kristensen<sup>1</sup>, Tanja van Mourik<sup>2</sup>, Tobias Bruun Pedersen<sup>1</sup>, Jens Laurids Sørensen<sup>1</sup>, Jens Muff<sup>1</sup>

<sup>1</sup>Aalborg university, Department of Chemistry and Bioscience, Section of Chemical Engineering, Denmark, sbk@bio.aau.dk.


<sup>2</sup>University of St. Andrews, School of Chemistry, North Haugh, St. Andrews, Fife KY16 9ST, Scotland (UK).

1


## BACKGROUND - Bioquinones as electrontransfer agents in redox flow batteries

Recent quinones has been investigated as organic electrolytes or use in redox flow battery technology[1-2]. none has however so far investigated other candidates than synthetical quinones. Quinoes are produced by several sources in the nature, where the compound act as electron transfer agent, antimicrobial agent and oxidant/antioxidant[3-4].

- The aim of this study is to screen and thereby reveal potential quinones produced by biological sources as the best candidates to use in redox flow battery technology.



2




## COMPUTATIONAL WORKFLOW

A Database with 990 different quinone structures with origin from various natural sources was compiled. The calibration model was prepared according to previous published[5]. The calibration model was utilized to calculate the reduction potentials based on the simulated energies  $\Delta H_f$ , using the PBE/6-31G\*\* functional and basis-set.

The solvation energies,  $\Delta G_{solv}$  was also calculated, using an implicit solvation model.


3




## RESULTS - Distribution of bioquinones

The distribution of all the predicted redox potentials ( $E^0$ ) of the obtained biological quinones is visualized. The predicted values stretch from the most negative value at -1.382V vs SHE to the most positive value at 1.666V vs SHE.

We divided the dataset into three groups: bacteria(green), fungi(red) and others(orange) (plants, algae, animals etc.). The fungal quinones are more widely distributed compared to the other two groups. Thus, the fungi-produced compounds constitute the most varied group.




4



## RESULTS - Investigation of solubility of bioquinones

The distribution of the solubility compared to the redox potentials, with the BQs(red), NQs(green), AQs(purple) and  $\geq 4$  ring structures(green) highlighted respectively. The BQs tend to be at the positive side of the distribution, whereas the NQs are more centrally located. The AQs are more distributed towards the negative side compared to the BQs and NQs, which indicates a higher solubility of these compounds



5

## DISCUSSION AND CONCLUSION - Top 7 bioquinones of both catholytes and anolytes

The quinones with the lowest and highest calculated redox potentials are listed. The quinones with the most positive potentials predominantly originated from fungi and are relatively small and simple compounds, whereas the quinones with the most negative potentials are larger and more complex quinone structures. The compound with the most positive redox potential was phenicin with 1.666V vs SHE, a molecule produced by several Penicillium species. The compound with the lowest redox potential is Pradimicin M with -1.382V vs SHE and is produced in a mutant of Actinomadura hibisca Bacteria.

Name	$E^0$ V vs SHE	$\Delta G_{solv}$ kJ/mol	Source	Structure	Name	$E^0$ V vs SHE	$\Delta G_{solv}$ kJ/mol	Source	Structure
Pradimicin M	-1.382	-74.94	[B] blocked mutant of Actinomadura hibisca p157-2		Phenicin	1.666	-31.74	[F] Penicillium spp.	
Stemphytoxin I	-1.298	-40.15	[F] Stemphylium botrysosum var. lactucum		Tridentoquinone	1.485	-38.10	[F] Suillus tridentinus (Boletales)	
Granaticin B	-1.144	-72.42	[B] Streptomyces spiroverticillatus N-9940 (FERM-p 2330)		Stemphylierylenol	1.338	-75.20	[F] Stemphylium botrysosum var. lactucum	
Altartoxin I	-1.094	-37.34	[F] Alternaria spp.		Citrinin hydrate	1.170	-60.82	[F] Penicillium spp. Aspergillus spp.	
Altartoxin II	-1.008	-41.94	[F] Stemphylium botrysosum var. lactucum. Alternaria spp.		Fusarnaphthoquinone A	1.135	-32.95	[F] Fusarium spp. PSU-F34 and PSU-F135	
3-O-Methylhydrofusarubin A	-1.001	-33.43	[F] Fusarium martii		5-Hydroxydihydrofusarubin B	1.126	-52.50	[F] endophytic Fusarium sp. BCC14842	
Stemphytoxin IV	-0.976	-36.40	[F] Stemphylium botrysosum var. lactucum		5-Hydroxy-3-methoxydihydrofusarubin D	1.028	-51.20	[F] endophytic Fusarium sp. BCC14842	

1. B. Huskinson et al., "A metal-free organic-inorganic aqueous flow battery," Nature, vol. 505, no. 7482, pp. 195–198, 2014.

2. K. Lin et al., "Alkaline quinone flow battery," Science (80-. ), vol. 349, no. 6255, pp. 1529–1532, 2015.

3. Thomson., R.H. (1987). Naturally occurring quinones III: recent advances (3rd ed.). London; New York: Chapman and Hall, 1987.

4. Medentsev, A. (1996). Fungal Naphthoquinone metabolites (review). Applied Microbiology and Biotechnology, 32(1), 7–29.

5. S. Er, C. Suh, M. P. Marshak, and A. Aspuru-Guzik, "Computational design of molecules for an all-quinone redox flow battery," Chem. Sci., vol. 6, no. 2, pp. 885–893, 2015.

