

Levels of Intestinal Inflammation and Fibrosis in Resection Specimens after Preoperative Anti-Tumor Necrosis Factor Alpha Treatment in Patients with Crohn's Disease

A Comparative Pilot Study

Torle, J; Dabir, P D; Korsgaard, U; Christiansen, J; Qvist, N; El-Hussuna, A

Published in:
Surgery research and practice

DOI (link to publication from Publisher):
[10.1155/2020/6085678](https://doi.org/10.1155/2020/6085678)

Creative Commons License
CC BY 4.0

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Torle, J., Dabir, P. D., Korsgaard, U., Christiansen, J., Qvist, N., & El-Hussuna, A. (2020). Levels of Intestinal Inflammation and Fibrosis in Resection Specimens after Preoperative Anti-Tumor Necrosis Factor Alpha Treatment in Patients with Crohn's Disease: A Comparative Pilot Study. *Surgery research and practice*, 2020, Article 6085678. <https://doi.org/10.1155/2020/6085678>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: July 04, 2025

Research Article

Levels of Intestinal Inflammation and Fibrosis in Resection Specimens after Preoperative Anti-Tumor Necrosis Factor Alpha Treatment in Patients with Crohn's Disease: A Comparative Pilot Study

J. Torle,¹ P. D. Dabir,² U. Korsgaard,² J. Christiansen,² N. Qvist,³ and A. El-Hussuna ⁴

¹Department of Surgery, Regional Hospital Randers, Midt-Jylland region, Randers, Denmark

²Department of Pathology, Regional Hospital Randers, Midt-Jylland region, Randers, Denmark

³IBD Care, Surgical Research Unit, Odense University Hospital, Southern University of Denmark, region Syd Danmark, Randers, Denmark

⁴Department of Surgery, Aalborg University Hospital, Aalborg, Nord-Jylland region, Denmark

Correspondence should be addressed to A. El-Hussuna; alaanewemail@gmail.com

Received 10 July 2019; Revised 13 January 2020; Accepted 4 February 2020; Published 21 February 2020

Academic Editor: Todd Pesavento

Copyright © 2020 J. Torle et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Strictures are a common complication in Crohn's disease (CD), found in more than 50% of patients. They are characterized by the excessive deposition of extracellular proteins in the tissue as a result of the chronic inflammatory process. The effect of anti-tumor necrosis factor alpha (TNF- α) therapy on the development of fibrosis is not yet fully understood. **Aim.** To investigate whether the degree of intestinal inflammation and fibrosis is correlated with preoperative anti-TNF- α therapy in patients with CD who are undergoing bowel resection. **Methods.** This unblinded, prospective, single tertiary center, pilot cohort study included all adult patients with CD who underwent elective, laparoscopic, or open intestinal resection. Preoperative investigations included measurement of blood TNF- α concentration, specific antidrug antibodies, and the concentration of selected inflammatory cytokines. Three pathologists independently examined the specimens and assessed the degree of inflammation and fibrosis. **Results.** Histopathological specimens from 10 patients with CD who underwent ileocecal or ileocolic resections were retrieved. Four of those patients were on anti-TNF- α treatment prior to surgery. The last dose of the anti-TNF- α agent was administered 1–9 weeks prior to bowel resection. Patients on anti-TNF- α treatment had a higher fibrosis score than controls ($p = 0.01$). Anti-TNF- α treatment was not associated with an increase in CD68- or CD163-positive macrophages. There was no significant relationship between the time from the final preoperative anti-TNF- α dose to surgery and the fibrosis score. No significant association was found between the concentration of major inflammatory cytokines, including TNF- α , and the fibrosis score or degree of inflammation. **Conclusions.** Patients who underwent preoperative anti-TNF- α treatment had a higher fibrosis score than controls.

1. Introduction

More than 50% of patients with Crohn's disease (CD) will develop intestinal fibrosis (stenosis and strictures) as a result of chronic inflammation [1].

Persistent and recurring tissue injury from inflammation, which is triggered and sustained by proinflammatory cytokines, leads to a continuous cycle of tissue breakdown

and repair. This results in the accumulation of fibroblasts and myofibroblasts, which may lead to fibrosis as a result of extracellular matrix production [2].

Clinically, strictures may occur as a result of inflammation-mediated swelling of the intestinal wall, fibrotic occlusions, or a combination of both [3]. It is well known that immunosuppressive drugs, including TNF- α inhibitors, may reduce the occurrence of inflammatory strictures; however,

their effect on the fibrotic reaction is not clear [4]. As chronic inflammation is closely associated with fibrosis and stricture formation, and considering the key role of TNF- α in initiating the inflammatory response [5], we hypothesized that anti-TNF- α therapy may lead to reduced fibrosis.

Intestinal macrophages have many important functions in maintaining gut homeostasis, but they also play a role in the development of inflammation. When identifying macrophages in tissue by immunohistochemistry, two widely used macrophage markers are CD68 and CD163 [6].

The aim of the present study was to investigate whether the degree of intestinal fibrosis, inflammation, and the number of macrophages are correlated with preoperative anti-TNF- α therapy in patients with CD undergoing bowel resection.

2. Materials and Methods

2.1. Study Design. This is an unblinded, prospective, non-interventional, pilot cohort study.

2.2. Study Hypothesis. Patients with CD who receive preoperative treatment with anti-TNF- α drugs will have less inflammation and a lower degree of intestinal fibrosis, assessed by CD163 concentration in the peripheral blood and histology, when compared to anti-TNF- α naïve patients.

2.3. Outcome Measures. The primary outcome measure was the degree of fibrosis. The secondary outcome was the grade of inflammation.

2.4. Inclusion Criteria. All adult patients with CD who were scheduled for elective small bowel or colon resection (open or laparoscopic) from a single tertiary center were included.

2.5. Exclusion Criteria. All patients with sepsis (abscess or fistula) or acute intestinal obstruction were excluded.

2.6. Extracted Data. This study is a continuation of the Inflammatory Bowel Disease Response study [7], in which resection specimens were obtained from patients undergoing elective ileocecal or ileocolic resections. Surgery and acquisition of the resection specimens were carried out as described previously [7].

Patient demographics, disease severity, nutritional status, comorbidities, preoperative medications, previous operations for CD, presence of preoperative intra-abdominal abscess or enteric fistula, preoperative nutritional support, disease duration, and disease localization were registered after obtaining oral and written informed consent. Blood samples were collected preoperatively and at 6, 24, and 48 hours after the surgical incision. Laboratory workup included serum analyses for TNF- α , interleukin- (IL-) 6, IL-8, IL-10, and IL-17A, in addition to white blood cell count, total iron-binding capacity, and C-reactive protein, albumin, hemoglobin, D-dimer, and cortisol levels. The drug concentration in the serum was measured in addition to the antidrug antibodies titer.

Out of the 46 patients included in the IBD Response study, bowel specimens from 10 patients with CD met the inclusion criteria and were eligible for analysis. Four of the 10 included patients had received TNF- α treatment preoperatively, with the last dose administered 1–9 weeks prior to surgical intervention. Representative hematoxylin and eosin (H&E)-stained slides from each of the resection specimens were chosen for further analysis. Fibrosis was assessed by Masson trichrome (MT) staining. Examples of fibrosis stained blue in Masson Trichrome special stain are shown in Figure 1.

To assess the number of macrophages, the immunohistochemical markers CD68 and CD163 were used. The slides were digitalized using a Hamamatsu slide scanner™ and image processing was subsequently performed in Visiopharm™. For both markers, deparaffinized sections were stained in a Ventana BenchMark ULTRA XT automatic stainer using 3,3'-diaminobenzidine (DAB) (OptiView universal DAB IHC detection kit; Ventana BioTek System, Tucson, AZ, USA), performed according to the manufacturer's instructions. Briefly, sections were demasked with cell conditioning buffer (CC1, Ventana Medical Systems, Tucson, AZ, USA) for 8 min followed by 16 min, and endogenous peroxidase activity was blocked with the UltraView inhibitor. Primary antibodies against CD68 (PREP KIT 54) and CD163 (CD163 [MRQ-26]) were applied for 32 min at 36°C. Sections were then incubated with OptiView horseradish peroxidase (HRP) multimer (Ventana BioTek Systems), containing a mixture of HRP-labeled goat anti-mouse and goat anti-rabbit antibodies. Bound antibodies were visualized by incubation with OptiView hydrogen peroxide substrate and DAB chromogen, and counterstained with hematoxylin. Positive signals for both CD68 and CD163 markers were evidenced by a brown color reaction in the cell cytoplasm.

Slides were evaluated by two consultant pathologists (PDD and JJC) and one trainee pathologist (UK) to minimize interobserver variation. The pathologists were blinded to the patients' clinical data and diagnosis. The three independent observers graded the degree of fibrosis on the MT-stained slides on a scale of 1–4. The grading was performed semi-quantitatively by assessing the amount of blue-stained fibers in the representative areas.

Acute and chronic inflammation was assessed in all four layers of the intestinal wall, and graded as absent, mild, moderate, or severe. The inflammation grades were determined semi-quantitatively by eyeballing, with a triple head microscope. Disagreements were resolved by consensus, and grades were assigned a numerical value from 0–3 for statistical analysis. The grades were defined as follows: absent (0), no traces of inflammation; mild (1), few inflammatory cells (mild acute inflammation in mucosa was defined as neutrophils in the lamina propria); moderate (2), moderate amount of inflammatory cells (moderate acute inflammation in mucosa was defined as the presence of cryptitis); and severe (3), many inflammatory cells (severe acute inflammation in mucosa was defined as the presence of crypt abscess).

Areas of ulceration were excluded from the analysis.

The immunohistochemical slides were assessed independent to the HE-stained sections in a blinded manner.




FIGURE 1: Example of fibrosis stained blue in Masson Trichrome special stain.

Positive immunohistochemical staining was defined as cytoplasmic staining for CD68 and CD163. To assess the number of CD68- and CD163-positive cells, the areas of mucosa, submucosa, and muscularis propria deemed to contain the most CD163-positive cells under $4\times$ magnification were selected. The CD68 and CD163 slides were then aligned, and each observer counted the number of positive CD68 and CD163 cells in the selected area ($110,000\ \mu\text{m}^2$) at $20\times$ magnification. A representative image of CD68 staining is presented in Figure 2.

2.7. Statistical Analysis. For the fibrosis analysis, the mean value of each independent observers' scores were calculated and used as the final result. Student's *t*-test was applied for all observations. The kappa score for inter-observer agreement was calculated using Fleiss' kappa for more than two raters. For the univariate analysis, Pearson's chi-square and Fisher's exact tests were employed. Continuous data were compared using the Mann-Whitney *U* test and one-way ANOVA. The relationship between continuous variables was analyzed using Pearson's chi-square correlation. Due to the small sample size of this pilot study, multivariate logistic regression was not used. SPSS version 19 software (IBM SPSS Statistics for Windows, Version 22.0; Armonk, NY: IBM Corp.2010) was used for all statistical analyses.

3. Results

3.1. Study Population Characteristics. Patients' characteristics are shown in Table 1. The mean age was $40.10 (\pm 12.97)$ years. Five patients were female. None of the patients had comorbidities.

3.2. Drug Levels at the Time of Operation. Four patients who underwent preoperative treatment with anti-TNF- α received their last dose 1–9 weeks prior to bowel resection. Three of the four patients who were on anti-TNF- α treatment had no detectable drug concentration at the time of operation. One patient was positive for antidrug antibodies at the time of operation, although this patient had an undetectable drug concentration.

3.3. Fibrosis. The patients' fibrosis scores are presented in Table 2. Patients on anti-TNF- α treatment had a higher fibrosis score compared to controls ($p=0.01$). There were no significant relationships between the number of weeks of administration of anti-TNF- α agents and the fibrosis score, CD68 or CD163 levels. The fibrosis score was not correlated with CD68 or CD163 levels. Preoperative and 6 h postoperative TNF- α concentrations were not correlated with the fibrosis score. The disease duration was not correlated with the degree of fibrosis.

3.4. Grade of Inflammation. Grades of acute and chronic inflammation are presented in Table 2. Patients on anti-TNF- α treatment showed no difference in inflammation compared to the controls. Neither the duration of anti-TNF- α treatment nor the anti-TNF- α drug concentration was associated with the grade of inflammation.

3.5. Macrophage Markers CD68 and CD163. The presence of CD68 and CD163 in the submucosa of patients with CD were significantly correlated ($p=0.014$), but not in the mucosa or muscularis propria. The concentrations of inflammatory markers (TNF- α , IL-6, IL-8, IL-10, IL-17, and cortisol) showed no significant correlations with CD68 or CD163, whether in the mucosa, submucosa or muscularis propria. Anti-TNF- α treatment was not associated with an increase in CD68 or CD163. No significant correlations were observed between preoperative C-reactive protein levels and fibrosis score, CD68 or CD163.

4. Discussion

There was no significant relationship between the duration of anti-TNF- α administration and the fibrosis score. This might be explained by differences in disease duration, the development of fibrosis prior to medication, or genotypic and phenotypic differences between patients. An allele that could lead to overexpression of the main profibrotic mediator, TGF- β , has been identified [8], which could lead to more rapid development of fibrosis and strictures. Another explanation could be that inflammation and fibrosis are initially linked in the pathogenesis, but later become two independent processes [9]. This might explain why fibrosis cannot be reversed by anti-inflammatory drugs.

The duration of anti-TNF- α treatment was not significantly correlated to either of the macrophage-associated receptors (CD68 and CD163). CD163 serves as a marker for activated macrophages in its soluble form [10]. Although the CD163 marker has been linked to the anti-inflammatory M2 macrophage phenotype, CD163-positive macrophages have been found to outnumber CD68-positive cells in a range of inflammatory events, including those that are associated with CD [6]. Dige et al. found that administration of anti-TNF- α led to a significant reduction in soluble CD163 when compared to biologic-naïve controls [10]. In the present study, we did not find such a correlation. This may be ascribed to the fact that plasma concentrations, rather than




FIGURE 2: Example of cell cytoplasmic staining of macrophages using the CD68 immunohistochemical marker.

tissue concentrations, were measured in the Dige et al. study [10]. In addition, none of the patients had detectable drug levels at the time of operation, which might also explain why CD163 concentrations were not attenuated.

In order to measure the number of macrophages, two cellular markers, CD68 and CD163, were chosen. CD68 is a pan-macrophage marker [11], whereas CD163 is a macrophage-specific scavenger receptor with anti-inflammatory abilities in M2 macrophages. When macrophages are activated, CD163 proteins are shed into the bloodstream [10]. By measuring soluble CD163, one can monitor the number of activated macrophages in the host. Consequently, elevated levels of CD163 are associated with conditions with a heightened inflammatory response, such as CD [10]. Using this method, however, no correlation between fibrosis and the number of macrophages was found. Additionally, no correlations were found between various cytokines (TNF- α , IL6 IL8, IL10, and IL17) and cortisol and CD68 and CD163. If this holds true in a larger population, it could indicate that macrophages play a more active role in inflammation than in the development of fibrosis. Alternatively, the question must be raised as to the accuracy of these markers for measuring the degree of fibrosis.

The lack of association between treatment with the anti-TNF- α drug and a decrease in inflammation is remarkable, as its main effect is to inhibit the proinflammatory cytokine TNF- α . This could indicate that the effect of the drug is subeffective or ineffective in the target tissue. Previous studies have concluded that inflammatory processes may

be confined to the fibrostenotic tissue, which may be beyond the reach of anti-TNF- α drugs [12]. Also, of the patients in the treatment group, half had received their last dose 9 weeks prior to surgery, which might be too long to detect a decrease in inflammation. If this is the case, it may explain why we observed no reductions in fibrosis or in macrophage activity.

The increase in fibrosis found in this study may seem counterintuitive considering its effectiveness against inflammation. This has been reported in other studies, and the general explanation is that TNF- α counterbalances the effects of TGF- β . Inhibition of TNF- α might therefore promote the profibrotic signaling cascade [13]. Although previous studies have found a significant decrease in inflammation in the treatment group, there was a significant increase in pauci-inflammatory fibrosis in the muscularis mucosae, as well as a trend toward increased fibrosis in the muscularis propria, albeit not statistically significant. This was ascribed to the classic “lattice-like” healing pattern also seen in endoscopy after anti-TNF- α treatment [13]. Since its introduction, theories related to the role of anti-TNF- α in the development of fibrosis have been subject to debate. Although several studies conducted on large populations have failed to produce evidence that support such a claim, concerns regarding the drug’s potential in causing fibrosis and strictures in these patients still remain [13].

We suspect that the increased fibrosis score observed in patients with CD on anti-TNF- α treatment in the current study indicates a nonresponse to the treatment because a

TABLE 1: Preoperative and intraoperative characteristics of 10 patients with Crohn's disease treated with anti-TNF- α compared to anti-TNF- α naïve patients.

Patients' characteristics	Anti-TNF- α treatment 4/10 patients (40%)	Anti-TNF- α naïve 6/10 patients (60%)	Univariate analysis
Age (years; mean \pm SD)	27.4 \pm 10.25	48.5 \pm 5.28	$p = 0.003$
Female	3/4 (75%)	2/6 (33.3%)	Ns
Body mass index (kg/m ² ; mean \pm SD)	26.58 \pm 11.8	22.5 \pm 2.17	Ns
Smoking, n (%):			Ns
Non- or ex-smoker	3/4 (75%)	6/6 (100%)	
Smoker	1/4 (25%)	0/6 (0%)	
Steroids, n (%)	0	3/6 (50%)	Ns
Immunomodulators, n (%)	1/4 (25%)	2/6 (33.3%)	Ns
Disease localization (Montreal classification)			Ns
L2	1/4 (25%)		
L3	3/4 (75%)		
Harvey-Bradshaw Index > median (7.5), n (%)	1/4 (25%)	3/6 (50%)	Ns
Disease duration (years; mean \pm SD)	5 \pm 3.16	14.33 \pm 9.05	Ns
Disease phenotype (Montreal classification)			Ns
B2 (stricturing)	4/4	3/6 (50%)	
B3 (penetrating)	0	3/6 (50%)	
Previous laparotomy/laparoscopy for CD	1/4 (25%)	3/6 (50%)	Ns
Preoperative parenteral nutrition, n (%)	0	1/6 (16.7%)	Ns
Preoperative sepsis	0	2/6 (33.3%)	Ns
Preoperative CRP mg/L (mean \pm SD)	32.18 \pm 37.5	19.35 \pm 19.28	Ns
Preoperative TNF- α ng/L (mean \pm SD)	1.22 \pm 2.14	0.37 \pm 0.35	Ns
Access to abdomen, n (%)			$p = 0.005$
Laparoscopic	4/4	0	
Open	0	6/6	
Type or resection, n (%)			Ns
Small bowel resection	0	2/6 (33.3%)	
Ileocolic resection	4/4*	4/6 (66.7%)*	

All operations were performed with a specialist surgeon in charge. Anti-TNF- α , anti-tumor necrosis factor alpha drugs; Ns, nonsignificant; CRP, C-reactive protein; SD, standard deviation. *In each group that underwent ileocolic resection, one patient also received colectomy.

TABLE 2: Drug administration, fibrosis score, and inflammation grade of patients with Crohn's disease.

Birth year	Gender (m/f)	Anti-TNF- α dose (mg)	TPS (weeks)	Fibrosis score			Acute inflammation grade				Chronic inflammation grade			
				P1	P2	P3	Mu	SM	MuP	SS	Mu	SM	MuP	SS
1997	f	40	1	4	3	4	2	1	0	0	3	3	1	2
1993	m	100	9	3	3	3	1	0	1	0	1	2	1	1
1974	m	0	—	2	2	3	2	1	0	0	2	2	2	1
				1	1	1	3	0	1	0	1	3	2	1
1963	m	0	—	1	1	2	1	0	0	0	1	1	1	0
1965	f	0	—	2	3	2	2	0	0	1	1	2	0	1
				2	3	4	1	0	0	1	2	3	2	2
1962	m	0	—	2	1	2	3	0	1	0	2	3	1	2
1982	f	40	1	3	3	4	1	0	0	1	2	2	1	1
1962	f	0	—	2	3	3	3	1	1	0	2	3	2	3
1972	m	0	—	1	2	1	3	1	0	1	1	1	0	1
				4	4	4	2	0	1		2	0	1	
1975	f	100	9	3	3	4	1	0	0	0	0	0	0	1

TPS, time prior to surgery; P1–3, pathologists 1–3; Mu, mucosa; SM, submucosa; MuP, muscularis propria; SS, subserosa.

response would have limited inflammation and fibrosis. This hypothesis is to be investigated in a large prospective cohort study.

5. Limitations

This study was a pilot study that can serve as a template for future larger-scale studies. The two groups were not

comparable with regard to age, gender, disease duration, preoperative events, and surgical approach. The patients in the control group seemed to be more sick (older, longer disease duration, and open surgery), which would translate to more inflammation and fibrosis. The study population was a group of patients in whom the medical treatment was not sufficient to avoid operation, and nonresponders to anti-TNF- α treatment may differ to responders with regard to the

formation of fibrosis. Logistic regression analysis to adjust for confounders was considered, but due to the small sample size, and after consulting a biomedical statistician, this was deemed inappropriate.

6. Conclusion

Patients who received preoperative anti-TNF- α treatment had a higher fibrosis score compared to controls. A significant correlation between the duration of drug administration and fibrosis score could not be found. Furthermore, there was no correlation between the duration of drug administration and the number of CD68 and CD163 macrophage receptors. The degree of acute and chronic inflammation was not significantly different between the two groups. There is a need for a large prospective study.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Additional Points

Perspectives. Histopathological examination of fibrosis and inflammation might indicate whether the patient will respond to biological treatment.

Ethical Approval

The study was approved by the ethics committee in the capital region (reference number H-2-2013-166) and the region of Zealand (SJ-399), and the Danish Data Protection Agency (Datatilsynet) in the capital region (reference number HVH-2013-046/02515) and in the region of Zealand (reference number REG-85-2013). The study was registered in clinicaltrials.gov (identifier: NCT01974869) and Trials Map on the ESCP website (<http://www.escp.eu.com/research/international-trials/trials-map>).

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

JT, PDD, UK, JC, NQ, and AE contributed to study design, data acquisition, analysis, and interpretation, as well as drafting of the manuscript. AE and PDD supervised the project. All authors performed critical revision of the manuscript for important intellectual content.

Acknowledgments

The authors would like to thank the Institute of Pathology, Regional Hospital, Randers, Denmark, for their technical assistance in carrying out this project and the Department of Pathology, Odense University Hospital, Denmark, for contributing the required materials for this project. This

study was supported by the Else og Mogens Wedell Wedellsborgs fund (Denmark).

References

- [1] F. Rieder, "Managing intestinal fibrosis in patients with inflammatory bowel disease," *Gastroenterology and Hepatology*, vol. 14, no. 14, pp. 120–122, 2018.
- [2] F. Rieder and C. Fiocchi, "Mechanisms of tissue remodeling in inflammatory bowel disease," *Digestive Diseases*, vol. 31, no. 2, pp. 186–193, 2013.
- [3] D. Bettenworth and F. Rieder, "Medical therapy of stricturing Crohn's disease: what the gut can learn from other organs - a systematic review," *Fibrogenesis & Tissue Repair*, vol. 7, no. 1, p. 5, 2014.
- [4] A. M. Schoepfer, E. Safroneeva, S. R. Vavricka, L. Peyrin-Biroulet, and C. Mottet, "Treatment of fibrostenotic and fistulizing Crohn's disease," *Digestion*, vol. 86, no. s1, pp. 23–27, 2012.
- [5] M. Cottone and V. Criscuoli, "Infliximab to treat Crohn's disease: an update," *Clinical and Experimental Gastroenterology*, vol. 4, no. 4, pp. 227–238, 2011.
- [6] M. H. M. Barros, F. Hauck, J. H. Dreyer, B. Kempkes, and G. Niedobitek, "Macrophage polarisation: an immunohistochemical approach for identifying M1 and M2 macrophages," *PLoS One*, vol. 8, no. 11, Article ID e80908, 2013.
- [7] A. El-Hussuna, N. Qvist, M. S. Zangenberg et al., "No effect of anti-TNF- α agents on the surgical stress response in patients with inflammatory bowel disease undergoing bowel resections: a prospective multi-center pilot study," *BMC Surgery*, vol. 18, no. 1, p. 91, 2018.
- [8] M. C. Visschedijk, L. M. Spekhorst, S.-C. Cheng et al., "Genomic and expression analyses identify a disease-modifying variant for fibrostenotic Crohn's disease," *Journal of Crohn's and Colitis*, vol. 12, no. 5, pp. 582–588, 2018.
- [9] G. Rogler and M. Hausmann, "Factors promoting development of fibrosis in Crohn's disease," *Frontiers in Medicine*, vol. 4, p. 96, 2017.
- [10] A. Dige, S. Støy, K. L. Thomsen et al., "Soluble CD163, a specific macrophage activation marker, is decreased by anti-TNF- α Antibody treatment in active inflammatory bowel disease," *Scandinavian Journal of Immunology*, vol. 80, no. 6, pp. 417–423, 2014.
- [11] Y. Kim, X. Wen, J. M. Bae, J. H. Kim, N.-Y. Cho, and G. H. Kang, "The distribution of intratumoral macrophages correlates with molecular phenotypes and impacts prognosis in colorectal carcinoma," *Histopathology*, vol. 73, no. 4, pp. 663–671, 2018.
- [12] R. Hundal, X. Gui, R. Panaccione et al., "Mo1238 resected fibrostenotic intestine from Crohn's disease patients express significant TNF in proliferative fibroblasts and in extracellular matrix," *Gastroenterology*, vol. 146, no. 5, 594 pages, 2014.
- [13] D. F. Schaeffer, J. C. Walsh, R. Kirsch, M. Waterman, M. S. Silverberg, and R. H. Riddell, "Distinctive histopathologic phenotype in resection specimens from patients with Crohn's disease receiving anti-TNF- α therapy," *Human Pathology*, vol. 45, no. 9, pp. 1928–1935, 2014.