Aalborg Universitet



Management of maxillofacial trauma in the elderly

A European multicenter study

Brucoli, Matteo; Boffano, Paolo; Romeo, Irene; Corio, Chiara; Benech, Arnaldo; Ruslin, Muhammad; Forouzanfar, Tymour; Starch-Jensen, Thomas; Rodríguez-Santamarta, Tanía; de Vicente, Juan Carlos; Snäll, Johanna; Thorén, Hanna; Tarle, Marko; Dediol, Emil; Pechalova, Petia; Pavlov, Nikolai; Daskalov, Hristo; Doykova, Iva; Kelemith, Kadri; Tamme, Tiia; Kopchak, Andrey; Shumynskyi, Ievgen; Corre, Pierre; Bertin, Helios; Goguet, Quentin; Anquetil, Marine; Louvrier, Aurélien; Meyer, Christophe; Dovšak, Tadej; Vozlič, David; Birk, Anže; Aničić, Boban; Konstantinovic, Vitomir S

Published in: Dental Traumatology

DOI (link to publication from Publisher): 10.1111/edt.12536

Publication date: 2020

Document Version Accepted author manuscript, peer reviewed version

Link to publication from Aalborg University

Citation for published version (APA):

Brucoli, M., Boffano, P., Romeo, I., Corio, C., Benech, A., Ruslin, M., Forouzanfar, T., Starch-Jensen, T., Rodríguez-Santamarta, T., de Vicente, J. C., Snäll, J., Thorén, H., Tarle, M., Dediol, E., Pechalova, P., Pavlov, N., Daskalov, H., Doykova, I., Kelemith, K., ... Konstantinovic, V. S. (2020). Management of maxillofacial trauma in the elderly: A European multicenter study. *Dental Traumatology*, *36*(3), 241-246. https://doi.org/10.1111/edt.12536

General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal -

Take down policy If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: July 08, 2025

TITLE: Management of maxillofacial trauma in the elderly: a European multicenter study.

AUTHORS: Matteo Brucoli MD DDS,¹ Paolo Boffano MD, ¹ Irene Romeo MD, ¹ Chiara Corio MD, ¹ Arnaldo Benech MD DDS, ¹ Muhammad Ruslin MD DDS PhD,² Tymour Forouzanfar MD DDS PhD,³ Thomas Starch- Jensen MD PhD,⁴ Tanía Rodríguez-Santamarta MD DDS,⁵ Juan Carlos de Vicente MD DDS PhD, ⁵ Johanna Snäll MD DDS PhD,⁶ Hanna Thorén MD DDS PhD,^{7,8} Marko Tarle MD,⁹ Emil Dediol MD PhD,⁹ Petia Pechalova MD DDS PhD,¹⁰ Nikolai Pavlov MD DDS,¹¹ Hristo Daskalov MD DDS,¹⁰ Iva Doykova MD DDS,¹² Kadri Kelemith DDS,¹³ Tiia Tamme MD PhD,¹⁴ Andrey Kopchak MD DDS PhD,¹⁵ levgen Shumynskyi MD DDS,¹⁶ Pierre Corre MD PhD,¹⁷ Helios Bertin MD PhD,¹⁷ Quentin Goguet MD,¹⁷ Marine Anquetil MD,¹⁷ Aurélien Louvrier MD,^{18,19} Christophe Meyer MD PhD,^{18,19} Tadej Dovšak MD PhD,²⁰ David Vozlič MD,²⁰ Anže Birk MD,²⁰ Boban Aničić MD DDS,²¹ Vitomir S. Konstantinovic DDS MD MSc PhD.²¹

¹ Division of Maxillofacial Surgery, University Hospital "Maggiore della Carità", University of Eastern Piedmont, Novara, Italy

 ² Department of Oral and Maxillofacial Surgery, Hasanuddin University, Makassar, Indonesia
 ³ Department of Oral and Maxillofacial Surgery/Oral Pathology, VU University Medical Center, Amsterdam, The Netherlands

⁴ Department of Oral and Maxillofacial Surgery, Aalborg University Hospital, Aalborg, Denmark
 ⁵ Servicio de Cirugía Maxilofacial, Hospital Universitario Central de Asturias, Oviedo, Spain
 ⁶ Department of Oral and Maxillofacial Diseases, University of Helsinki and Helsinki University
 Hospital, Helsinki, Finland.

⁷ Department of Oral and Maxillofacial Surgery, Institute of Dentistry, University of Turku.

⁸ Department of Oral and Maxillofacial Diseases, Turku University Hospital, Turku, Finland

⁹ Department of Maxillofacial Surgery, University Hospital Dubrava, Zagreb, Croatia

¹⁰ Department of Oral surgery, Faculty of Dental Medicine, Medical University, Plovdiv, Bulgaria

¹¹ Private practice of oral surgery, Plovdiv, Bulgaria

¹² Department of maxillofacial surgery, Faculty of Dental Medicine, Medical University, Plovdiv,
 Bulgaria

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the <u>Version of Record</u>. Please cite this article as <u>doi:</u> 10.1111/EDT.12536

- ¹³ Department of maxillo-facial surgery, North Estonia Medical Centre Foundation, Tallinn, Estonia.
- ¹⁴ Faculty of Medicine, University of Tartu, Tartu, Estonia
- ¹⁵ Bogomolets National Medical University, Stomatological medical center, Kyiv, Ukraine.
- ¹⁶ Bogomolets National Medical University, Kyiv City Clinical Emergency Hospital, Kyiv, Ukraine
- ¹⁷ Division of Maxillofacial Surgery, Chu de Nantes, Nantes, France
- ¹⁸ Department of Oral and Maxillofacial Surgery Hospital Dentistry Unit, University Hospital of Besançon – France
- ¹⁹ University of Bourgogne Franche-Comté, EA 4662 Nanomedicine Lab Imagery and Therapeutics, F-25000 Besançon – France
- ²⁰ Department of *Maxillofacial* and Oral Surgery of the University Medical Centre, Ljubljana, Slovenia
- ²¹ Department of Maxillofacial surgery, School of Dental Medicine, University of Belgrade, Serbia

Address correspondence and reprint requests to Dr Paolo Boffano: Division of Maxillofacial Surgery, University Hospital "Maggiore della Carità", University of Eastern Piedmont, Novara, Italy

E-mail address: paolo.boffano@gmail.com

Disclosure: The authors have no financial interest to declare in relation to the content of this article.

No funding

DR PAOLO BOFFANO (Orcid ID : 0000-0002-0782-9933)

PROFESSOR MUHAMMAD RUSLIN (Orcid ID : 0000-0003-0943-4000)

Article type : Original Article

ABSTRACT:

BACKGROUND / AIMS: Management of maxillofacial trauma in the geriatric population poses a great challenge due to anatomical variations and medical comorbidities. The aim of this study was to analyze the management variables, timing, and outcomes of facial fractures in elderly patients (aged 70 years or more) at several European departments of oral and maxillofacial surgery.

MATERIALS AND METHODS: This study was based on a systematic computer-assisted database that allowed the recording of data from all geriatric patients with facial fractures from the involved maxillofacial surgical units across Europe between 2013 and 2017. **RESULTS**: A total of 1334 patients were included in the study: 665 patients underwent closed or open surgical treatment. A significant association (p < .005) was found between the presence of concomitant injuries and a prolonged time between hospital admission and treatment. The absence of indications to treatment was associated with comorbidities and an older age (p < .00005).

CONCLUSIONS: Elderly patients require specific attention and multidisciplinary collaboration in the diagnosis and sequencing of trauma treatment. A prudent attitude may be kept in selected cases, especially when severe comorbidities are associated and function is not impaired.

KEYWORDS: elderly; maxillofacial trauma; management; geriatric.

Article

INTRODUCTION:

Management of maxillofacial trauma in the geriatric population poses a great challenge due to anatomical variations and medical comorbidities.¹⁻⁹ Moreover, the frequency of old patients sustaining craniofacial trauma due to an increase in the ageing population is leading to a progressive increase of importance of this topic in clinical maxillofacial practice.¹⁰⁻²⁰ The principles of management of facial fractures in the elderly population and adults remain basically the same. However, the way they are really managed in the geriatric population may become noticeably different due to anatomic and physiologic alterations in this population.⁶⁻¹¹ Bone atrophy, inadequate blood supply, reduced capacity for tissue repair, declining baseline functions, and above all the prevalence of pre-existing diseases and comorbidities, may change not only timing of surgery but also the indications for surgery in elderly patients.⁶⁻⁸

The management of geriatric patients with facial fractures is often challenging, as they are more severely injured, hospital stay is prolonged, and deaths following trauma occur more frequently compared with younger adults.⁶⁻⁹

The peri-operative management of the acutely injured elderly patient is also more complex than that of younger patients, with a disproportionate consumption of health care resources.⁹⁻¹¹

Publications regarding knowledge and up-to-date management of geriatric facial fractures are still rare.⁶⁻¹¹

Therefore, several European centers that had already shown research experience in maxillofacial trauma decided to collaborate on a multicenter research project about maxillofacial fracture management in elderly patients.¹⁻⁵

The aim of this study was to analyze the management variables, timing and outcomes of facial fractures in elderly patients (aged 70 years or more) at several European departments of oral and maxillofacial surgery. The results of this multicenter collaboration on maxillofacial trauma management in the elderly over a 5-year period are presented in this study.

MATERIALS AND METHODS

The present multicenter study was conducted in 12 European departments of oral and maxillofacial surgery (Table 1).

This study was based on a systematic computer-assisted database that allowed the recording of data from all geriatric patients (70 years or more) with facial fractures from the participating maxillofacial surgical units across Europe between January 1, 2013, and December 31, 2017.

The following data were recorded for each patient: gender, age, comorbidities, site of facial fractures, synchronous body injuries, timing of intervention, type of intervention (no treatment or "expectative", closed treatment, open reduction and internal fixation), length of hospital stay, outcome and complications.

The Facial Injury Severity Score according to Bagheri et al¹¹ was calculated for each patient. Facial fractures were determined from computed tomography scans at admission to the hospital and classified as fractures of the mandible, orbital-zygomatic-maxillary complex (MZO), orbit, nose, Le Fort, frontal sinus, and naso-orbital-ethmoid (NOE) fracture. Orbital fractures were sub-classified according to the involved walls. Fractures of the mandible were sub-classified into fractures of the symphysis, parasymphysis, body, angle, ramus, coronoid, or condyle. Data regarding the timing of intervention (within 24 h from hospitalization, between 24 and 72 h, after 72 h from hospitalization) were collected. As a retrospective study, outcome was rated good in case of re-establishment of an appropriate (more than 3 cm) and painless mouth opening, without complications. The following were considered as complications: infraorbital nerve paresthesia, inferior alveolar nerve paresthesia, infection, visual disturbances such as diplopia or loss of visual acuity, fracture malunion, dehiscence.

Patient characteristics were analyzed using descriptive statistics. Statistical analysis was used to search for associations among multiple variables. Statistical significance was determined using the X2 test or the Fisher exact test, if the sample sizes were too small. Statistical significance was set at 0.05. The Helsinki Declaration guidelines were followed, according to local laws. The study was exempt from requiring institutional review board approval as a retrospective study, according to a local institution.

RESULTS

A total of 1334 patients (599 male and 735 female patients) were included in the study. Mean age was 79.3 years (range, 70 to 100 years).

Within the study population, 66% of patients (881) reported one or more comorbidities, the most frequent being hypertension (50%), followed by diabetes (14%), atrial fibrillation (9%), heart ischemic disease (6%), and dementia (5%).

MZO fractures were the most frequently observed injuries with 515 fractures, followed by mandibular fractures (414 fractures), orbital fractures (373), Le Fort fractures (174), nasal fractures (165), and frontal sinus fractures (30). The FISS mean score of the whole study population was 1.88 (range, 1 to 14).

Concomitant injuries were observed in 27.3% of patients (364 patients). Most frequently observed concomitant injuries were orthopedic injuries (172 patients), followed by encephalic (155), thoracic (48), and ocular injuries (44).

On the whole, 665 patients underwent closed or open surgical treatment, whereas in 669 cases an expectative approach without surgery or closed treatment was decided. Among the 665 treated patients, 174 (26%) subjects underwent closed treatment (such as MMF or zygomatic arch closed reduction) whereas 491 patients (74%) underwent open reduction and internal fixation (ORIF) of facial fractures.

Timing of treatment since hospital admission is depicted in Table 2 and Figure 1: almost half of the patients (44% and 43%, respectively) underwent closed treatment or ORIF beyond 72 hours after hospital admission.

A statistically significant association (p < .005) was found between the presence of concomitant injuries and a prolonged (beyond 72 hours) time between hospital admission and treatment.

Table 3 shows the relationship between the performance of treatment and the following variables: age, FISS, and comorbidities. The absence of indications for treatment (an expectative approach) was significantly associated with comorbidities (p < .000005) and an older age (p < .000005). Mean length of hospital stay in specialized medical care was 4.49 days (range, 0 - 7; median, 3; SD, 5.1). Figure 2 shows the mean FISS values and mean hospital stay in the study centers.

The relationship between the length of hospital stay and the variables age, FISS, and comorbidities is outlined in Table 4. A longer hospital stay was significantly associated with a higher FISS (p < .000005), and with the presence of concomitant injuries (p < .000005). Twenty two patients (1.6%) died during their hospital stay: in 18 of these cases, severe concomitant injuries had been diagnosed (14 encephalic, 6 spine, 6 thoracic).

In 1165 cases, a good outcome was obtained with no complications. The most frequently observed complications were: infraorbital nerve paresthesia (45 patients), inferior alveolar nerve paresthesia (27 patients), infection (24 patients), visual disturbances such as diplopia or loss of visual acuity (14 patients).

DISCUSSION

The objective of the present multicenter study was to assess the management and treatment outcome of facial fractures in the elderly population.

The principles of maxillofacial trauma treatment are almost identical, regardless of age. However, the management of elderly patients with maxillofacial fractures needs appropriate and specific adjustments due to anatomic and physiologic variations, which increases the complexity and surgical risks. First of all, the pre-operative assessment of an aging patient should be focused not only on the facial condition but also (if not above all) on the physiological and general status of such patients.⁶⁻¹¹

When the maxillofacial trauma team judges that complex surgical intervention may represent a threat to life or that the risks of surgery may overwhelm its advantages, withholding treatment may be a prudent alternative.⁶⁻⁸ Surgeons, patients and family members have to acknowledge when form and function can be incompletely restored without substantial interference in the quality of life of elderly patients.⁷⁻⁹ To this aim, the surgical management of mandible fractures interfering with mastication is more likely to be indicated, as nourishment is critical to the health and well-being of the geriatric patient. Instead, the surgical management of a slightly displaced zygomatic fracture without associated functional disorders may be avoided in geriatric patients with severe comorbidities, if the only residual defect is represented by a slight cosmetic alteration. The results of this multicenter study highlight that half of the patients underwent a closed or open treatment, whereas in the other half an expectative approach was decided. It is interesting to note that, among the treated patients, 26% of subjects underwent closed treatment (such as MMF or zygomatic arch closed reduction). Such percentage of closed treatment may represent the first important finding that could confirm the trend to reduce the invasiveness of treatment option in elderly patients that have severe comorbidities. The second interesting finding regards the timing of treatment, which was postponed beyond 72 hours after hospital admission in almost half of the patients. This is explained by the need for additional specialized consultations before surgery in elderly patients with polytrauma, which is confirmed by the statistically significant association between the presence of concomitant injuries and a prolonged time between hospital admission and treatment.

Furthermore, the absence of indications to treatment was significantly associated with comorbidities (p < .000005) and an older age (p < .000005).

As a multicenter retrospective study, no uniform indication for different treatment options could be established. However, the aim of the study was to identify a possible trend, that was in fact the obtained result: when important comorbidities or an older age was encountered, the absence of treatment was more likely to be suggested.

As aforementioned, despite the universal validity of the principles of maxillofacial trauma treatment that can be applied to all trauma victims, surgeons have to consider the real indication in elderly patients on a case by case basis. Facing an elderly trauma patient with

severe comorbidities (for example, Alzheimer disease or dementia, or severe heart disease), the physician may reasonably keep a prudent attitude and an expectative approach, especially if the sequelae consist in minor esthetic alterations without functional compromise. The results confirm that this is a widespread attitude in older patients with severe comorbidities.

As for FISS values and hospital stay, results are quite uniform across the participating European centers, as shown in Figure 2.

This study highlights that longer hospital stays are significantly associated with a higher FISS and with the presence of concomitant injuries (p < .000005). This represents a crucial finding, that confirms that peri-operative management of the polytrauma elderly patient may be more complex than that of younger patients, with in some cases a disproportionate consumption of health care resources.

Some (1.6%) of the elderly trauma patients died during their hospital stay. Such a percentage, although it may be considered low, has still to be assessed, especially since in most of these deaths severe concomitant injuries had been diagnosed. In 1165 cases, a good outcome was obtained with no complications. The most frequently observed complications were: infraorbital nerve paresthesia (45 patients), inferior alveolar nerve paresthesia (27 patients), infection (24 patients), visual disturbances such as diplopia or loss of visual acuity (14 patients).

Therefore, the results of this European multicenter study confirm that the management of facial trauma in the elderly requires selected adjustments in the algorithms commonly applied to the care of trauma victims.⁶⁻¹¹

A prudent attitude and a careful assessment to pre-injury intercurrent diseases, medical history, nutrition, and psychosocial conditions are critical in elderly patients.

Furthermore, surgeons must take into consideration that several older patients and their families may refuse aggressive treatment unless function is seriously impaired.

The withholding of surgical treatment for facial injuries in selected gravely ill patients may be considered when delaying or omitting care will not result in substantial function loss.⁶⁻¹⁰

CONCLUSIONS

In conclusion, the management of maxillofacial trauma in elderly patient is often challenging, as elderly patients require specific attention and multidisciplinary collaboration in the diagnosis and sequencing of trauma treatment. A prudent attitude for treatment may be kept in selected cases, especially when severe comorbidities are associated and function is not impaired. More resources are often needed to be allocated for supportive care during hospitalization and assistive care after discharge in elderly patients.

REFERENCES

- Brucoli M, Boffano P, Pezzana A, Benech A, Corre P, Bertin H, et al. The "European Mandibular Angle" research project: the analysis of complications after unilateral angle fractures. Oral Surg Oral Med Oral Pathol Oral Radiol. 2019;128:14-7.
- Brucoli M, Boffano P, Broccardo E, Benech A, Corre P, Bertin H, et al. The "European zygomatic fracture" research project: The epidemiological results from a multicenter European collaboration. J Craniomaxillofac Surg. 2019;47:616-21.
- Brucoli M, Boffano P, Pezzana A, Benech A, Corre P, Bertin H, et al. The "European mandibular angle" research project: the epidemiologic results from a multicenter European collaboration. J Oral Maxillofac Surg. 2019;77:791.e1-e7.
- Ruslin M, Brucoli M, Boffano P, Benech A, Dediol E, Uglešić V, et al. Motor vehicle accidents-related maxillofacial injuries: a multicentre and prospective study. Oral Surg Oral Med Oral Pathol Oral Radiol. 2019;128:199-204.
- Boffano P, Kommers SC, Karagozoglu KH, Forouzanfar T. Aetiology of maxillofacial fractures: a review of published studies during the last 30 years. Br J Oral Maxillofac Surg. 2014;52:901-6.
- 6. Yamamoto K, Matsusue Y, Murakami K, Horita S, Sugiura T, Kirita T. Maxillofacial fractures in older patients. J Oral Maxillofac Surg. 2011;69:2204-10.
- Marciani RD. Critical systemic and psychosocial considerations in management of trauma in the elderly. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 1999;87:272-80.

- Atisha DM, Burr Tv, Allori AC, Puscas L, Erdmann D, Marcus JR. Facial fractures in the aging population. Plast Reconstr Surg. 2016;137:587-93.
- Shumate R, Portnof J, Amundson M, Dierks E, Batdorf R, Hardigan P. Recommendations

for care of geriatric maxillofacial trauma patients following a retrospective 10year multicenter review. J Oral Maxillofac Surg. 2018;76:1931-6.

- Toivari M, Helenius M, Suominen AL, Lindqvist C, Thorén H. Etiology of facial fractures in elderly Finns during 2006-2007. Oral Surg Oral Med Oral Pathol Oral Radiol. 2014;118:539-45.
- Bagheri SC, Dierks EJ, Kademani D, Holmgren E, Bell RB, Hommer L, et al. Application of a facial injury severity scale in craniomaxillofacial trauma. J Oral Maxillofac Surg. 2006;64:408-14.
- 12. Brucoli M, Boccafoschi F, Boffano P, Broccardo E, Benech A. The Anatomage Table and the placement of titanium mesh for the management of orbital floor fractures. Oral Surg Oral Med Oral Pathol Oral Radiol. 2018;126:317-21.
- 13. Giarda M, Tavolaccini A, Arcuri F, Brucoli M, Benech A. Surgical approach to isolated bilateral orbital floor fractures. Acta Otorhinolaryngol Ital. 2015;35:362-4.
- Arcuri F, Brucoli M, Baragiotta N, Benech R, Ferrero S, Benech A. Analysis of complications following endoscopically assisted treatment of mandibular condylar fractures. J Craniofac Surg. 2012;23:e196-8.
- 15. Brucoli M, Nestola DF, Baragiotta N, Boffano P, Benech A. Maxillofacial fractures:
 Epidemiological analysis of a single-center experience. Otorinolaringologia.
 2018;68:132-7.
- Brucoli M, Boffano P, Magnano M, Mistretta R, Benech R, Benech A. The management of a high-risk patient with edentulous mandibular fractures. Otorinolaringologia. 2019;68:42-4.
- Boffano P, Benech R, Gallesio C, Arcuri F, Benech A. Current opinions on surgical treatment of fractures of the condylar head. Craniomaxillofac Trauma Reconstr. 2014;7:92-100.
- Boffano P, Corre P, Righi S. The Role of intra-articular surgery in the management of mandibular condylar head fractures. Atlas Oral Maxillofac Surg Clin North Am. 2017;25:25-34.

- 19. Kommers SC, Boffano P, Forouzanfar T. Consensus or controversy? The classification and treatment decision-making by 491maxillofacial surgeons from around the world in three cases of a unilateral mandibular condyle fracture. J Craniomaxillofac Surg. 2015;43:1952-60.
- 20. Brucoli M, Boffano P, Bonaso M, Benech A. The management of a Y-shaped fracture of the mandibular ramus. Otorinolaringologia. 2019;69:192-5.

LEGENDS:

Figure 1: Timing of treatment since hospital admission according to type of treatment.

Figure 2: Mean FISS values and mean hospital stay in the study centers

Table 1: Participating centers of maxillofacial surgery

	City	Country
Departments of Oral and Maxillofacial Surgery,	Plovdiv	Bulgaria
Faculty of Dental medicine, Medical University		
Department of Maxillofacial Surgery, University	Zagreb	Croatia
Hospital Dubrava		
Department of Oral and Maxillofacial Surgery,	Aalborg	Denmark
Aalborg University Hospital		
Department of Maxillofacial Surgery, North Estonia	Tallinn	Estonia
Medical Centre Foundation		
Department of Oral and Maxillofacial Surgery,	Helsinki	Finland
Helsinki University Hospital		
Service de Stomatologie et Chirurgie Maxillo-faciale,	Nantes	France
CHU de Nantes		
Oral and Maxillofacial Surgery – Hospital Dentistry	Besançon	France
Unit, University Hospital of Besançon		
Division of Maxillofacial Surgery, University of	Novara	Italy
Eastern Piedmont		
Clinic of Maxillofacial Surgery, School of Dentistry,	Belgrade	Serbia
University of Belgrade		
Department of Maxillofacial and Oral Surgery of the	Ljubljana	Slovenia
University Medical Centre		
Maxillofacial Department, Hospital Universitario	Oviedo	Spain
Central de Asturias		
Department for Oral and Maxillofacial Surgery,	Kiev	Ukraine

		Closed		ORIF	
		Ν	%	Ν	%
	Within 24	73	42%	162	33%
	Within 72	24	14%	119	24%
	Beyond 72	77	44%	210	43%
	TOTAL	174		491	

Table 2: Timing of closed and open treatment since hospital admission

Table 3: Indications for surgery according to decades of age, FISS, and presence of comorbidities

		No treatment	Closed or surgical treatment	
Age	70-79 years	314	440	
	80-89 years	278	200	
	90-99 years	77	25	P < . 000005
FISS		1,65	2,12	p>.05
Comorbidities	NO	173	280	
				P < . 000005

Table 4: Length of hospital stay according to decades of age, FISS, and presence of comorbidities

			Mean hospital stay (days)	
	Age	70-79 years	5	
		80-89 years	4	
0		90-99 years	3	p>.05
	FISS	≤1	3,6	
		>1	5,2	p < .000005
	Comorbidities	NO	4,47	
		YES	4	p>.05
	Concomitant	NO	3,6	
-	injuries			p < .000005



edt_12536_f1.tif



edt_12536_f2.tif