

## **Eccentric Exercise Reduces Upper Trapezius Muscle Stiffness Assessed by Shear Wave Elastography and Myotonometry**

Kisilewicz, Aleksandra; Madeleine, Pascal; Ignasiak, Zofia; Cizek, Bogdan; Kawczynski, Adam; Larsen, Ryan Godsk

*Published in:*  
Frontiers in Bioengineering and Biotechnology

*DOI (link to publication from Publisher):*  
[10.3389/fbioe.2020.00928](https://doi.org/10.3389/fbioe.2020.00928)

*Creative Commons License*  
CC BY 4.0

*Publication date:*  
2020

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Kisilewicz, A., Madeleine, P., Ignasiak, Z., Cizek, B., Kawczynski, A., & Larsen, R. G. (2020). Eccentric Exercise Reduces Upper Trapezius Muscle Stiffness Assessed by Shear Wave Elastography and Myotonometry. *Frontiers in Bioengineering and Biotechnology*, 8, Article 928.  
<https://doi.org/10.3389/fbioe.2020.00928>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.





# Eccentric Exercise Reduces Upper Trapezius Muscle Stiffness Assessed by Shear Wave Elastography and Myotonometry

Aleksandra Kisilewicz<sup>1</sup>, Pascal Madeleine<sup>2</sup>, Zofia Ignasiak<sup>3</sup>, Bogdan Cizek<sup>4</sup>, Adam Kawczynski<sup>1\*</sup> and Ryan Godsk Larsen<sup>2\*</sup>

<sup>1</sup> Department of Paralympics Sports, University School of Physical Education, Wrocław, Poland, <sup>2</sup> Sport Sciences – Performance and Technology, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark, <sup>3</sup> Department of Biostructure, University School of Physical Education, Wrocław, Poland, <sup>4</sup> Department of Descriptive and Clinical Anatomy, Medical University of Warsaw, Warsaw, Poland

## OPEN ACCESS

### Edited by:

Yih-Kuen Jan,  
University of Illinois  
at Urbana–Champaign, United States

### Reviewed by:

Fu-Lien Wu,  
National Yang-Ming University, Taiwan  
Cleiton Augusto Libardi,  
Federal University of São Carlos,  
Brazil  
Ai Lyn Tan,  
University of Leeds, United Kingdom

### \*Correspondence:

Adam Kawczynski  
kawczynski.a@gmail.com  
Ryan Godsk Larsen  
rl@hst.aau.dk

### Specialty section:

This article was submitted to  
Biomechanics,  
a section of the journal  
Frontiers in Bioengineering and  
Biotechnology

**Received:** 14 May 2020

**Accepted:** 20 July 2020

**Published:** 05 August 2020

### Citation:

Kisilewicz A, Madeleine P, Ignasiak Z, Cizek B, Kawczynski A and Larsen RG (2020) Eccentric Exercise Reduces Upper Trapezius Muscle Stiffness Assessed by Shear Wave Elastography and Myotonometry. *Front. Bioeng. Biotechnol.* 8:928. doi: 10.3389/fbioe.2020.00928

In this study, we tested the hypotheses that unaccustomed eccentric exercise (ECC) would reduce the elastic modulus and dynamic stiffness of the upper trapezius muscle and that these changes would correlate with increases in muscle thickness, reflecting muscle edema. Shear wave elastography was used to measure elastic modulus, dynamic stiffness was assessed using myotonometry, and muscle thickness was measured using ultrasonography. All measurements were performed at four locations over the upper trapezius before and 24 h after a single bout of ECC. Fourteen healthy participants (11 males and 3 females;  $23.2 \pm 3.0$  years; height  $175.1 \pm 10.4$  cm; body mass  $73.8 \pm 11.3$  kg) took part in the study. Overall, ECC resulted in decreased elastic modulus (from  $45.8 \pm 1.6$  to  $39.4 \pm 1.2$  kPa,  $p < 0.01$ ) and dynamic muscle stiffness (from  $369.0 \pm 7.3$  to  $302.6 \pm 6.0$  N/m,  $p < 0.01$ ). Additionally, ECC resulted in increased muscle thickness (from  $6.9 \pm 0.4$  to  $7.3 \pm 0.4$  mm,  $p < 0.01$ ). Spatial changes (across the four locations) were found for elastic modulus, stiffness and thickness. No significant correlations were found between changes in measures of muscle stiffness, or between changes in stiffness and changes in thickness. In conclusion, the present pilot study showed that ECC altered biomechanical muscle properties, reflected by decreased elastic modulus and dynamic muscle stiffness 24 h after ECC.

**Keywords:** eccentric exercise, trapezius muscle, muscle stiffness, ultrasonography, myotonometry

## INTRODUCTION

In a biomechanical context, stiffness refers to the resistance of tissue during passive stretching which depends on the type of external force applied and the deformation of the structure caused by this force (Baumgart, 2000). Increased level of stiffness has been related to a higher risk of strain and stress injuries (Pruyn et al., 2012). A recent study on overhead athletes have shown that neck and shoulder pain is related to higher stiffness of upper trapezius (UT) muscle (Leong et al., 2016). More specifically, the shoulder girdle and the trapezius are of special interest in both sport and

occupational perspectives when one considers the high prevalence of neck-shoulder disorders and their socioeconomic burden (Picavet and Hazes, 2003).

Unaccustomed eccentric muscle contractions place high strain on the muscle-tendon complex resulting in neuromuscular functional impairments, muscle damage and finally delayed onset muscle soreness (DOMS) (Guilhem et al., 2010). According to the literature, muscle soreness peaks between 12- and 36-h after exercise (Nosaka et al., 2002). Soreness sensation is related to the inflammation process due to efflux of substances from the damaged tissue to the extracellular space that sensitize free nerve endings and activate groups III and IV pain receptors (Zainuddin et al., 2005). Eccentric exercise (ECC) has also been reported to increase muscle stiffness (Hoang et al., 2007). It has been suggested that post-exercise stiffness is caused by muscle edema related to fiber damage. On the other hand, change in muscle stiffness precedes any evidence of swelling (Proske and Morgan, 2001). Particularly, muscle edema occurs subsequent to the inflammatory response and prior to the first symptoms of DOMS (Peake et al., 2005). Moreover, several studies show no correlation between exercise-induced change in muscle stiffness and typical indicators of muscle damage, such as edema and soreness (Lacourpaille et al., 2014; Yanagisawa et al., 2015). Therefore, there is a need for further investigating the DOMS related changes in mechanical properties of skeletal muscles.

The assessment of biomechanical properties of the musculoskeletal system is challenging because striated muscle is an anisotropic and viscoelastic complex composed of both active and passive structures (Gennisson et al., 2010). However, relatively new methods based on shear wave elastography (SWE) and myotonometry have been reported to provide reliable assessments of biomechanical properties of individual muscles (Lacourpaille et al., 2012; Kawczynski et al., 2018). Ultrasonic SWE provides real-time, direct measurement of muscle elastic modulus. The SWE estimates shear elastic modulus linearly related to Young's modulus, as a substitute for passive muscle stiffness (Lacourpaille et al., 2017; Xie et al., 2019; Zhang et al., 2019). In recent years, the application of SWE for muscle tissue evaluation has increased rapidly, with good-to-excellent intra-observer, inter-observer, and inter-day reliability of SWE for elastic modulus measurements (Tas et al., 2017). The SWE methodology has been proposed to non-invasively quantify exercise-induced muscle damage, with relevance for clinical evaluation of athletes (Lacourpaille et al., 2017). Myotonometry is an alternative method that quantifies soft tissue viscoelastic properties, including muscle stiffness. Importantly, myotonometry is less expensive than SWE, hand-held and easy to use, with application for use in sport fields and every-day evaluation of training or treatment. Similarly, to SWE, myotonometry has been successfully employed in the evaluation of viscoelastic muscle parameters (Gervasi et al., 2017), showing good validity and high reliability for e.g., the trapezius muscle (Viir et al., 2011; Kawczynski et al., 2018). Although both SWE and myotonometry operate using the principle of Young's modulus, the depth of measurements varies, as SWE provide measurements of elastic modulus (i.e., passive muscle stiffness) from deeper structures, while myotonometry

measures dynamic muscle stiffness superficially (Kelly et al., 2018). Further, these measurements made at defined locations enabled inverse-distance-based interpolation to generate topographical maps of the viscoelastic properties (Kawczynski et al., 2018; Heredia-Rizo et al., 2020).

To date, no prior study has compared the spatial changes in biomechanical properties of skeletal muscle, using both SWE and myotonometry, following unaccustomed eccentric muscle contractions. Therefore, the present study was designed to quantify the spatial changes of UT elastic modulus, muscle stiffness and thickness, 24 h after unaccustomed ECC of UT. Based on our previous work (Kawczynski et al., 2018), we hypothesized that ECC would reduce the elastic modulus and stiffness of UT and that these changes would correlate with increases in muscle thickness, reflecting exercise-induced muscle edema.

## MATERIALS AND METHODS


### Participants

Fourteen participants (11 males and 3 females; aged  $23.2 \pm 3.0$  years; height  $175.1 \pm 10.4$  cm; body mass  $73.8 \pm 11.3$  kg) volunteered to take part in this pilot study. All participants were recruited from mixed student population, with no professional training experience in disciplines involving overhead actions or occupations with significant involvement of the upper limbs. All participants were right-handed and maintained normal daily activity during the course of the study. The inclusion criteria were: no pain in the shoulder region prior to the experiment, no history of neck or shoulder disorders, and no strength training in the past month. The study was reviewed and approved by the Research Ethics Committee of North Denmark (N-2016-0023). The participants provided their written informed consent to participate in this study.

### Study Design

All measurements were carried out by the same investigator, trained in muscle ultrasonography and myotonometry. The study was performed in the spring season, with an average indoor temperature of 25°C. The Young's modulus and stiffness of UT muscle on the dominant side, reported by each participant prior to the experiment. During the experiment, participants sat on a chair with the back supported in an upright position. Forearms were supported on the table, and the head was facing forward. The skin overlying UT muscle was exposed for testing. Thickness measurements were acquired first followed by myotonometry and SWE measurements. The measurements were performed twice: before the ECC protocol, and  $24 (\pm 1)$  h post ECC.

Myotonometry, SWE and muscle thickness measurements were performed on the same four points marked on the skin overlying the UT muscle (**Figure 1**). To properly locate four points of measurement, the spinous process of the C7 vertebra and the angle of acromion were identified by palpation. The distance between C7 spinous process and acromion: "d," was used to calculate the between-points distance (mean  $21.8 \pm 1.8$  cm). Adjacent points were separated by 1/6 (mean  $3.6 \pm 0.3$  cm) of



**FIGURE 1** | Location of the four measurement points marked on the skin overlying the upper trapezius muscle for the assessment of muscle dynamic stiffness measured by myotonometry, elastic modulus measured by shear-wave elastography and thickness measured by ultrasonography C7, 7th cervical vertebrae spinous process; Acr., acromion; “d”, the distance between C7 spinous process and acromion.

“d” distance (Kawczynski et al., 2018). The B-mode ultrasound images were used to locate the greatest muscle thickness and confirm correct points location. The probe of the MyotonPRO device was positioned directly over each point. The device would only start the measurement when the investigator applied the appropriate pre-compression force and the probe was placed perpendicular to the surface of the skin. For SWE measurements, the transducer was oriented in parallel to the UT muscle fibers, with the center positioned over the skin mark. During data acquisition, the transducer was held stationary for 10 s during which the SWE sonogram was recorded with minimal pressure applied on the skin (Kot et al., 2012). A rectangular region of interest (ROI) selected to determine the shear elastic modulus of the UT muscle was defined by the thickness of the muscle belly, excluding aponeurosis and unfilled regions within the elasticity map (Ates et al., 2015). Due to differences in anatomy, there were slight variation in the ROI used across participants. All ultrasonic images were imported to a personal laptop and given specific codes in order to blind the identity of the data. This process was done by an investigator who was not involved in data acquisition or analyses. To obtain highest reliability of measurements we used an average of three SWE elasticity measurements of elastic modulus (Kelly et al., 2018), and a MyotonPRO Multiscan mode of five continuous measurements of dynamic muscle stiffness (Vain and Kums, 2002).

Three LOGIQ longitudinal view images of UT muscle belly at C7 level were performed before and 24 h post ECC. Thickness of UT was determined using MicroDicom viewer software (MicroDicom DICOM Viewer, Bulgaria). Specifically, four parallel straight lines were drawn between the superficial and deep fascia of UT muscle belly. Averages from three images for each of four measurement points were calculated and used for further analyses. The muscle stiffness, SWE elastic modulus and thickness maps were created using the averaged values for each

of the assessed points (Figure 3). The interpolation was done using an inverse distance weighted interpolation to obtain a 3D graphical representation (Binderup et al., 2010b).

In addition to measurements of UT dynamic stiffness, elastic modulus, and thickness, measurements of range of movement (ROM) for shoulder elevation, maximum voluntary contraction (MVC) in isometric condition, and soreness intensity was obtained before and 24 h post ECC. Each participant reported individual soreness level, using a 10-cm standardized visual analog scale where 0 indicated “no soreness” and 10 “maximal soreness intensity.”

## Ultrasound Measurements

An ultrasound system (LOGIQ S8, General Electric, Norwalk, CT, United States) was used for acquiring images (B-mode, 8.5–10.0 MHz) of the UT muscle. For the assessment of UT muscle thickness, the LOGIQ longitudinal view feature was used to acquire an image over the full length of the UT muscle at the C7 level of the cervical spine. Specifically, a linear transducer (9L) was positioned over the right UT, parallel to the arrangement of muscle fibers for measurements of thickness and SWE. The orientation of the transducer longitudinal to muscle fibers is required to achieve accurate and reliable measurements (Gennisson et al., 2010). The velocity of the shear wave propagation tracked by a pulse-echo ultrasound, provides a characterization of elastic properties as velocity of shear waves increase with increase in passive muscle stiffness (Eby et al., 2013). Assuming a linear and elastic behavior, the shear elastic modulus (kPa) is a function of shear velocity ( $V_s$ ) as follows:  $E = \rho V_s^2$ , where  $\rho$  is the density of muscle ( $1000 \text{ kg/m}^3$ ) (Ates et al., 2015).

## Myotonometry Measurements

The MyotonPRO (Myoton AS, Myoton Ltd., Estonia) is a non-invasive, portable device using superficial mechanical deformation for assessment of biomechanical characteristics of soft tissues (Aird et al., 2012). The device has been used to measure dynamic stiffness (N/m) of trapezius muscles, quantified with the damped oscillation method (Viir et al., 2011). Dynamic stiffness is defined as the resistance of a soft tissue to a contraction or an external force that deforms its initial shape (Schneider et al., 2015). The probe of the device was placed perpendicular to the surface of the skin overlying the UT, and applied slight pressure with a short mechanical impulse (0.4 N for 15 ms), with a constant pre-compression force of 0.18 N. The mechanical impulse generated damped oscillations within the muscle soft tissue, recorded by an accelerometer. The acceleration signal is processed to provide an oscillation curve from which the dynamic stiffness is calculated as follows: the period  $T$ , is defined as the time elapsed between the first two adjacent acceleration peaks following the mechanical impulse; the oscillation frequency  $f$  is calculated from the equation:  $f = 1/T$ ; the angular frequency  $\omega$  ( $\omega = 2\pi f$ ), is related to stiffness  $K$  and mass  $m$ , and allows for calculating stiffness:  $K = \omega^2 m$ , substituting into:  $K = 4\pi^2 f^2 m$  (Sohirad et al., 2017).



## Eccentric Exercise Protocol

A dynamic shoulder dynamometer (Aalborg University, Aalborg, Denmark) was used to induce DOMS in the UT muscle (Madeleine et al., 2006). The dynamic shoulder dynamometer consists of an actuator, a load cell, a control unit, a cylinder, a shoulder contact pad, and an adjustable seat fixed on a stainless-steel frame. The ECC protocol consisted of 50 ECC performed in five bouts of 10 contractions at 100% level of maximum voluntary contraction (MVC), separated by a 2-min rest (Kawczynski et al., 2012). Due to the one-sided design of the dynamometer, the ECC was performed only with the right (dominant) UT muscle.

Before exercising, the ROM for shoulder elevation was assessed. All participants were asked to elevate their right shoulder to their highest position and then lower it as much as possible. Both positions were measured and recorded by the dynamometer. The ROM was calculated as the difference between the highest and lowest position. In the eccentric mode, the dynamometer produces a constant vertical downward force at a given level of MVC. Three trials were performed to determine the MVC in neutral position, i.e., the maximal shoulder elevation force under isometric conditions for 3 s, separated by 2 min rest. The contact point between the dynamometer and shoulder was approximately 3 cm medial to the acromion. The contact pad of the shoulder dynamometer was padded to avoid inhibition of force exertion because of pressure pain (Kawczynski et al., 2007). While exercising, participants were instructed to counteract the vertical force, generated by the dynamometer, along the shoulder ROM. All participants wore a corset during the exercise to prevent lateral bending.

## Statistical Analysis

The sample size was calculated using G\*Power software (Kiel University, Kiel, Germany) with an expected “medium” effect size ( $f^2 = 0.25$ ) for changes over time,  $\alpha$  level of 0.05, power ( $1-\beta$ ) of 0.9, and correlation for repeated measures of 0.6 (Faul et al., 2007; Kawczynski et al., 2018). Sample size calculations revealed that 11 participants were required. Accounting for potential dropouts, 14 participants were recruited. The dynamic stiffness measured with myotonometry, elastic modulus assessed by SWE, and muscle thickness were introduced in a two-way repeated measure analysis of variance (RM-ANOVA). Time (before and 24 h post ECC) and points (1–4) marked on the skin overlying UT muscle were introduced as within-subject factors. Soreness intensity, ROM, and MVC were analyzed using a 1-way RM-ANOVA with time (before and 24 h post ECC) as within-subject factor. Normality of the data distribution was tested by Shapiro–Wilk tests. If there was an interaction between variables, Bonferroni adjustment for multiple comparisons was used for *post hoc* tests. Pearson’s product-moment correlation was used to assess the relationship between relative changes in stiffness measured by MyotonPRO and elastic modulus determined by SWE, and the association between relative changes in muscle stiffness and muscle thickness. These analyses were done for each of the four points of measurement. For all statistical tests,  $p$ -value  $< 0.05$  was considered significant. Data are presented as mean  $\pm$  SD in the text and the figures. All statistical analyses were

performed using SPSS 22.0 software (IBM SPSS Inc., Armonk, NY, United States).

## RESULTS

### Myotonometry

There was a time-by-point interaction for muscle dynamic stiffness ( $F_{1,13} = 7.1$ ;  $p = 0.022$ ). *Post hoc* analyses revealed that muscle dynamic stiffness decreased from before ( $369.0 \pm 7.3$  N/m) to 24 h post ECC ( $302.6 \pm 6.0$  N/m) for all four points of measurement ( $p < 0.001$ ) (Figure 2). There was a main effect of point for muscle dynamic stiffness ( $F_{1,13} = 62.3$ ;  $p < 0.001$ ). *Post hoc* analyses showed that muscle dynamic stiffness was higher in point 4 compared with all other points (all  $p < 0.001$ ).

### Ultrasonography

There was no time-by-point interaction for SWE muscle elastic modulus ( $F_{1,13} = 0.1$ ,  $p = 0.943$ ). However, there was a main effect of time for SWE muscle elastic modulus ( $F_{1,13} = 12.4$ ;  $p = 0.005$ ), such that muscle elastic modulus decreased from before ( $45.8 \pm 1.6$  kPa) to 24 h post ECC ( $39.4 \pm 1.2$  kPa) (Figure 2). There was also a main effect of point for muscle elastic modulus ( $F_{1,13} = 5.4$ ;  $p = 0.022$ ), such that muscle elastic modulus was higher in point 1 compared with point 2 ( $p < 0.05$ ). There was no time-by-point interaction for muscle thickness ( $F_{1,13} = 0.1$ ,  $p = 0.97$ ). However, there was a main effect of time for muscle thickness ( $F_{1,13} = 7.9$ ;  $p < 0.001$ ), such that muscle thickness increased from before ( $6.9 \pm 0.4$  mm) to 24 h post ECC ( $7.3 \pm 0.4$  mm) (Figure 2). There was also a main effect of point ( $F_{1,13} = 7.7$ ;  $p < 0.001$ ) such that muscle thickness was lower for point 1 compared with points 2 and 3 ( $p < 0.05$ ).

### Soreness Intensity, Maximal Voluntary Contraction, Range of Motion


Soreness intensity increased from before ( $0.0 \pm 0.0$ ) to 24 h after ECC ( $4.6 \pm 1.4$ ;  $p < 0.001$ ). There was no significant change in MVC from before ( $622.4 \pm 243.0$  N) to 24 h post ECC ( $521.7 \pm 239.2$  N;  $p = 0.10$ ). Similarly, there was no change in shoulder elevation ROM from before ( $67.9 \pm 16.4$  mm) to 24 h post ECC ( $66.0 \pm 9.2$  mm;  $p = 0.70$ ).

### Correlations Among Stiffness Measures and Between Stiffness and Thickness

Pearson’s product-moment correlations showed no correlation between relative changes in stiffness measured by the MyotonPRO and SWE (Table 1). Also, there were no correlations between relative changes in SWE and muscle thickness, or relative changes in myotonometry and muscle thickness (Table 1).

## DISCUSSION

In this pilot study, we showed a decrease in both dynamic muscle stiffness assessed by myotonometry and the elastic modulus




measured by SWE at 24 h after ECC. Accompanying these changes, UT muscle thickness was increased at 24 h after ECC, indicating muscle edema (**Figure 2**). Our results confirmed the hypothesis concerning spatial changes in UT elastic modulus, stiffness and thickness 24 h after ECC. However, we found no significant correlations between decreases in muscle stiffness measured by SWE and myotonometry, or between decreases in stiffness and increases in thickness.

## Effect of ECC on Muscle Stiffness

In line with recent studies, we found a reduction in elastic modulus and dynamic stiffness (Andonian et al., 2016; Kawczynski et al., 2018; Xu et al., 2019). Kawczynski et al. (2018) observed a decrease in UT muscle dynamic stiffness at 24 h after a single bout of ECC, using the same exercise protocol. They also investigated test-retest reliability of MyotonPRO device, obtaining ICC (95% CI) values from 0.59 to 0.96, with standard error of measurement ( $\pm$ SD) of 18.5 ( $\pm$ 7.3) and minimal detectable change ( $\pm$ SD) of 51.2 ( $\pm$ 20.3), respectively. Andonian et al. (2016) demonstrated that prolonged, low-intensity, and mainly ECC induced a decrease in quadriceps muscle stiffness quantified by SWE. They showed good-to-high intra-session reliability (ICC = 0.88–0.92) of the shear modulus, with standard error of measurement from 0.12 to 0.20. Although the ECC protocol differed from our study, we used the same measurement conditions, such that the muscle

was investigated in slack position (i.e., no muscle tension). As proposed by Andonian et al. (2016), the measurement conditions (slack versus stretched) could have contributed to post-exercise changes in elastic modulus. Consistent with this interpretation, Lacourpaille et al. (2017) suggested that the effect of ECC on muscle stiffness is muscle length dependent. Another study by Lacourpaille et al. (2014) showed a significant increase in elastic modulus of the elbow extensors, but only when the investigated muscles were in a stretched position. This could be explained by perturbations of calcium homeostasis, as muscle fiber sensitivity to  $\text{Ca}^{2+}$  increases with muscle elongation. Therefore, measurements performed in slack position versus stretched position could possibly contribute to discrepancies between studies (Green et al., 2012; Lacourpaille et al., 2017; Xie et al., 2019). However, the reduction in elastic modulus and dynamic stiffness stands inconsistent with the general idea of an increase in muscle stiffness after a single bout of ECC (Green et al., 2012; Heredia-Rizo et al., 2020). Green et al. (2012) reported a 21% increase in gastrocnemius medialis muscle elastic modulus, but no significant changes in soleus muscle after a single bout of ECC. Nevertheless, they used a different measurement technique (i.e., magnetic resonance elastography). Furthermore, as proposed by the authors, differences in the articular anatomy and fiber composition of various muscles may contribute to muscle dependent changes in stiffness (Green et al., 2012). In line with this idea, Xu et al. (2019) showed a significant



decrease in elastic modulus of the vastus lateralis, but not the other quadriceps muscle heads after a single bout of 75 knee flexions performed in a dynamometer.

## The Relationship Between Post-exercise Changes of Muscle Stiffness and Thickness and Correlations Among Stiffness Measures

By definition, the shear elastic modulus is a function of muscle tissue density and shear wave propagation speed (Ates et al., 2015). Thereby, exercise-induced changes in muscle tissue density could affect elastic modulus after ECC. It is known that the rate of collagen turnover increases with exercise (Hjorth et al., 2015). The elevated collagen turnover increases the density of extracellular matrix (ECM) within muscle fibers, causing a highly negative interstitial fluid pressure, and subsequently a muscle edema (Kjaer, 2004). Likewise, according to Hyl Dahl and Hubal (2014) and Mackey et al. (2004), the changes in muscle biomechanical properties following unaccustomed ECC are most likely related to the remodeling of the ECM. However, it is surprising how little is known about ECM remodeling, despite its' important role in myofibrillar adaptation to exercise (Kjaer, 2004). As proposed by Andonian et al. (2016), an increase in muscle stiffness after ECC could be counterbalanced by changes in extracellular water volume. As suggested by Madeleine et al. (2018), changes in the muscle physical milieu at 24 h after ECC

could be related to increased muscle thickness. However, we observed no significant correlation between changes in muscle stiffness and thickness. Similarly, there was no correlation among stiffness measures, although both SWE and myotonometry showed a decrease in UT muscle stiffness 24 h after ECC. In agreement with our results, Akagi and Kusama (2015) showed no correlation between stiffness measured by SWE and a mechanical stiffness meter in the assessment of neck and shoulder muscles. On the contrary, Kelly et al. (2018) observed a significant positive correlation between myotonometry and SWE in the evaluation of infraspinatus, erector spinae and gastrocnemius muscles stiffness at rest. However, none of these studies assessed the correlation between changes in stiffness parameters after ECC. Due to the small sample size in this pilot study, there is a need for further research with larger sample sizes to investigate the relationship between muscle stiffness assessed by SWE and myotonometry.

## Post-exercise Changes of Soreness Intensity, Muscle Force, Range of Shoulder Elevation and the Spatial Heterogeneity in UT Muscle Stiffness and Thickness

To induce DOMS, we used the ECC protocol established and validated in previous studies (Madeleine et al., 2006, 2011, 2018; Kawczynski et al., 2007, 2012, 2018). We observed an increase in



**TABLE 1** | Correlation coefficients between relative changes in upper trapezius muscle dynamic stiffness measured by myotonometry, elastic modulus measured by shear-wave elastography and muscle thickness measured by ultrasonography, respectively.

Point	Mean ± SD		Mean ± SD		Mean ± SD		Correlation coefficients					
	Dynamic stiffness (N/m)		Elastic modulus (kPa)		Thickness (mm)		Δ Dynamic stiffness (%) / Δ elastic modulus (%)	Δ Dynamic stiffness (%) / Δ thickness (%)	Δ Elastic modulus (%) / Δ thickness (%)			
	Before	24 h after ECC	Before	24 h after ECC	Before	24 h after ECC	<i>R</i>	<i>p</i>	<i>R</i>	<i>p</i>	<i>R</i>	<i>p</i>
1	309.29 ± 42.83	268.64 ± 17.72	41.96 ± 10.67	38.82 ± 11.40	5.60 ± 1.84	6.11 ± 1.96	0.08	0.78	−0.08	0.79	−0.26	0.38
2	327.07 ± 63.95	260.64 ± 31.55	47.83 ± 9.13	46.19 ± 5.99	7.19 ± 1.38	7.50 ± 1.46	−0.14	0.64	0.42	0.14	0.04	0.90
3	373.71 ± 68.79	248.64 ± 25.38	41.69 ± 9.61	40.74 ± 10.84	7.53 ± 1.42	7.87 ± 1.44	−0.25	0.39	0.14	0.63	0.08	0.79
4	500.43 ± 65.22	424.36 ± 83.24	42.47 ± 10.85	40.60 ± 13.11	7.33 ± 1.88	7.67 ± 1.63	0.50	0.07	−0.44	0.11	−0.10	0.73

soreness intensity 24 h after ECC in line with previous research (Madeleine et al., 2011, 2018), but no changes in MVC and ROM. The lack of changes in MVC and ROM can be explained by recruitment of additional motor unit, increased motor unit synchronization or altered contractile properties (Madeleine et al., 2011; Penailillo et al., 2015). In agreement with current findings, Nosaka et al. (2002) showed that DOMS appears to be independent of decline in MVC and ROM. Moreover, Yanagisawa et al. (2015) reported the absence of significant correlations between muscle stiffness evaluated by elastography and other exercise-induced muscle damage indicators, such as decreased joint ROM and muscle soreness. However, their results showed a significant increase in serum creatine kinase activity, suggesting that muscle damage did occur as a result of ECC. Similarly, Kawczynski et al. (2018) observed no decline in MVC of the UT muscle, despite a significant decrease in muscle belly stiffness and an increase in soreness intensity. In addition, we showed spatial heterogeneity in UT muscle dynamic stiffness, elastic modulus and thickness depicted in the 3D topographical maps (Figure 3) underlining the relevance of a bioengineering approach to map the ROI (Binderup et al., 2010a). We observed a consistent spatial distribution from before to 24 h after ECC for all three parameters, i.e., the UT dynamic stiffness remained highest for point 4 (most distal) after ECC, while elastic modulus showed highest values in more proximal regions (point 2). As suggested in previous studies (Kelly et al., 2018; Madeleine et al., 2018), point 2 location corresponds to a thicker muscle belly site, where the SWE technique provides measurements from deeper tissue structures. The most distal point (point 4) is located closer to a superficial musculotendinous site, where stiffness of superficial structures can be assessed with myotonometry (Kawczynski et al., 2018). The spatial heterogeneity may be explained by differences in local structural features such as pennation angles, and fiber type (Damon et al., 2008). As suggested by other authors, – the spatial heterogeneity in elastic modulus could also result from non-uniform muscle damage induced by ECC (Green et al., 2012) or heterogeneous activation of muscle fibers during exercise (Kinugasa et al., 2006). Furthermore, Alfuraih et al. (2018) confirmed that the depth of measurement may influence the SWE elastic modulus. They showed that variability of the measurements increases quadratically with the depth of acquisition. Due to the anatomy of UT muscle, the depth

of measurements could differ across the four points of data acquisition causing spatial heterogeneity in UT elastic modulus.

## Strengths and Limitations

The selection of UT as the muscle of interest was based on the fact that this body region is greatly affected by DOMS and musculoskeletal pain (Madeleine et al., 2018). As suggested in previous research, monitoring of UT muscle stiffness is essential for the prevention of rotator cuff tendinopathy and other sport-related injuries (Leong et al., 2016). Therefore, our results add new important information for clinicians and applied researchers with a focus on injury prevention and rehabilitation. Showing how a single muscle adapts to ECC could also support training load programming. Conducting multiple, instead of a single, measurement for each point for SWE and myotonometry, allowed us to obtain high reliability of these measurements (Vain and Kums, 2002; Kelly et al., 2018). All ultrasound measurements were blinded prior analysis, which allowed us to avoid subjectivity bias. As reported by Andonian et al. (2016) measurements performed in a slack resting position are more reliable and easier to standardize. However, as showed by Huang et al. (2018), testing in a functional position could better reflect real situations. Therefore, performing our measurements only in a resting position should be considered as a first limitation of the current study. Secondly, we did not investigate the intra-operator and day-to-day reliability of myotonometry and SWE in measuring the UT muscle stiffness. We based our reliability on previous results from our research group (Kawczynski et al., 2018), and others (Andonian et al., 2016). Therefore, we acknowledge the need for further research including the test-retest reliability, coefficient of variation and typical error. Thirdly, we conducted our measurements only 24 h after ECC. Adding measurements immediately after ECC would inform about the specific effects of exercise on the results. Notably, Kawczynski et al. (2018) showed that UT muscle stiffness decreased by 6.5% immediately after ECC, and by 14.2% at 24 h after ECC. The small change in stiffness immediately after ECC suggests that exercise in itself had a small influence on the results at 24 h. Moreover, according to Alfuraih et al. (2017), using different ROI sizes, probe orientations and measurement locations produce variability of SWE estimation. These factors can be considered as limitations. However, as proved by the same authors, SWE demonstrates

the strongest internal agreement when the probe is placed longitudinally to the muscle fibers, with ROI of medium-to-large size, excluding myotendinous or myoaponeurotic structures (Alfuraih et al., 2017), which is in line with our settings. Finally, due to differences in anatomy, the available area for calculating SWE was not always uniform across subjects and some variations may have occurred due to the dimensions of the ROI.

## CONCLUSION

This pilot study shows for the first time that biomechanical muscle properties, represented by elastic modulus and dynamic muscle stiffness, decreased 24 h after ECC. Moreover, we present a novel 3D topographical maps showing spatial heterogeneity of muscle dynamic stiffness, elastic modulus and thickness, which may provide insight about within-muscle changes following ECC. Findings from the present study demonstrate that monitoring UT muscle stiffness using both myotonometry and SWE contribute to the understanding of how a single muscle adapts to ECC.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## REFERENCES

- Aird, L., Samuel, D., and Stokes, M. (2012). Quadriceps muscle tone, elasticity and stiffness in older males: reliability and symmetry using the MyotonPRO. *Arch. Gerontol. Geriatr.* 55, e31–e39. doi: 10.1016/j.archger.2012.03.005
- Akagi, R., and Kusama, S. (2015). Comparison between neck and shoulder stiffness determined by shear wave ultrasound elastography and a muscle hardness meter. *Ultrasound. Med. Biol.* 41, 2266–2271. doi: 10.1016/j.ultrasmedbio.2015.04.001
- Alfuraih, A. M., O'Connor, P., Hensor, E., Tan, A. L., Emery, P., and Wakefield, R. J. (2018). The effect of unit, depth, and probe load on the reliability of muscle shear wave elastography: variables affecting reliability of SWE. *J. Clin. Ultrasound.* 46, 108–115. doi: 10.1002/jcu.22534
- Alfuraih, A. M., O'Connor, P., Tan, A. L., Hensor, E., Emery, P., and Wakefield, R. J. (2017). An investigation into the variability between different shear wave elastography systems in muscle. *Med. Ultrasonogr.* 19, 392–400. doi: 10.11152/mu-1113
- Andonian, P., Viallon, M., Le Goff, C., De Bourguignon, C., Tourel, C., Morel, J., et al. (2016). Shear-wave elastography assessments of quadriceps stiffness changes prior to, during and after prolonged exercise: a longitudinal study during an extreme mountain ultra-marathon. *PLoS One* 11:e0161855. doi: 10.1371/journal.pone.0161855
- Ates, F., Hug, F., Bouillard, K., Jubeau, M., Frappart, T., Couade, M., et al. (2015). Muscle shear elastic modulus is linearly related to muscle torque over the entire range of isometric contraction intensity. *J. Electromyogr. Kinesiol.* 25, 703–708. doi: 10.1016/j.jelekin.2015.02.005
- Baumgart, E. (2000). Stiffness—an unknown world of mechanical science? *Injury* 31(Suppl. 2), B14–B23.
- Binderup, A. T., Arendt-Nielsen, L., and Madeleine, P. (2010a). Cluster analysis of pressure pain threshold maps from the trapezius muscle. *Comput. Methods. Biomech. Biomed. Engin.* 13, 677–683. doi: 10.1080/10255840903446979

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Research Ethics Committee of North Denmark (N-2016-0023). The study was conducted in accordance with the Declaration of Helsinki. Informed consent was obtained from each participant. The patients/participants provided their written informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

AKi, PM, AKa, and RL planned the study. PM and RL carried out the phases of the protocol and supervised the stiffness and thickness measurements. AKa performed the statistical analysis. AKi collected the data and drafted the manuscript. ZI and BC supervised methods and manuscript preparation. All authors read and approved the final version of the manuscript.

## FUNDING

The study was partly supported by the Danish Rheumatism Association (R77-5214).

## ACKNOWLEDGMENTS

We thank the volunteers for their participation in the study.

- Binderup, A. T., Arendt-Nielsen, L., and Madeleine, P. (2010b). Pressure pain threshold mapping of the trapezius muscle reveals heterogeneity in the distribution of muscular hyperalgesia after eccentric exercise. *Eur. J. Pain.* 14, 705–712. doi: 10.1016/j.ejpain.2009.11.001
- Damon, B. M., Wadington, M. C., Lansdown, D. A., and Hornberger, J. L. (2008). Spatial heterogeneity in the muscle functional MRI signal intensity time course: effect of exercise intensity. *Magn. Reson. Imaging* 26, 1114–1121. doi: 10.1016/j.mri.2008.01.023
- Eby, S. F., Song, P., Chen, S., Chen, Q., Greenleaf, J. F., and An, K. N. (2013). Validation of shear wave elastography in skeletal muscle. *J. Biomech.* 46, 2381–2387. doi: 10.1016/j.jbiomech.2013.07.033
- Faul, F., Erdfelder, E., Lang, A. G., and Buchner, A. (2007). G\*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav. Res. Methods* 39, 175–191. doi: 10.3758/bf03193146
- Gennissou, J. L., Deffieux, T., Mace, E., Montaldo, G., Fink, M., and Tanter, M. (2010). Viscoelastic and anisotropic mechanical properties of in vivo muscle tissue assessed by supersonic shear imaging. *Ultrasound. Med. Biol.* 36, 789–801. doi: 10.1016/j.ultrasmedbio.2010.02.013
- Gervasi, M., Sisti, D., Amatori, S., Andreazza, M., Benelli, P., Sestili, P., et al. (2017). Muscular viscoelastic characteristics of athletes participating in the European Master Indoor Athletics Championship. *Eur. J. Appl. Physiol.* 117, 1739–1746. doi: 10.1007/s00421-017-3668-z
- Green, M. A., Sinkus, R., Gandevia, S. C., Herbert, R. D., and Bilston, L. E. (2012). Measuring changes in muscle stiffness after eccentric exercise using elastography. *NMR Biomed.* 25, 852–858. doi: 10.1002/nbm.1801
- Guilhem, G., Cornu, C., and Guevel, A. (2010). Neuromuscular and muscle-tendon system adaptations to isotonic and isokinetic eccentric exercise. *Ann. Phys. Rehabil. Med.* 53, 319–341. doi: 10.1016/j.rehab.2010.04.003
- Heredia-Rizo, A. M., Petersen, K. K., Arendt-Nielsen, L., and Madeleine, P. (2020). Eccentric training changes the pressure pain and stiffness maps of the upper

- trapezius in females with chronic neck-shoulder pain: a preliminary study. *Pain Med.* pnz360. doi: 10.1093/pmn/pnz360
- Hjorth, M., Norheim, F., Meen, A. J., Pourteymour, S., Lee, S., Holen, T., et al. (2015). The effect of acute and long-term physical activity on extracellular matrix and serglycin in human skeletal muscle. *Physiol. Rep.* 3:e12473. doi: 10.14814/phy2.12473
- Hoang, P. D., Herbert, R. D., and Gandevia, S. C. (2007). Effects of eccentric exercise on passive mechanical properties of human gastrocnemius in vivo. *Med. Sci. Sports. Exerc.* 39, 849–857. doi: 10.1249/MSS.0b013e318033499b
- Huang, J., Qin, K., Tang, C., Zhu, Y., Klein, C. S., Zhang, Z., et al. (2018). Assessment of passive stiffness of medial and lateral heads of gastrocnemius muscle, achilles tendon, and plantar fascia at different ankle and knee positions using the MyotonPRO. *Med. Sci. Monit.* 24, 7570–7576. doi: 10.12659/MSM.909550
- Hyldahl, R. D., and Hubal, M. J. (2014). Lengthening our perspective: morphological, cellular, and molecular responses to eccentric exercise. *Muscle Nerve* 49, 155–170. doi: 10.1002/mus.24077
- Kawczynski, A., Mroczek, D., Andersen, R. E., Stefaniak, T., Arendt-Nielsen, L., and Madeleine, P. (2018). Trapezius viscoelastic properties are heterogeneously affected by eccentric exercise. *J. Sci. Med. Sport* 21, 864–869. doi: 10.1016/j.jsams.2018.01.005
- Kawczynski, A., Nie, H., Jaskolska, A., Jaskolski, A., Arendt-Nielsen, L., and Madeleine, P. (2007). Mechanomyography and electromyography during and after fatiguing shoulder eccentric contractions in males and females. *Scand. J. Med. Sci. Sports* 17, 172–179. doi: 10.1111/j.1600-0838.2006.00551.x
- Kawczynski, A., Samani, A., Fernandez-De-Las-Penas, C., Chmura, J., and Madeleine, P. (2012). Sensory mapping of the upper trapezius muscle in relation to consecutive sessions of eccentric exercise. *J. Strength Cond. Res.* 26, 1577–1583. doi: 10.1519/JSC.0b013e318234e589
- Kelly, J. P., Koppenhaver, S. L., Michener, L. A., Proulx, L., Bisagni, F., and Cleland, J. A. (2018). Characterization of tissue stiffness of the infraspinatus, erector spinae, and gastrocnemius muscle using ultrasound shear wave elastography and superficial mechanical deformation. *J. Electromyogr. Kinesiol.* 38, 73–80. doi: 10.1016/j.jelekin.2017.11.001
- Kinugasa, R., Kawakami, Y., and Fukunaga, T. (2006). Quantitative assessment of skeletal muscle activation using muscle functional MRI. *Magn. Reson. Imaging.* 24, 639–644. doi: 10.1016/j.mri.2006.01.002
- Kjaer, M. (2004). Role of extracellular matrix in adaptation of tendon and skeletal muscle to mechanical loading. *Physiol. Rev.* 84, 649–698. doi: 10.1152/physrev.00031.2003
- Kot, B. C., Zhang, Z. J., Lee, A. W., Leung, V. Y., and Fu, S. N. (2012). Elastic modulus of muscle and tendon with shear wave elastography: variations with different technical settings. *PLoS One* 7:e44348. doi: 10.1371/journal.pone.0044348
- Lacourpaille, L., Hug, F., Bouillard, K., Hogrel, J. Y., and Nordez, A. (2012). Supersonic shear imaging provides a reliable measurement of resting muscle shear elastic modulus. *Physiol. Meas.* 33, N19–N28. doi: 10.1088/0967-3334/33/3/N19
- Lacourpaille, L., Nordez, A., Hug, F., Couturier, A., Dibie, C., and Guilhem, G. (2014). Time-course effect of exercise-induced muscle damage on localized muscle mechanical properties assessed using elastography. *Acta Physiol.* 211, 135–146. doi: 10.1111/apha.12272
- Lacourpaille, L., Nordez, A., Hug, F., Doguet, V., Andrade, R., and Guilhem, G. (2017). Early detection of exercise-induced muscle damage using elastography. *Eur. J. Appl. Physiol.* 117, 2047–2056. doi: 10.1007/s00421-017-3695-9
- Leong, H. T., Hug, F., and Fu, S. N. (2016). Increased upper trapezius muscle stiffness in overhead athletes with rotator cuff tendinopathy. *PLoS One* 11:e0155187. doi: 10.1371/journal.pone.0155187
- Mackey, A. L., Donnelly, A. E., Turpeenniemi-Hujanen, T., and Roper, H. P. (2003). Skeletal muscle collagen content in humans after high-force eccentric contractions. *J. Appl. Physiol.* 97, 197–203. doi: 10.1152/japplphysiol.01174.2003
- Madeleine, P., Hansen, E. A., Andersen, R. E., Kumorek, M., Mroczek, D., Samani, A., et al. (2018). Eccentric exercise induces spatial changes in the mechanomyographic activity of the upper trapezius muscle. *Scand. J. Med. Sci. Sports* 28, 1661–1670. doi: 10.1111/sms.13067
- Madeleine, P., Nie, H., and Arendt-Nielsen, L. (2006). Dynamic shoulder dynamometry: a way to develop delay onset muscle soreness in shoulder muscles. *J. Biomech.* 39, 184–188. doi: 10.1016/j.jbiomech.2004.10.027
- Madeleine, P., Samani, A., Binderup, A. T., and Stensdotter, A. K. (2011). Changes in the spatio-temporal organization of the trapezius muscle activity in response to eccentric contractions. *Scand. J. Med. Sci. Sports* 21, 277–286. doi: 10.1111/j.1600-0838.2009.01037.x
- Nosaka, K., Newton, M., and Sacco, P. (2002). Delayed-onset muscle soreness does not reflect the magnitude of eccentric exercise-induced muscle damage. *Scand. J. Med. Sci. Sports* 12, 337–346. doi: 10.1034/j.1600-0838.2002.10178.x
- Peake, J., Nosaka, K., and Suzuki, K. (2005). Characterization of inflammatory responses to eccentric exercise in humans. *Exer. Immunol. Rev.* 11, 64–85.
- Penailillo, L., Blazevidich, A., Numazawa, H., and Nosaka, K. (2015). Rate of force development as a measure of muscle damage. *Scand. J. Med. Sci. Sports* 25, 417–427. doi: 10.1111/sms.12241
- Picavet, H. S., and Hazes, J. M. (2003). Prevalence of self reported musculoskeletal diseases is high. *Ann. Rheum. Dis.* 62, 644–650. doi: 10.1136/ard.62.7.644
- Proske, U., and Morgan, D. L. (2001). Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical applications. *J. Physiol.* 537(Pt 2), 333–345. doi: 10.1111/j.1469-7793.2001.00333.x
- Pruyn, E. C., Watsford, M. L., Murphy, A. J., Pine, M. J., Spurr, R. W., Cameron, M. L., et al. (2012). Relationship between leg stiffness and lower body injuries in professional Australian football. *J. Sports Sci.* 30, 71–78. doi: 10.1080/02640414.2011.624540
- Schneider, S., Peipsi, A., Stokes, M., Knicker, A., and Abeln, V. (2015). Feasibility of monitoring muscle health in microgravity environments using Myoton technology. *Med. Biol. Eng. Comput.* 53, 57–66. doi: 10.1007/s11517-014-1211-5
- Sohirad, S., Wilson, D., Waugh, C., Finnamore, E., and Scott, A. (2017). Feasibility of using a hand-held device to characterize tendon tissue biomechanics. *PLoS One* 12:e0184463. doi: 10.1371/journal.pone.0184463
- Tas, S., Onur, M. R., Yilmaz, S., Soylu, A. R., and Korkusuz, F. (2017). Shear wave elastography is a reliable and repeatable method for measuring the elastic modulus of the rectus femoris muscle and patellar tendon. *J. Ultrasound. Med.* 36, 565–570. doi: 10.7863/ultra.16.03032
- Vain, A., and Kums, T. (2002). Criteria for preventing overtraining of the musculoskeletal system of gymnasts. *Biol. Sport* 19, 329–345.
- Viir, R., Laiho, K., Kramarenko, J., and Mikkelsen, M. (2011). Repeatability of trapezius muscle tone assessment by a myometric method. *J. Mech. Med. Biol.* 06, 215–228. doi: 10.1142/s0219519406001856
- Xie, Y., Thomas, L., Hug, F., Johnston, V., and Coombes, B. K. (2019). Quantifying cervical and axioapical muscle stiffness using shear wave elastography. *J. Electromyogr. Kinesiol.* 48, 94–102. doi: 10.1016/j.jelekin.2019.06.009
- Xu, J., Fu, S. N., Zhou, D., Huang, C., and Hug, F. (2019). Relationship between pre-exercise muscle stiffness and muscle damage induced by eccentric exercise. *Eur. J. Sport. Sci.* 19, 508–516. doi: 10.1080/17461391.2018.1535625
- Yanagisawa, O., Sakuma, J., Kawakami, Y., Suzuki, K., and Fukubayashi, T. (2015). Effect of exercise-induced muscle damage on muscle hardness evaluated by ultrasound real-time tissue elastography. *Springerplus* 4:308. doi: 10.1186/s40064-015-1094-4
- Zainuddin, Z., Newton, M., Sacco, P., and Nosaka, K. (2005). Effects of massage on delayed-onset muscle soreness, swelling, and recovery of muscle function. *J. Athl. Train.* 40, 174–180.
- Zhang, J., Yu, J., Liu, C., Tang, C., and Zhang, Z. (2019). Modulation in elastic properties of upper trapezius with varying neck angle. *Appl. Bionics. Biomech.* 2019:6048562. doi: 10.1155/2019/6048562

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Kisilewicz, Madeleine, Ignasiak, Cizek, Kawczynski and Larsen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.