



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Long-term outcomes after perioperative treatment with omega-3 fatty acid supplements in colorectal cancer

Sørensen, L Schmidt; Rasmussen, S Ladefoged; Calder, P C; Yilmaz, M Nytoft; Schmidt, E Berg; Thorlacius-Ussing, O

Published in:
BJS Open

DOI (link to publication from Publisher):
[10.1002/bjs5.50295](https://doi.org/10.1002/bjs5.50295)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Sørensen, L. S., Rasmussen, S. L., Calder, P. C., Yilmaz, M. N., Schmidt, E. B., & Thorlacius-Ussing, O. (2020). Long-term outcomes after perioperative treatment with omega-3 fatty acid supplements in colorectal cancer. *BJS Open*, 4(4), 678-684. Advance online publication. <https://doi.org/10.1002/bjs5.50295>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Long-term outcomes after perioperative treatment with omega-3 fatty acid supplements in colorectal cancer

L. Schmidt Sørensen^{1,4,5}, S. Ladefoged Rasmussen^{1,4,5} , P. C. Calder^{6,7}, M. Nytoft Yilmaz², E. Berg Schmidt^{3,5} and O. Thorlacius-Ussing^{1,4,5}

Departments of ¹Gastrointestinal Surgery, ²Oncology and ³Cardiology, and ⁴Clinical Cancer Research Centre, Aalborg University Hospital, and ⁵Department of Clinical Medicine, Aalborg University, Aalborg, Denmark, and ⁶Human Development and Health, Faculty of Medicine, University of Southampton, and ⁷National Institute for Health Research Southampton Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust and University of Southampton, Southampton, UK

Correspondence to: Dr S. Ladefoged Rasmussen, Department of Gastrointestinal Surgery, Aalborg University Hospital, Hobrovej 18-22, 9000 Aalborg, Denmark (e-mail: simon.rasmussen@rn.dk)

Background: This study aimed to evaluate the effect of perioperative supplementation with omega-3 fatty acids (*n*-3 FA) on perioperative outcomes and survival in patients undergoing colorectal cancer surgery.

Methods: Patients scheduled for elective resection of colorectal cancer between 2007 and 2010 were randomized to either an *n*-3 FA-enriched oral nutrition supplement (ONS) twice daily or a standard ONS (control) for 7 days before and after surgery. Outcome measures, including postoperative complications, 3-year cumulative incidence of local or metastatic colorectal cancer recurrence and 5-year overall survival, were compared between the groups.

Results: Of 148 patients enrolled in the study, 125 (65 patients receiving *n*-3 FA-enriched ONS and 60 receiving standard ONS) were analysed. There were no differences in postoperative complications after surgery ($P = 0.544$). The risk of disease recurrence at 3 years was similar (relative risk 1.66, 95 per cent c.i. 0.65 to 4.26). The 5-year survival rate of patients treated with *n*-3 FA was 69.2 (95 per cent c.i. 56.5 to 78.9) per cent, compared with 81.7 (69.3 to 89.4) per cent in the control group ($P = 0.193$). After adjustment for age, stage of disease and adjuvant chemotherapy, *n*-3 FA was associated with higher mortality compared with controls (hazard ratio 1.73, 95 per cent c.i. 1.06 to 2.83; $P = 0.029$). The interaction between *n*-3 FA and adjuvant chemotherapy was not statistically significant.

Conclusion: Perioperative supplementation with *n*-3 FA did not confer a survival benefit in patients undergoing colorectal cancer surgery. *n*-3 FA did not benefit the subgroup of patients treated with adjuvant chemotherapy or decrease the risk of disease recurrence.

Funding information

Obel Family Foundation
Aase and Ejnar Danielsens Foundation
Augustinus Foundation
L. F. Foghts Foundation

Paper accepted 30 March 2020

Published online 11 May 2020 in Wiley Online Library (www.bjsopen.com). DOI: 10.1002/bjs5.50295

Introduction

Colorectal cancer surgery has been associated with a phase of hyperinflammation followed by a relative immune incompetence^{1,2}. Patients' nutritional status and, in particular, the availability of specific biologically active nutrients, such as the marine omega-3 fatty acids (*n*-3 FA) (eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA)) and docosapentaenoic acid, have been investigated in

the literature^{3–7} as factors influencing the postoperative course. In particular, *n*-3 FA have been associated with an improved immunoproliferative response⁸.

Several colorectal cancer cell line studies have investigated also whether *n*-3 FA could improve the response to chemotherapeutic agents. A previous study⁹ demonstrated a synergistic anticancer effect between EPA and a regimen of 5-fluorouracil (5-FU) and oxaliplatin *in vitro*

and *in vivo* against the human colonic cancer cells. In line with this, other researchers¹⁰ hypothesized that EPA could have a role in adjuvant therapy for the prevention of metastatic colorectal cancer. Oral administration of EPA reduced the growth of experimental colorectal cancer liver metastases and, importantly, preoperative treatment with EPA resulted in improved disease-free survival¹⁰. Finally, it has been proposed¹¹ that *n*-3 FA might increase the susceptibility of tumour cells to chemotherapeutic regimens: in an experimental study conducted on mice with colonic cancer, the administration of 5-FU and *n*-3 FA increased both DNA damage and the apoptotic index by activation of extrinsic and intrinsic apoptotic pathways. The increased proapoptotic effect by the synergism of 5-FU and *n*-3 FA could be attributed to the incorporation of *n*-3 FA into the cancer cell membranes, altering membrane fluidity and thereby chemosensitizing the tumour cells.

The aim of the present study was to investigate whether oral supplementation with *n*-3 FA in the perioperative period might be associated with improved postoperative outcomes (decreased number of complications), decreased risk of disease recurrence, and improved survival in patients undergoing colorectal cancer surgery.

Methods

A randomized, double-blind, placebo-controlled, single-centre trial was conducted to study the effect of an *n*-3 FA-enriched oral nutrition supplement (ONS) on long-term outcomes in patients scheduled for colorectal cancer surgery.

Patients scheduled for colorectal cancer surgery at the Department of Gastrointestinal Surgery, Aalborg University Hospital, between July 2007 and January 2010 were screened for eligibility. Aalborg University Hospital treats some 260–350 patients with colorectal cancer each year. Each patient was referred from the surgical outpatient clinic and managed according to the current Danish guidelines. The inclusion and exclusion criteria have been described previously¹² (ClinicalTrials identifier NCT00488904).

Briefly, patients in both the *n*-3 FA and control group received the ONS as a sip feed (200 ml twice daily, once in the morning and once in the afternoon) for 7 days before and 7 days after surgery. The feeds were isocaloric (1.5 kcal/ml) and isonitrogenous (Supportan®; Fresenius Kabi, Bad Homburg, Germany). Both feeds contained the same amounts of carbohydrate, protein, total fat and *n*-6 FA, as well as vitamins and minerals. However, the fatty acid composition differed between the two feeds: both

contained medium-chain triglycerides, sunflower oil and safflower oil, but the *n*-3 FA ONS also contained fish oil. Thus, patients in the *n*-3 FA group received 2.0 g EPA and 1.0 g DHA per day from the ONS, whereas no *n*-3 FA was provided to controls.

Outcome measures


Outcomes included perioperative results (pneumonia, wound infection, urinary tract infection, peritonitis (including anastomotic leakage) and sepsis) and survival. Five-year overall survival (defined as being alive 5 years after colorectal cancer surgery) and the risk of disease recurrence (defined as the cumulative incidence of local or metastatic colorectal cancer recurrence) at 3 years according to treatment allocation were evaluated. According to the Danish national guidelines for colorectal cancer treatment, each patient was followed with CT of the thorax and abdomen 1 and 3 years after treatment. If the patient did not have full colonoscopy before surgery, a new colonoscopy was planned for 3 months after surgery; otherwise, all patients had colonoscopy 5 years after treatment. Data on patient survival and disease recurrence were collected from patient records and the Danish National Patient Registry.

After evaluation of the surgical resection specimen, the patients were restaged and evaluated at a new multidisciplinary team conference, after which those with high-risk stage II and III disease were referred for adjuvant chemotherapy. Patients with stage IV disease with no surgically treatable liver metastasis received chemotherapy with palliative intent only. A subgroup analysis of all patients receiving adjuvant chemotherapy with curative intent was also performed, and therefore excluded patients who had either neoadjuvant radiochemotherapy or postoperative palliative chemotherapy (patients with stage IV disease).

Post-trial analysis of survival was approved by the North Denmark Committee on Health Research Ethics Committee (N-20140064) and the Danish Data Protection Agency.

Statistical analysis

The study was designed to detect a 20 per cent difference in postoperative complication rates between groups (a reduction from 30 to 10 per cent was hypothesized). The required sample size was 72 patients in each group (80 per cent power, $\alpha = 0.05$). Baseline characteristics were compared using Student's *t* test or Fisher's exact test, as appropriate. The median length of hospital stay was



FA, fatty acids.

compared using the Wilcoxon–Mann–Whitney rank test. The Kaplan–Meier estimator was used to depict the association between treatment with the n -3 FA ONS and overall survival. Overall 5-year survival was calculated from Kaplan–Meier estimates and compared with the log rank test. To evaluate the effects of a supplement with n -3 FA along with interaction with adjuvant chemotherapy, a generalized linear model was constructed using the pseudo-observation method¹³. Estimates were presented as hazard ratios (HRs) with 95 per cent confidence intervals. The risk of disease recurrence was estimated using competing risk regression. The Aalen–Johansen estimator was used to produce the cumulative incidence plot, and the pseudo-observation method was employed to determine the relative risk (RR) of disease recurrence 3 years after surgery. All analyses were conducted using STATA® V.15.1 (StataCorp, College Station, Texas, USA).

Results

Of 610 patients who had colorectal cancer surgery during the study period, 201 did not meet the inclusion criteria, 31 declined to participate, and 230 could not be included because they had surgery within 4–5 days of diagnosis and therefore could not receive the allocated 7-day intervention before surgery (Fig. 1). The remaining 148 patients were randomized (in line with the power calculation); however, 23 patients were subsequently excluded owing to patient refusal, postoperative death, logistical reasons, or because they did not have a standard colorectal cancer resection. Accordingly, 65 patients were enrolled in the n -3 FA treatment arm and 60 patients in the control arm.

Baseline characteristics of the patients are shown in Table 1. Twenty-two patients received adjuvant chemotherapy, either as monotherapy with 5-FU (5 patients) or as combined treatment (17) with folinic acid, 5-FU and oxaliplatin (FOLFOX).

Table 1 Study population			
	Enriched ONS (n = 65)	Control ONS (n = 60)	P†
Age (years)*	68.3(11.3)	70.6(10.1)	0.246‡
Sex ratio (M : F)	27 : 38	30 : 30	0.373
Smoking status			
Non-smoker	47 (72)	51 (85)	
Smoker	17 (26)	8 (13)	
Unknown	1 (1.5)	1 (2)	0.148
Cancer location			
Colon	36 (55)	34 (57)	
Rectum	29 (45)	26 (43)	1.000
UICC stage			
I	14 (22)	10 (17)	
II	26 (40)	29 (48)	
III	18 (28)	19 (32)	
IV	7 (11)	2 (3)	0.348
Adjuvant therapy	11 (17)	11 (18)	1.000

Values in parentheses are percentages unless indicated otherwise; *values are mean(s.d.). ONS, oral nutrition supplement. †Fisher's exact test, except ‡Student's *t* test.


Perioperative outcomes

Data on perioperative outcomes have been reported elsewhere¹². In brief, there was no significant difference in the number of infectious (35.4 per cent in patients treated with *n*-3 FA versus 38.3 per cent in the control group; $P = 0.853$) or non-infectious (29.2 and 23.3 per cent respectively; $P = 0.544$) complications, 30-day mortality rate (0 per cent in both groups; $P = 1.000$) or median length of hospital stay (9 versus 10 days respectively; $P = 0.807$).

Survival

The 5-year overall survival rate was 69.2 (95 per cent c.i. 56.5 to 78.9) per cent in patients treated with *n*-3 FA, compared with 81.7 (69.3 to 89.4) per cent in the control group ($P = 0.193$). The Kaplan–Meier survival curves (Fig. 2) indicate that the assumption of proportional hazards was violated over the recorded time period (which was also evident from the log–log plot). Therefore, the pseudo-observation method was used to construct a generalized linear model to evaluate the effect of treatment with *n*-3 FA and adjuvant chemotherapy on 5-year survival. The model was adjusted for age and stage of disease at the time of diagnosis. The unadjusted HR 5 years after surgery was 1.68 (95 per cent c.i. 0.88 to 3.21), suggesting no effect of *n*-3 FA treatment on survival of surgically treated patients with colorectal cancer (Table 2). After adjustment for adjuvant chemotherapy, age and stage at time of diagnosis, supplementation with *n*-3 FA was associated

Fig. 2 Kaplan–Meier analysis of overall survival according to treatment allocation




n-3 FA, omega-3 fatty acids. $P = 0.193$ (log rank test).

with a higher overall mortality (HR 1.73 (1.06 to 2.83); $P = 0.029$).

The interaction between treatment with *n*-3 FA and adjuvant chemotherapy was, however, not statistically significant, suggesting a spurious association. Of patients treated with adjuvant chemotherapy, the stratified HR for patients in the *n*-3 FA group was 2.09 (95 per cent c.i. 0.92

	Univariable estimates		Multivariable estimates	
	Hazard ratio	P	Hazard ratio	P
Age	1.05 (1.01, 1.09)	0.011	1.07 (1.03, 1.12)	0.001
<i>n</i> -3 fatty acids	1.68 (0.88, 3.21)	0.118	1.73 (1.06, 2.83)	0.029
Adjuvant chemotherapy	2.23 (1.23, 4.06)	0.009	1.52 (0.87, 2.65)	0.138
UICC stage				
II	1.75 (0.40, 7.66)	0.461	3.89 (0.43, 34.71)	0.223
III	5.19 (1.30, 20.69)	0.020	12.57 (1.66, 95.37)	0.014
IV	6.67 (1.55, 28.58)	0.011	7.87 (1.03, 60.82)	0.047

Values in parentheses are 95 per cent confidence intervals. *n*-3 FA, omega-3 fatty acids.



There were 55 patients in the omega-3 fatty acids (*n*-3 FA) group and 55 in the control group. Relative risk 1.66 (95 per cent c.i. 0.65 to 4.26).

to 4.75) compared with 1.29 (0.51 to 3.25) in the control group ($P = 0.138$).

Disease recurrence

The effect of treatment with *n*-3 FA on the risk of disease recurrence, using death as a competing risk, was calculated for the 110 patients who had an R0 resection (55 in each group) (Fig. 3). The risk of disease recurrence did not differ between groups (RR 1.66, 95 per cent c.i. 0.65 to 4.26), and adjustment for stage and adjuvant chemotherapy did not alter the estimate.

Discussion

This study aimed to evaluate perioperative treatment with *n*-3 FA on the outcomes of patients undergoing colorectal

cancer surgery in a randomized clinical setting. Of note, the investigation was designed as an RCT, enrolling blinded participants; the sip feeds were nutritionally similar and had an identical appearance and taste, apart from content of marine *n*-3 FA. Compliance with the allocated intervention (*n*-3 FA versus control) was acceptable, and the study population was relatively homogeneous¹².

However, the study was powered to show a difference in postoperative complications and not a difference in survival in patients receiving chemotherapy and *n*-3 FA; thus, the number of patients might have been too low for this outcome. The low number of patients receiving adjuvant chemotherapy (11 in each group) also limited the study power, and any association with reduced survival or other events should be interpreted with caution. Follow-up on postoperative dietary habits, other disease states and medications was not conducted, and this should also be regarded as a limitation. The intake of long-chain *n*-3 FA in Western countries is generally low, but is traditionally higher in Scandinavian countries, and therefore the results may not be directly comparable for other populations. Finally, other *n*-3 FA dosages and different lengths of treatment duration were not considered. Prolonged treatment might be needed to ensure an effect on postoperative outcomes.

Some meta-analyses^{14,15} identified that so-called immunonutrition, containing arginine, glutamine, nucleotides and *n*-3 FA, could reduce postoperative infections in selected patient groups. However, it is unclear which of the components is responsible for the clinical effect. In the present study, daily supplementation with 3 g EPA plus DHA for 7 days before and 7 days after colorectal cancer surgery did lead to increased incorporation of these *n*-3 FA in the cell membranes, as well as a significant increase in the production of leukotriene B5 from EPA and a significant decrease in the production of leukotriene B4, but had no effect on postoperative complications^{12,16,17}.

It has been suggested¹⁸ that supplementation with *n*-3 FA could increase chemotherapeutic efficacy and thereby

contribute to increased survival. In a combined analysis of the Nurses' Health Study and the Health Professionals Follow-up Study, it was reported¹⁸ that a high intake of *n*-3 FA after colorectal cancer diagnosis was associated with a lower risk of mortality from the disease. Thus, patients consuming at least 0.3 g *n*-3 FA daily had an adjusted HR for colorectal cancer-specific mortality of 0.59 (95 per cent c.i. 0.35 to 1.01). Patients who increased their marine *n*-3 FA intake by at least 0.15 g/day after diagnosis had a HR of 0.30 for colorectal cancer death, compared with those who did not change their intake of marine *n*-3 FA.

A previous phase II double-blind placebo-controlled RCT enrolled patients with colorectal cancer and liver metastases who received 2 g EPA daily or placebo twice daily from randomization to the day of liver resection¹⁰. The *n*-3 FA EPA was successfully incorporated into liver metastases removed at operation, and a trend towards reduced tumour vascularization in *n*-3 FA-treated colorectal liver metastases was recorded. The authors report a non-significant trend towards improved overall survival from *n*-3 FA in the first 18 months after surgery, and early colorectal cancer recurrence was similar in the groups¹⁰.

A meta-analysis¹⁹ compared RCTs using pharmaconutrition with those using standard nutrition in elective adult surgical patients operated on between 1980 and 2011. No differences were reported in postoperative mortality, but there was a significant reduction in infectious complications and length of hospital stay. Regarding perioperative administration, there was a reduction in anastomotic dehiscence. Furthermore, a reduction in non-infectious complications was detected with postoperative administration. Pharmaconutrition given before surgery did not demonstrate a notable advantage compared with standard nutrition for any of the assessed clinical outcomes¹⁹. The importance of timing was also shown in another review²⁰ of published meta-analyses, implying improved outcome after oncological gastrointestinal surgery, but before issuing recommendations further studies are warranted. Additionally, two meta-analyses^{19,21} showed that preoperative pharmaconutrition did not confer any benefit over standard formulations. The earlier seen benefits of pharmaconutrition (reduction in infectious complications) were reported only with perioperative or postoperative administration.

Acknowledgements

The authors thank study nurse A. Madsen and the staff at the Lipid Research Clinic at Aalborg University Hospital for invaluable assistance in completing the study, and H. C. Brix Noergaard for helping with the initial phase of the

study. Fresenius Kabi (Bad Homburg, Germany) supplied sip feeds.

This study was funded by the Obel Family Foundation, Aase and Ejnar Danielsen's Foundation, Augustinus Foundation and L. F. Foghts Foundation.

Disclosure: The authors declare no conflict of interest.

References

- 1 Kehlet H, Dahl JB. Anaesthesia, surgery, and challenges in postoperative recovery. *Lancet* 2003; **362**: 1921–1928.
- 2 Calder PC. *n*-3 fatty acids, inflammation, and immunity – Relevance to postsurgical and critically ill patients. *Lipids* 2004; **39**: 1147–1161.
- 3 Bozzetti F, Braga M, Gianotti L, Gavazzi C, Mariani L. Postoperative enteral *versus* parenteral nutrition in malnourished patients with gastrointestinal cancer: a randomised multicentre trial. *Lancet* 2001; **358**: 1487–1492.
- 4 Braga M, Gianotti L, Vignali A, Di Carlo V. Preoperative oral arginine and *n*-3 fatty acid supplementation improves the immunometabolic host response and outcome after colorectal resection for cancer. *Surgery* 2002; **132**: 805–814.
- 5 Zhu MW, Tang DN, Hou J, Wei JM, Hua B, Sun JH *et al.* Impact of fish oil enriched total parenteral nutrition on elderly patients after colorectal cancer surgery. *Chin Med J (Engl)* 2012; **125**: 178–181.
- 6 Jie B, Jiang ZM, Nolan MT, Zhu SN, Yu K, Kondrup J. Impact of preoperative nutritional support on clinical outcome in abdominal surgical patients at nutritional risk. *Nutrition* 2012; **28**: 1022–1027.
- 7 Ryan AM, Reynolds JV, Healy L, Byrne M, Moore J, Brannelly N *et al.* Enteral nutrition enriched with eicosapentaenoic acid (EPA) preserves lean body mass following esophageal cancer surgery: results of a double-blinded randomized controlled trial. *Ann Surg* 2009; **249**: 355–363.
- 8 Marimuthu K, Varadhan KK, Ljungqvist O, Lobo DN. A meta-analysis of the effect of combinations of immune modulating nutrients on outcome in patients undergoing major open gastrointestinal surgery. *Ann Surg* 2012; **255**: 1060–1068.
- 9 Vasudevan A, Yu Y, Banerjee S, Woods J, Farhana L, Rajendra SG *et al.* Omega-3 fatty acid is a potential preventive agent for recurrent colon cancer. *Cancer Prev Res* 2014; **7**: 1138–1148.
- 10 Cockbain AJ, Volpato M, Race AD, Munarini A, Fazio C, Belluzzi A *et al.* Anticorectal cancer activity of the omega-3 polyunsaturated fatty acid eicosapentaenoic acid. *Gut* 2014; **63**: 1760–1768.
- 11 Rani I, Sharma B, Kumar S, Kaur S, Agnihotri N. Apoptosis mediated chemosensitization of tumor cells to 5-fluorouracil on supplementation of fish oil in experimental colon carcinoma. *Tumor Biol* 2017; **39**: 101042831769501.
- 12 Sorensen LS, Thorlacius-Ussing O, Schmidt EB, Rasmussen HH, Lundbye-Christensen S, Calder PC *et al.* Randomized clinical trial of perioperative omega-3 fatty acid supplements

- in elective colorectal cancer surgery. *Br J Surg* 2014; **101**: 33–42.
- 13 Hansen SN, Andersen PK, Parner ET. Events per variable for risk differences and relative risks using pseudo-observations. *Lifetime Data Anal* 2014; **20**: 584–598.
- 14 Cerantola Y, Hübner M, Grass F, Demartines N, Schäfer M. Immunonutrition in gastrointestinal surgery. *Br J Surg* 2011; **98**: 37–48.
- 15 Heys SD, Walker LG, Smith I, Eremin O. Enteral nutritional supplementation with key nutrients in patients with critical illness and cancer: a meta-analysis of randomized controlled clinical trials. *Ann Surg* 1999; **229**: 467–477.
- 16 Sorensen LS, Rasmussen HH, Aardestrup IV, Thorlacius-Ussing O, Lindorff-Larsen K, Schmidt EB *et al*. Rapid incorporation of ω -3 fatty acids into colonic tissue after oral supplementation in patients with colorectal cancer: a randomized, placebo-controlled intervention trial. *J Parenter Enteral Nutr* 2014; **38**: 617–624.
- 17 Sorensen L, Thorlacius-Ussing O, Rasmussen HH, Lundbye-Christensen S, Calder PC, Lindorff-Larsen K *et al*. Effects of perioperative supplementation with omega-3 fatty acids on leukotriene B4 and leukotriene B5 production by stimulated neutrophils in patients with colorectal cancer: a randomized, placebo-controlled intervention trial. *Nutrients* 2014; **6**: 4043–4057.
- 18 Song M, Zhang X, Meyerhardt JA, Giovannucci EL, Ogino S, Fuchs CS *et al*. Marine ω -3 polyunsaturated fatty acid intake and survival after colorectal cancer diagnosis. *Gut* 2017; **66**: 1790–1796.
- 19 Osland E, Hossain MB, Khan S, Memon MA. Effect of timing of pharmaconutrition (immunonutrition) administration on outcomes of elective surgery for gastrointestinal malignancies: a systematic review and meta-analysis. *J Parenter Enteral Nutr* 2014; **38**: 53–69.
- 20 Osland E, Memon B, Memon MA. Pharmaconutrition administration on outcomes of elective oncological surgery for gastrointestinal malignancies: is timing everything? – a review of published meta-analyses until the end of 2016. *Transl Gastroenterol Hepatol* 2018; **3**: 52–52.
- 21 Song G-M, Tian X, Zhang L, Ou Y-X, Yi L-J, Shuai T *et al*. Immunonutrition support for patients undergoing surgery for gastrointestinal malignancy. *Medicine (Baltimore)* 2015; **94**: e1225.