



AALBORG UNIVERSITY
DENMARK

Aalborg Universitet

Green Protein from locally grown crops

Steenfeldt, Sanna; Ambye-Jensen, Morten; Lübeck, Mette

Creative Commons License
CC BY 4.0

Publication date:
2020

Document Version
Også kaldet Forlagets PDF

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Steenfeldt, S., Ambye-Jensen, M., & Lübeck, M. (2020). Green Protein from locally grown crops. <https://orgprints.org/37034/1/PA013-Steenfeldt-2019-green-protein-final.pdf>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Green protein from locally grown crops

Problem

Organic poultry producers are often confronted with high feed costs and a lack of alternative high-quality protein sources for poultry.

Solution

Green protein concentrate can be produced from locally grown crops such as clover-grass (see figure 1) or alfalfa. It can be used in the diets of organic broilers and layers. In a bio-refinery, protein concentrate is obtained by pressing fresh green material (see figure 2), heating/fermenting the juice to precipitate protein and finally putting it in a centrifuge. The concentrated green protein can be dried and added to poultry feed.

Benefits

Concentrate from clover/clover grass and alfalfa has a high protein content and an optimal amino acid profile for poultry, which makes the feed formulation of organic diets more optimal. An increase in locally grown protein sources can improve the sustainability of the farm and make the farmer less dependent upon imported protein, such as soya, from overseas.

Practical recommendation

- Choose an appropriate type of green crop, such as clover-grass or alfalfa, with an expected high protein and amino acid content. Consider soil types and weather patterns to grow a crop with a good and high quality yield.
- Harvest the field at regular intervals in order to achieve good plant growth and to obtain batches with more high quality protein and less fibre



Figure 1: Harvesting of locally grown clover-grass. Photo: Erik Fog, SEGES



Figure 2: Screw pressing of fresh clover grass into green juice and press cake. Photo: Erik Fog, SEGES

Applicability box

Theme

Layers; feeding, processing and handling of harvested feed

Geographical coverage

In temperate climates. Middle and Northern Europe.

Application time

The product can be used at any time of the year if the protein paste are dried and stored under optimal conditions.

Required time

Harvest time of green protein during spring, summer and autumn, processing time in a bio-refinery and time to dry it.

Period of impact

Immediate Impact

Equipment

Machinery required for harvest of green material (clover/grass/ alfalfa) and for transportation to a bio-refinery plant or to storage facilities.

Best in

Choice of crop for production of green protein depends on the country, soil type and weather conditions during preferred harvest time. Advantageous in crop rotation.

- Harvesting procedures, which minimise soil content in the green material obtained from the field are necessary to obtain good quality green protein and to avoid wear of machinery and technical equipment
- Cooperation with a bio-refinery plant is a prerequisite in order to concentrate the protein into a green paste that can be dried and used in poultry feed.
- If not dried, the wet green paste can be stored in closed containers/plastic bags in cool conditions for a shorter period.
- Chemical analysis of the green protein concentrate is important in order to replace other protein sources such as soya and to carry out the correct feed formulation. This can be done together with advisors or feed companies.

Further information

Further readings

Video

- Video "[GRASS PROTEIN - a golden chance to improve organic farming](#)" from Seges

Links

- Report on "[Green Biomass – Protein Production Through Bio-refining](#)"
- OrganoFinery: [Organic growth with biorefined organic protein feed, fertilizer and energy](#)
- Check the [Organic Farm Knowledge](#) platform for more practical recommendations.

About this practice abstract and OK-Net EcoFeed

Publishers:

Dept. of Animal Science (ANIS) and Dept. of Engineering (AU)
Aarhus University (AU), Blichers Allé 20, Postboks 50, DK-8830 Tjele
Phone +45 8715 6000/+45 8715 8074, anis.au.dk, eng.au.dk

Section for Sustainable Biotechnology, Dept. of Chemistry and Bioscience
Aalborg University (AAU), A. C. Meyers Vænge 15, DK-2450 Copenhagen
Phone +45 2447 0418, bio.aau.dk

Research Institute of Organic Agriculture (FiBL)
Ackerstrasse 113, Postfach 219, CH-5070 Frick
Phone +41 62 865 72 72, info.suisse@fibl.org, www.fibl.org

IFOAM EU, Rue du Commerce 124, BE-1000 Brussels
Phone +32 2 280 12 23, info@ifoam-eu.org, www.ifoam-eu.org

Author: Sanna Steinfeldt, Morten Ambye-Jensen, Mette Lübeck

Contact: sanna.steenfeldt@anis.au.dk, maj@eng.au.dk, mel@bio.aau.dk

Permalink: organic-farmknowledge.org/tool/37034



OK-Net EcoFeed: This practice abstract was elaborated in the Organic Knowledge Network on Monogastric Animal Feed project. The project is running from January 2018 to December 2020. The overall aim of OK-Net EcoFeed is to help farmers, breeders and the organic feed processing industry in achieving the goal of 100% use of organic and regional feed for monogastrics.

Project website: ok-net-ecofeed.eu

Project partners: IFOAM EU Group (project coordinator), BE; Aarhus University (ICROFS), DK; Organic Research Centre (ORC), UK; Institut Technique de l'Agriculture Biologique (ITAB), FR; Research Institute of Organic Agriculture (FiBL), CH; Bioland, DE; Associazione Italiana per l'Agricoltura Biologica (AIAB), IT; Donau Soja DS, AT; Swedish University of Agricultural Sciences, SE; ECOVALIA, ES; Soil Association, UK.

© 2020

