

## Home parenteral nutrition provision modalities for chronic intestinal failure in adult patients

*An international survey*

Pironi, Loris; Steiger, Ezra; Brandt, Chrisoffer; Joly, Francisca; Wanten, Geert; Chambrier, Cecile; Aimasso, Umberto; Sasdelli, Anna Simona; Zeraschi, Sarah; Kelly, Darlene; Szczepanek, Kinga; Jukes, Amelia; Di Caro, Simona; Theilla, Miriam; Kunecki, Marek; Daniels, Joanne; Serlie, Mireille; Poullenot, Florian; Wu, Jian; Cooper, Sheldon C; Rasmussen, Henrik H; Compher, Charlene; Seguy, David; Crivelli, Adriana; Pagano, Maria C; Hughes, Sarah-Jane; Guglielmi, Francesco W; Kozjek, Nada Rotovnik; Schneider, Stéphane M; Gillanders, Lyn; Ellegard, Lars; Thibault, Ronan; Matras, Przemysław; Zmarzly, Anna; Matysiak, Konrad; Van Gossum, André; Forbes, Alastair; Wyer, Nicola; Taus, Marina; Virgili, Nuria M; O'Callaghan, Margie; Chapman, Brooke; Osland, Emma; Cuerda, Cristina; Sahin, Peter; Jones, Lynn; Won Lee, Andre Dong; Masconale, Luisa; Orlandoni, Paolo; Izbéki, Ferenc; Spaggiari, Corrado; Bueno, Marta; Dotchinova-Simeonova, Maryana; Garde, Carmen; Serralde-Zúñiga, Aurora E.; Oliveira, Gabriel; Murilo, Zelijko; Czako, Laszlo; Kekstas, Gintautas; Sanz-Paris, Alejandro; Jáuregui, Estrella Petrina; Murillo, Ana Zugasti; Schafer, Eszter; Arends, Jann; Suárez-Llanos, José P.; Lal, Simon; Home Artificial Nutrition and Chronic Intestinal Failure Special Interest Group of ESPEN

*Published in:*  
Clinical Nutrition

*DOI (link to publication from Publisher):*  
[10.1016/j.clnu.2019.03.010](https://doi.org/10.1016/j.clnu.2019.03.010)

*Creative Commons License*  
CC BY-NC-ND 4.0

*Publication date:*  
2020

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Pironi, L., Steiger, E., Brandt, C., Joly, F., Wanten, G., Chambrier, C., Aimasso, U., Sasdelli, A. S., Zeraschi, S., Kelly, D., Szczepanek, K., Jukes, A., Di Caro, S., Theilla, M., Kunecki, M., Daniels, J., Serlie, M., Poullenot, F., Wu, J., ... Home Artificial Nutrition and Chronic Intestinal Failure Special Interest Group of ESPEN (2020). Home parenteral nutrition provision modalities for chronic intestinal failure in adult patients: An international survey. *Clinical Nutrition*, 39(2), 585-591. <https://doi.org/10.1016/j.clnu.2019.03.010>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

#### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from [vbn.aau.dk](http://vbn.aau.dk) on: July 04, 2025

# Home parenteral nutrition provision modalities for chronic intestinal failure in adult patients: An international survey

[Loris Pironi](#)<sup>1</sup>, [Ezra Steiger](#)<sup>2</sup>, [Chrisoffer Brandt](#)<sup>3</sup>, [Francisca Joly](#)<sup>4</sup>, [Geert Wanten](#)<sup>5</sup>, [Cecile Chambrier](#)<sup>6</sup>, [Umberto Aimasso](#)<sup>7</sup>, [Anna Simona Sasdelli](#)<sup>8</sup>, [Sarah Zeraschi](#)<sup>9</sup>, [Darlene Kelly](#)<sup>10</sup>, [Kinga Szczepanek](#)<sup>11</sup>, [Amelia Jukes](#)<sup>12</sup>, [Simona Di Caro](#)<sup>13</sup>, [Miriam Theilla](#)<sup>14</sup>, [Marek Kunecki](#)<sup>15</sup>, [Joanne Daniels](#)<sup>16</sup>, [Mireille Serlie](#)<sup>17</sup>, [Florian Poullenot](#)<sup>18</sup>, [Jian Wu](#)<sup>19</sup>, [Sheldon C Cooper](#)<sup>20</sup>, [Henrik H Rasmussen](#)<sup>21</sup>, [Charlene Compheer](#)<sup>22</sup>, [David Seguy](#)<sup>23</sup>, [Adriana Crivelli](#)<sup>24</sup>, [Maria C Pagano](#)<sup>25</sup>, [Sarah-Jane Hughes](#)<sup>26</sup>, [Francesco W Guglielmi](#)<sup>27</sup>, [Nada Rotovnik Kozjek](#)<sup>28</sup>, [Stéphane M Schneider](#)<sup>29</sup>, [Lyn Gillanders](#)<sup>30</sup>, [Lars Ellegard](#)<sup>31</sup>, [Ronan Thibault](#)<sup>32</sup>, [Przemysław Matras](#)<sup>33</sup>, [Anna Zmarzly](#)<sup>34</sup>, [Konrad Matysiak](#)<sup>35</sup>, [Andr  Van Gossum](#)<sup>36</sup>, [Alastair Forbes](#)<sup>37</sup>, [Nicola Wyr](#)<sup>38</sup>, [Marina Taus](#)<sup>39</sup>, [Nuria M Virgili](#)<sup>40</sup>, [Margie O'Callaghan](#)<sup>41</sup>, [Brooke Chapman](#)<sup>42</sup>, [Emma Osland](#)<sup>43</sup>, [Cristina Cuerda](#)<sup>44</sup>, [Peter Sahin](#)<sup>45</sup>, [Lynn Jones](#)<sup>46</sup>, [Andre Dong Won Lee](#)<sup>47</sup>, [Luisa Masconale](#)<sup>48</sup>, [Paolo Orlandoni](#)<sup>49</sup>, [Ferenc Izb ki](#)<sup>50</sup>, [Corrado Spaggiari](#)<sup>51</sup>, [Marta Bueno](#)<sup>52</sup>, [Maryana Doitchinova-Simeonova](#)<sup>53</sup>, [Carmen Garde](#)<sup>54</sup>, [Aurora E Serralde-Z niga](#)<sup>55</sup>, [Gabriel Olveira](#)<sup>56</sup>, [Zeljko Krznaric](#)<sup>57</sup>, [Laszlo Czako](#)<sup>58</sup>, [Gintautas Kekstas](#)<sup>59</sup>, [Alejandro Sanz-Paris](#)<sup>60</sup>, [Estrella Petrina J uregui](#)<sup>61</sup>, [Ana Zugasti Murillo](#)<sup>62</sup>, [Eszter Schafer](#)<sup>63</sup>, [Jann Arends](#)<sup>64</sup>, [Jos  P Su rez-Llanos](#)<sup>65</sup>, [Simon Lal](#)<sup>66</sup>, [Home Artificial Nutrition and Chronic Intestinal Failure Special Interest Group of ESPEN](#); [European Society for Clinical Nutrition and Metabolism](#)

<sup>1</sup> St. Orsola University Hospital, Bologna, Italy. Electronic address: [loris.pironi@unibo.it](mailto:loris.pironi@unibo.it).

<sup>2</sup> Cleveland Clinic Foundation, Cleveland, OH, USA.

<sup>3</sup> Rigshospitalet, Copenhagen, Denmark.

<sup>4</sup> Beaujon Hospital, Clichy, France.

<sup>5</sup> Radboud University Medical Center, Nijmegen, the Netherlands.

<sup>6</sup> Hospices Civils de Lyon, Centre Hospitalier Lyon Sud, Lyon, France.

<sup>7</sup> Citt  della Salute e della Scienza, Torino, Italy.

<sup>8</sup> St. Orsola University Hospital, Bologna, Italy.

<sup>9</sup> Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom.

<sup>10</sup> Mayo Clinic College of Medicine, Rochester, MN, USA.

<sup>11</sup> Stanley Dudrick's Memorial Hospital, Skawina, Poland.

<sup>12</sup> University Hospital of Wales, Cardiff, United Kingdom.

<sup>13</sup> University College Hospital, London, United Kingdom.

<sup>14</sup> Rabin Medical Center, Petach Tikva, Israel.

<sup>15</sup> M. Pirogow Hospital, Lodz, Poland.

<sup>16</sup> Nottingham University Hospital NHS Trust, Nottingham, United Kingdom.

<sup>17</sup> Academic Medical Center, Amsterdam, the Netherlands.

<sup>18</sup> CHU de Bordeaux, H pital Haut-L v que, Pessac, France.

<sup>19</sup> University Hospital Southampton NHS Foundation Trust, Southampton, United Kingdom.

<sup>20</sup> University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom.

<sup>21</sup> Center for Nutrition and Bowel Disease, Aalborg University Hospital, Aalborg, Denmark.

<sup>22</sup> Hospital of the University of Pennsylvania, Philadelphia, PA, USA.

<sup>23</sup> CHRU de Lille, Lille, France.

<sup>24</sup> Hospital Universitario Fundacion Favaloro, Buenos Aires, Argentina.

<sup>25</sup> Federico II University, Napoli, Italy.

- <sup>26</sup> Regional Intestinal Failure Service, Belfast Health and Social Care Trust, Northern Ireland, United Kingdom.
- <sup>27</sup> San Nicola Pellegrino Hospital, Trani, Italy.
- <sup>28</sup> Institute of Oncology, Ljubljana, Slovenia.
- <sup>29</sup> CHU Archet, Nice, France.
- <sup>30</sup> Auckland City Hospital, Auckland, New Zealand.
- <sup>31</sup> Sahlgrenska University Hospital, Gothenburg, Sweden.
- <sup>32</sup> Nutrition unit, CHU Rennes, Nutrition Metabolisms and Cancer Institute, NuMeCan, INRA, INSERM, Université Rennes, Rennes, France.
- <sup>33</sup> Medical University of Lublin, Lublin, Poland.
- <sup>34</sup> J. Gromkowski City Hospital, Wrocław, Poland.
- <sup>35</sup> H.Święcicki University Hospital, Poznań, Poland.
- <sup>36</sup> Hôpital Erasme, Brussels, Belgium.
- <sup>37</sup> Norfolk and Norwich University Hospital, Norwich, United Kingdom.
- <sup>38</sup> University Hospital, Coventry, United Kingdom.
- <sup>39</sup> Ospedali Riuniti, Ancona, Italy.
- <sup>40</sup> Hospital Universitari de Bellvitge, Barcelona, Spain.
- <sup>41</sup> Flinders Medical Centre, Adelaide, Australia.
- <sup>42</sup> Austin Health, Melbourne, Argentina.
- <sup>43</sup> Royal Brisbane and Women's Hospital, Herston, Australia.
- <sup>44</sup> Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- <sup>45</sup> St. Imre Hospital, Budapest, Hungary.
- <sup>46</sup> Royal Prince Alfred Hospital, Sydney, Australia.
- <sup>47</sup> Hospital das Clinicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil.
- <sup>48</sup> ULSS 22 Ospedale Orlandi, Bussolengo, Italy.
- <sup>49</sup> IRCCS - INRCA, Ancona, Italy.
- <sup>50</sup> Szent György Teaching Hospital of County Fejér, Székesfehérvár, Hungary.
- <sup>51</sup> AUSL di Parma, Parma, Italy.
- <sup>52</sup> Hospital Universitario Arnau de Vilanova, Lleida, Spain.
- <sup>53</sup> Bulgarian Executive Agency of Transplantation, Sofia, Bulgaria.
- <sup>54</sup> Hospital Universitario Donostia, San Sebastian, Spain.
- <sup>55</sup> Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, México City, Mexico.
- <sup>56</sup> IBIMA, Hospital Regional Universitario de Málaga, Universidad de Málaga, Málaga, Spain.
- <sup>57</sup> University Hospital Centre Zagreb, Zagreb, Croatia.
- <sup>58</sup> University of Szeged, Szeged, Hungary.
- <sup>59</sup> Vilnius University Hospital Santariskiu Clinics, Vilnius, Lithuania.
- <sup>60</sup> Miguel Servet Hospital, Zaragoza, Spain.
- <sup>61</sup> Complejo Hospitalario de Navarra, Pamplona, Spain.
- <sup>62</sup> Hospital Virgen del Camino, Pamplona, Spain.
- <sup>63</sup> Magyar Honvedseg Egészségügyi Központ (MHEK), Budapest, Hungary.
- <sup>64</sup> Tumor Biology Center, Freiburg, Germany.
- <sup>65</sup> Hospital Universitario Nuestra Señora de Candelaria, Santa Cruz de Tenerife, Spain.
- <sup>66</sup> Salford Royal NHS Foundation Trust, Salford, United Kingdom.

## **HOME PARENTERAL NUTRITION PROVISION MODALITIES FOR CHRONIC INTESTINAL FAILURE IN ADULT PATIENTS: AN INTERNATIONAL SURVEY**

Loris Pironi, Denise Konrad, Chrisoffer Brandt, Francisca Joly, Geert Wanten, Cecile Chambrier, Umberto Aimasso, Anna Simona Sasdelli, Sarah Zeraschi, Darlene Kelly, Kinga Szczepanek, Amelia Jukes, Simona Di Caro, Miriam Theilla, Marek Kunecki, Joanne Daniels, Mireille Serlie, Florian Poullenot, Jian Wu, Sheldon C. Cooper, Henrik H. Rasmussen, Charlene Compher, David Seguy, Adriana Crivelli, Maria C. Pagano, Sarah-Jane Hughes, Francesco W. Guglielmi, Nada Rotovnik Kozjek, Stéphane M. Schneider, Lyn Gillanders, Lars Ellegard, Ronan Thibault, Przemysław Matras, Anna Zmarzly, Konrad Matysiak, André Van Gossum, Alastair Forbes, Nicola Wyer, Marina Taus, Nuria M. Virgili, Margie O'Callaghan, Brooke Chapman, Emma Osland, Cristina Cuerda, Peter Sahin, Lynn Jones, Andre Dong Won Lee, Valentino Bertasi, Paolo Orlandoni, Ferenc Izbéki, Corrado Spaggiari, Marta Bueno Díez, Maryana Doitchinova-Simeonova, Carmen Garde, Aurora E. Serralde-Zúñiga, Gabriel Oliveira, Zeljko Krznaric, Laszlo Czako, Gintautas Kekstas, Alejandro Sanz-Paris, Estrella Petrina Jáuregui, Ana Zugasti Murillo, Eszter Schafer, Jann Arends, José P. Suárez-Llanos and Simon Lal.

The Home Artificial Nutrition and Chronic Intestinal Failure special interest group of ESPEN, the European Society for Clinical Nutrition and Metabolism

Corresponding author

Loris Pironi  
Center for Chronic Intestinal Failure  
Department of Digestive System  
St. Orsola-Malpighi Hospital, University of Bologna  
Via Massarenti, 9 - 40138 Bologna, Italy  
Tel: +39 - 051 - 6363073  
Fax: +39 - 051 - 6364193  
Email: loris.pironi@unibo.it

## Abstract

**Background and aim.** The safety and effectiveness of an HPN program depends on both the expertise and the management procedures of the HPN center. We aimed to know the modalities needed to provide the home parenteral nutrition (HPN)-program and the types of intravenous supplementation (IVS)-admixtures supplied to patients with chronic intestinal failure (CIF) in different countries.

**Methods.** In March 2015, 65 centers from 22 countries enrolled 3239 patients (benign disease 90.1%, malignant disease 9.9%), recording the patient, CIF and HPN characteristics in a structured database. The HPN-provider was categorized as health care system local pharmacy (LP) or home care company (HCC). The IVS-admixture was categorized as fluids and electrolytes alone (FE) or parenteral nutrition, either commercially premixed (PA) or customized to the individual patient (CA), alone or plus extra FE (PAFE or CAFE).

**Results.** HPN-provider: HCC 66%, LP 34%; no difference between benign-CIF and malignant-CIF. LP was the main modality in 11 Countries; HCC prevailed in 4 European countries, Israel, USA, South America and Oceania ( $p < 0.001$ ). IVS-admixture: FE 10%, PA 17%, PAFE 17%, CA 38%, CAFE 18%. PA+PAFE use was greater in malignant-CIF and CA+CAFE use was greater in benign-CIF ( $p < 0.001$ ). PA+PAFE prevailed in those Countries where LP was the main HPN-provider and CA+CAFE prevailed where the main HPN-provider was HCC ( $p < 0.001$ ).

**Conclusions.** The HPN provision and the IVS-admixture types differ greatly among countries, among HPN centers and between benign-CIF and malignant-CIF. As both HPN provider and IVS-admixture types may play a role in the safety and effectiveness of HPN therapy, criteria to homogenize HPN programs are needed, to give patients the same opportunity to receive appropriate HPN therapy.

**Keywords:** Intestinal failure, home parenteral nutrition, management, intravenous supplementation, cancer.

## Introduction

Home parenteral nutrition (HPN) is the primary and life-saving treatment for patients with chronic intestinal failure (CIF) [1]. Intestinal failure (IF) is defined as the “reduction of gut function below the minimum necessary for the absorption of macronutrients and/or water and electrolytes, such that intravenous supplementation (IVS) is required to maintain health and/or growth” [2]. Chronic IF can be due to five pathophysiological mechanisms (short bowel, intestinal fistulas, intestinal dysmotility, intestinal mechanical occlusion, or extensive small bowel mucosa disease) which can originate from either non-malignant (benign-CIF) or malignant (malignant-CIF) diseases [2]. Patients with CIF require IVS for months, years or lifelong [1,2]. They are discharged onto HPN programs which aim to provide evidence-based therapy, to prevent HPN-related complications, such as central venous catheter (CVC)-related infections and metabolic complications, and to maximize the patient/family quality of life (QoL) [3,4]. The European Society for Clinical Nutrition and metabolism (ESPEN) guidelines on benign-CIF recommend that at discharge: patients are metabolically stable, able physically and emotionally to cope with the HPN therapy, and have an adequate home environment; patients should be cared for by a multidisciplinary team with skills and experience in IF and HPN management; patient/caregiver training for HPN management should be patient-centered with a multidisciplinary approach, together with written guidelines; HPN patients should have access to infusion pumps or devices with specified safety features together with ancillary products, safe compounding and delivery systems. Thus, the safety and effectiveness of a HPN program depends on the expertise and the management modalities of the HPN center.

It is known that the management and the provision of HPN programs differ greatly among countries and among HPN centers. However, only one study, performed in 2010, objectively described this feature [5]. Using the ESPEN database for CIF, we have carried out an international

cross-sectional survey to know the modalities used to provide the HPN-program and the types of IVS-admixtures supplied to patients with CIF.

Accepted Manuscript



## Materials and methods

This international cross-sectional observational study comprises the largest study protocol approved by the Home Artificial Nutrition and Chronic Intestinal Failure (HAN&CIF) special interest group of ESPEN, which aims to investigate the applicability of the clinical classification of CIF [6]. The HPN centers' recruitment, the patient inclusion criteria, the modalities of data collection and the recorded items have been already extensively described [6] and are summarized below.

### *Participating centers and patient enrollment criteria*

Sixty-five HPN centers from 22 countries enrolled all adult patients ( $\geq 18$  year old) who were on HPN for either benign-CIF or malignant-CIF on March 1st 2015. The term malignant-CIF indicates the presence of an active malignant disease at time of enrollment on the study (and thus excludes, for example, patients with IF as a result of radiotherapy in whom the malignancy has been cured; these patients were surveyed within the benign-CIF group).

### *Data collection and schedule*

Data were collected into a structured questionnaire embedded in an Excel (Microsoft Co., 2013) database, termed "the CIF Action day", available at the web page of the HAN&CIF group on the ESPEN website [7].

Demographic, clinical, CIF, underlying disease, IVS and HPN program characteristics were gathered, and the clinical classification of CIF was calculated for each patient [6]. The HPN-provider was categorized as health care system local pharmacy (LP) or home care company (HCC). By HPN-provider was meant the supplier to the patient of the IVS-admixture, the infusion pump or regulatory device and the ancillaries for infusion-line management and CVC medication. The IVS-admixture was categorized as: fluids and electrolytes (FE); commercially premixed ready-to-use parenteral nutrition admixture (PA); commercially premixed parenteral nutrition admixture plus extra FE (PAFE);

parenteral nutrition admixture customized (tailored) to the individual patient requirements (CA); parenteral nutrition admixture customized to the individual patient requirements alone or plus extra FE (CAFE).

#### *Ethical statement*

The research was based on anonymized information taken from patient records at the time of data collection. The study was conducted with full regard to confidentiality of the individual patient. Ethical committee approval was obtained by the individual HPN centers according to local regulations. The collected data were used only for the study purpose. The identity of the contributing centers has also been anonymized for data analysis and presentation.

#### *Statistical analysis*

The daily mean volume and energy of IVS were calculated as follows: daily total volume (mL/day) or energy (kcal/day) = amount per day of infusion x number of infusions per week / 7; daily volume or energy per kg of patient body weight (mL/kgBW/day or kcal/kgBW/day) = amount per day of infusion x number of infusions per week) / 7 / kg patient body weight. The patients' body mass index (BMI) was calculated by Quetelet's formula (weight (kg) / height (m<sup>2</sup>).

Data are reported as mean  $\pm$  standard deviation (SD) and as absolute and relative frequencies. The non-parametric Kruskal Wallis test, the Fisher's exact test and the Chi-square test were applied where appropriate.

The IBM SSPS Statistics package for Windows, version 23.0 (BM Co., Armonk, NY, USA) was used for the analyses. Two-tailed p values less than 0.05 were considered as statistically significant.

## Results

### *Participating centers and patient cohorts*

A total of 3239 patients were included, 2919 benign-CIF (90.1%) and 320 malignant-CIF (9.9%) (**Table 1**). All the HPN centers enrolled benign-CIF patients, while only 45 of them enrolled malignant-CIF. The malignant-CIF cohort had statistically significant older age, lower BMI, shorter duration of HPN, IVS of greater daily volume and energy, and a 10 times greater percentage of patients with IF due to mechanical occlusions, (**Table 2**). In both benign-CIF and malignant-CIF, two-thirds of patients were female.

In the benign-CIF cohort, the underlying diseases were Crohn's disease 22.4%, mesenteric ischemia 17.7%, surgical complications 15.8%, primary chronic intestinal pseudo-obstruction 9.7%, radiation enteritis 7.3%, others (<3% each-one) 21.3%, and not reported 5.9%. In the malignant-CIF cohort, the type of cancer was not specified 62%, gastrointestinal 28%, extra-abdominal 10%. Concurrent enteritis due to radio- or chemo-therapy was described in 5% of cases, and peritoneal carcinomatosis was reported in 12%.

### *HPN-providers and IVS-admixture types in the total group*

The HPN-provider was not reported in 11 cases and was LP in 1111 (34.4%) and HCC in 2117 (65.6%) cases. The IVS-admixture type was FE in 312 (9.7%), PA in 556 (17.2%) and PAFE in 541 (16.8%), CA in 1227 (37.9%) and CAFE in 595 (18.4%) cases. The IVS-admixture types significantly differed between the two modalities of HPN provision. When the HPN was provided by a HCC, the IVS-admixtures were CA or CAFE in two-thirds of cases, while PA or PAFE accounted for more than 50% of the IVS-admixtures provided by the LPs (**Table 3**).

#### *HPN-providers and IVS-admixture types by countries*

HCCs provided all HPN in the UK and Israel, were almost exclusive providers ( $\geq 80\%$  of patients) in the USA, Mexico and South America, and were the main providers (56-63% of cases) in France, Italy, Poland and Oceania. LPs provided all the HPN programs in Denmark, two thirds of programs in The Netherlands and more than 90% of cases in the other 9 European countries which contributed to the survey (**Figure 1**).

In those countries, excepting Poland, where most or all the HPN programs were provided by a HCC, CA and CAFE represented more than 50% of the IVS-admixtures. Where the LP was the main HPN-provider, PA and PAFE prescription prevailed (**Figure 2**).

#### *HPN-providers and IVS-admixture types by the nature of the underlying disease*

The percentage split of the two HPN-providers did not differ between benign-CIF and malignant-CIF, while CA and CAFE were the IVS-admixture types in almost two-thirds of benign-CIF and PA and PAFE were the IVS-admixture types in more than 50% of malignant-CIF (**Figure 3**).

## Discussion

This international survey shows that the modalities of HPN provision and the type of IVS-admixture supplied significantly differ among countries and between benign-CIF and malignant-CIF. Although it is well known that HPN management is not homogeneous among countries or among HPN centers within an individual country, this is the first study that gives objective data on this aspect. The strengths of the study are the large numbers of participating countries, and the worldwide distribution of contributing HPN-centers and enrolled patients. The main limitation of the study is the relatively small number of malignant-CIF patients, in comparison with those with benign-CIF (**Table 1**). Actually, the percentage of enrolled patients with malignant-CIF is lower than that expected on the basis of previous early and recent surveys on HPN [8-13]. This could be due to the voluntary basis of the HPN center participation, that could have attracted primarily those centers mainly caring for benign-CIF. Indeed, it may be that in individual countries, most of the patients on HPN for cancer are managed by specialists other than those caring for CIF, such as oncologists or internists. Another explanation could be that, in previous surveys on HPN prevalence, a percentage of patients with a diagnosis of cancer may have not had CIF, being on HPN because of refusal of an otherwise working gut, or just because they already had a CVC positioned for chemotherapy. However, as expected, the benign-CIF and the malignant-CIF cohorts of the present study consistently differed for all the clinical and IVS characteristics, thus supporting the representativeness of the malignant-CIF cohort (**Table 2**).

Overall, the prescribed IVS-admixture type and the HPN-providers looked associated. Commercially PA or PAFE were more frequently used when the LP was the HPN-provider, while CA or CAFE were more frequently used when the HPN-provider was a HCC (**Table 3**). This would indicate that, when required, a HCC makes it easier to supply a tailored IVS-admixture. Indeed, as CIF is a rare

condition, not all the LP may have developed the expertise and/or implemented the facilities to produce CA in a sufficient quantity to overcome the production costs.

The data confirm that the HPN provision modalities differ greatly among the individual countries, with a range of 0 to 100% of cases for both HCC and LP (**Figure 1**). The association between the HPN provision modality and the IVS-admixture types reported in the total cohort, was observed also in the individual countries (**Figure 2**). The primary aims of an HPN program are prevention of HPN-related complications and maximization of the patient/family QoL [3]. The protocol for patient/caregiver training and the facilities and ancillaries for IVS management may be very relevant to the CVC-related complications, and the availability of a portable infusion pump may significantly change the QoL of patients [3]. Differences between means of HPN provision may therefore have implications to the safety and efficacy of an HPN program. This suggests that criteria for the implementation of HPN provision should be formally devised, in order to homogenize this feature of the HPN program and to give patients the same opportunity to receive appropriate HPN therapy regardless of where they live.

The results showed that IVS-admixture types but not the HPN-provider differed between benign-CIF and malignant-CIF. The IVS-admixtures tailored to the patient requirements (CA and CAFE) were mainly used in benign-CIF, while premixed, ready-to-use, IVS-admixture (PA and PAFE) were mainly used in malignant-CIF (**Figure 3**). This difference may be due to the characteristics of the two patient populations, in terms of pathophysiological mechanisms of IF as well as in the aims of the HPN program and the expected patient outcome. The clinical scenarios of benign-CIF and malignant-CIF are quite different. In malignant-CIF, the cause of IF was more homogeneous, being represented by mechanical obstruction in almost 50% of cases, often due to peritoneal carcinomatosis. In benign-CIF, the mechanisms of IF were represented by SBS and fistula in almost 70% of patients. The oral food

and beverage intakes and the intestinal fluid and electrolytes losses may greatly differ among patients [6]. Patients with benign CIF have a high survival probability, may have good probabilities of intestinal rehabilitation and programs aim for social-working-familial rehabilitation and good QoL. Most of the patients are independent from a caregiver and do not need any home healthcare assistance [3]. This scenario often needs a fine tuning of the HPN program, requiring tailored IVS-admixture in order to maximize the prevention of long-term metabolic complications as well as the daily time free of IVS infusion. In patients with malignant-CIF, HPN may be required while waiting for or during cancer-directed treatment, on symptomatic treatment, and/or receiving palliative care [4]. The expected duration of HPN is much shorter, either because of a transient need related to cancer treatment plans or to short life expectancy of advanced cancer. These patients are often home-bound, dependent on a caregiver and require home healthcare assistance. Safety of the line infusion is a priority of any HPN program, while the risk of metabolic complications may be lower in the short term. In this scenario, premixed ready-use IVS-admixtures may adequately fit with the requirements of the patient with malignant-CIF. The lack of differences in HPN-provider between benign-CIF and malignant-CIF was probably due to the bias in the enrollment of the HPN centers, that were mostly devoted to benign-CIF, so that the same HPN provider may have been used for the few patients with malignant-CIF enrolled.

In conclusion, the HPN provision modalities and the IVS-admixture types differ greatly among countries and between benign-CIF and malignant-CIF. As both HPN provider and IVS-admixture types may play a role in the safety and effectiveness of a HPN program, these data indicate the need to devise criteria to homogenize these essential features of HPN, in order to give all patients with CIF the same opportunity to receive this life-saving therapy in the appropriate modality.

## **Acknowledgements**

### **Contributing coordinators and centers by country**

#### **Argentina**

Adriana N. Crivelli, Hector Solar Muñiz; Hospital Universitario Fundacion Favaloro, Buenos Aires

#### **Australia**

Brooke R. Chapman; Austin Health, Melbourne

Lynn Jones; Royal Prince Alfred Hospital, Camperdown

Margie O'Callaghan; Flinders Medical Centre, Adelaide

Emma Osland, Ruth Hodgson, Siobhan Wallin, Kay Lasenby; Royal Brisbane and Women's Hospital, Herston

#### **Belgium**

Andre Van Gossum; Hôpital Erasme, Brussels

#### **Brasil:**

Andre Dong Won Lee; Hospital da Universidade de São Paulo, São Paulo

#### **Bulgaria**

Maryana Doitchinova-Simeonova; Bulgarian Executive Agency of Transplantation, Sofia

#### **Croatia**

Zeljko Krznaric; University Hospital Centre Zagreb, Zagreb

#### **Denmark**

Henrik Højgaard Rasmussen; Aalborg University Hospital, Aalborg

Chrisoffer Brandt, Michael Staun; Rigshospitalet, Copenhagen

#### **France**

Cecile Chambrier; Hospices Civils de Lyon, Hôpital de la Croix Rousse, Lyon

Francisca Joly; Beaujon Hospital, Clichy

Florian Poullenot; CHU de Bordeaux, Hôpital Haut-Lévêque, Pessac

Stéphane M. Schneider; CHU Archet, Nice

David Seguy; CHRU de Lille, Lille

Ronan Thibault; University Hospital of Rennes, Rennes

#### **Germany**

Jann Arends; Tumor Biology Center, Freiburg

#### **Hungary**

Laszlo Czako, Tomas Molnar, Mihaly Zsilak-Urban; University of Szeged, Szeged

Ferenc Izbéki; Szent György Teaching Hospital of County Fejér, Székesfehérvár

Peter Sahin, Gábor Udvarhelyi; St. Imre Hospital, Budapest

Eszter Schafer; Magyar Honvedseg Egészségügyi Központ (MHEK), Budapest

#### **Israel**

Miriam Theilla; Rabin Medical Center, Petach Tikva

#### **Italy**

Anna Simona Sasdelli, Loris Pironi; S. Orsola University Hospital, Bologna

Umberto Aimasso; Città della Salute e della Scienza, Torino

Valentino Bertasi, Luisa Mosconale; ULSS 22 Ospedale Orlandi, Bussolengo

Francesco W. Guglielmi, Nunzia Regano; San Nicola Pellegrino Hospital, Trani

Paolo Orlandoni; INRCA – IRCCS, Ancona



Maria C. Pagano, Santarpia Lidia, Lucia Alfonsi; Federico II University, Italy  
Corrado Spaggiari; AUSL di Parma, Parma  
Marina Taus, Debora Busni; Ospedali Riuniti, Ancona

#### **Lithuania**

Gintautas Kekstas; Vilnius University Hospital Santariskiu Clinics, Vilnius

#### **México**

Aurora E. Serralde-Zúñiga; Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, México City

#### **New Zealand**

Lyn Gillanders; Auckland City Hospital, Auckland

#### **Poland**

Marek Kunecki; M. Pirogow Hospital, Lodz  
Przemysław Matras; Medical University of Lublin, Lublin  
Konrad Matysiak; H.Święcicki University Hospital, Poznań  
Kinga Szczepanek; Stanley Dudrick's Memorial Hospital, Skawina  
Anna Zmarzly; J. Gromkowski City Hospital, Wrocław

#### **Slovenia**

Nada Rotovnik Kozjek; Institute of Oncology, Ljubljana

#### **Spain**

Marta Bueno Díez; Hospital Universitario Arnau de Vilanova, Lleida  
Cristina Cuerda; Hospital General Universitario Gregorio Marañón, Madrid  
Carmen Garde; Hospital Universitario Donostia, San Sebastian  
Nuria M. Virgili; Hospital Universitari de Bellvitge, Barcelona  
Gabriel Oliveira; Hospital Regional Universitario de Málaga, Málaga  
M<sup>a</sup> Estrella Petrina Jáuregui; Complejo Hospitalario de Navarra, Pamplona  
Alejandro Sanz-Paris; Miguel Servet Hospital, Zaragoza  
José P. Suárez-Llanos; Hospital Universitario Nuestra Señora de Candelaria, Santa Cruz de Tenerife  
Ana Zugasti Murillo; Hospital Virgen del Camino, Pamplona

#### **Sweden**

Lars Ellegard; Sahlgrenska University Hospital, Gothenburg

#### **The Netherlands**

Mireille Serlie, Cora Jonker; Academic Medical Center, Amsterdam  
Geert Wanten; Radboud University Medical Center, Nijmegen

#### **United Kingdom**

Sheldon C. Cooper; University Hospitals Birmingham NHS Foundation Trust, Birmingham  
Joanne Daniels; Nottingham University Hospital NHS Trust, Nottingham  
Simona Di Caro, Niamh Keane, Pinal Patel; University College Hospital, London  
Alastair Forbes; Norfolk and Norwich University Hospital, Norwich  
Sarah-Jane Hughes; Belfast Health and Social Care Trust, Belfast  
Amelia Jukes, Rachel Lloyd; University Hospital of Wales, Cardiff  
Simon Lal, Arun Abraham, Gerda Garside, Michael Taylor; Salford Royal NHS Foundation Trust, Salford  
Jian Wu, Trevor Smith, Charlotte Pither, Michael Stroud; University Hospital Southampton NHS Foundation Trust, Southampton  
Nicola Wyer, Reena Parmar, Nicola Burch; University Hospital, Coventry  
Sarah Zeraschi; Leeds Teaching Hospitals NHS Trust, Leeds

**United States of America**

Charlene Compher; Hospital of the University of Pennsylvania, Philadelphia, PA

Darlene Kelly; Mayo Clinic College of Medicine, Rochester, MN

Denise Konrad, Ezra Steiger; Cleveland Clinic Foundation, Cleveland, OH

**Statistical analysis performed by:** Dr. Marianna Mastroroberto, MD, PhD; Department of Medical and Surgical Sciences, University of Bologna, Italy

**Funding source**

The project of the ESPEN database for Chronic Intestinal Failure was promoted by the ESPEN

Executive Committee in 2013, was approved by the ESPEN Council and was supported by an ESPEN grant.

**Statement of authorship**

LP devised the study protocol, collected the data, analyzed the results and drafted the manuscript.

The Home Artificial Nutrition & Chronic Intestinal Failure Special Interest Group of ESPEN discussed and approved the protocol study, discussed the results and reviewed the manuscript before submission. According to the authorship rules described in the protocol study, all the coordinators of the participating centers were considered coauthors of the study and received the manuscript upon submission. All authors approved the final version of the manuscript before submission.

**Conflict of interest statements**

None declared.

## References

1. Staun M, Pironi L, Bozzetti F, Baxter J, Forbes A, Joly F, Jeppesen P, Moreno J, Hébuterne X, Pertkiewicz M, Mühlebach S, Shenkin A, Van Gossum A; ESPEN. ESPEN Guidelines on Parenteral Nutrition: home parenteral nutrition (HPN) in adult patients. Clin Nutr. 2009 Aug;28(4):467-79. doi: 10.1016/j.clnu.2009.04.001. Epub 2009 May 22. PubMed PMID: 19464089.
2. Pironi L, Arends J, Baxter J, Bozzetti F, Peláez RB, Cuerda C, Forbes A, Gabe S, Gillanders L, Holst M, Jeppesen PB, Joly F, Kelly D, Klek S, Irtun Ø, Olde Damink SW, Panisic M, Rasmussen HH, Staun M, Szczepanek K, Van Gossum A, Wanten G, Schneider SM, Shaffer J; Home Artificial Nutrition & Chronic Intestinal Failure; Acute Intestinal Failure Special Interest Groups of ESPEN. ESPEN endorsed recommendations. Definition and classification of intestinal failure in adults. Clin Nutr. 2015 Apr;34(2):171-80.
3. Pironi L, Arends J, Bozzetti F, Cuerda C, Gillanders L, Jeppesen PB, Joly F, Kelly D, Lal S, Staun M, Szczepanek K, Van Gossum A, Wanten G, Schneider SM; Home Artificial Nutrition & Chronic Intestinal Failure Special Interest Group of ESPEN.. ESPEN guidelines on chronic intestinal failure in adults. Clin Nutr. 2016 Apr;35(2):247-307. doi:10.1016/j.clnu.2016.01.020. PubMed PMID: 26944585.
4. Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, Bozzetti F, Fearon K, Hütterer E, Isenring E, Kaasa S, Krznaric Z, Laird B, Larsson M, Laviano A, Mühlebach S, Muscaritoli M, Oldervoll L, Ravasco P, Solheim T, Strasser F, de van der Schueren M, Preiser JC. ESPEN guidelines on nutrition in cancer patients. Clin Nutr. 2017 Feb;36(1):11-48. doi: 10.1016/j.clnu.2016.07.015. Epub 2016 Aug 6. PubMed PMID: 27637832.
5. Baxter JP, Gillanders L, Angstmann K, Staun M, O'Hanlon C, Smith T, Joly F, Thul P, Jonkers C, Wanten G, Gardiner K, Klek S, Cuerda C, Magambo W, Hawthorne AB, Lukes A, Van Gossum A,

Theilla M, Singer P, Shamir R, Pironi L. Home parenteral nutrition: An international benchmarking exercise. *e-SPEN Journal* 7 (2012) e211-e214.

6. Pironi, L., Konrad, D., Brandt, C., Joly, F., Wanten, G., Agostini, F., Chambrier, C., Aimasso, U., Zeraschi, S., Kelly, D., Szczepanek, K., Jukes, A., Di Caro, S., Theilla, M., Kunecki, M., Daniels, J., Serlie, M., Poullenot, F., Wu, J., Cooper, S.C., Rasmussen, H.H., Compher, C., Seguy, D., Crivelli, A., Pagano, M.C., Hughes, S.J., Guglielmi, F.W., Kozjek, N.R., Schneider, S.M., Gillanders, L., Ellegard, L., Thibault, R., Matras, P., Zmarzly, A., Matysiak, K., Van Gossum, A., Forbes, A., Wyer, N., Taus, M., Virgili, N.M., O'Callaghan, M., Chapman, B., Osland, E., Cuerda, C., Sahin, P., Jones, L., Lee, A.D.W., Bertasi, V., Orlandoni, P., Izbéki, F., Spaggiari, C., Díez, M.B., Doitchinova-Simeonova, M., Garde, C., Serralde-Zúñiga, A.E., Oliveira, G., Krznaric, Z., Czako, L., Kekstas, G., Sanz-Paris, A., Jáuregui, E.P., Murillo, A.Z., Schafer, E., Arends, J., Suárez-Llanos, J.P., Shaffer, J., Lal, S. 2017, Clinical classification of adult patients with chronic intestinal failure due to benign disease: an international multicenter cross-sectional survey. *Clin Nutr.* 2018 Apr;37(2):728-738. doi: 10.1016/j.clnu.2017.04.013. Epub 2017 Apr 19.
7. <http://www.espen.org/education/special-interest>
8. Howard L, Ament M, Fleming CR, Shike M, Steiger E. Current use and clinical outcome of home parenteral and enteral nutrition therapies in the United States. *Gastroenterology*. 1995;109:355-65
9. Van Gossum, A., Bakker, A, De Francesco, A., Ladefoged, K., Leon-Sanz, M., Messing, M., Pironi, L., Pertkiewicz, M., Shaffer, J., Thul, P., Wood, S. Home parenteral nutrition at home in adults: a multicentre survey in Europe in 1993. *Clinical Nutrition* 1996; 15, 53-59.

10. Pironi L and SINPE regional co-ordinators. Development of home artificial nutrition in Italy over a seven year period: 2005–2012. BMC Nutrition. December 2017, 3:63. <https://link.springer.com/article/10.1186/s40795-016-0118-y>.
11. British Association of Parenteral and Enteral Nutrition (BANS) Report 2016, Artificial Nutrition Support in the UK 2005-2015. Adult Home Parenteral Nutrition & Home Intravenous Fluids. <http://www.bapen.org.uk/>.
12. Brandt CF, Hvistendahl M, Naimi RM, Tribler S, Staun M, Brobech P, et al. Home Parenteral Nutrition in Adult Patients With Chronic Intestinal Failure: The Evolution Over 4 Decades in a Tertiary Referral Center. JPEN Journal of parenteral and enteral nutrition. 2017;41:1178-87. PMID: 28483328-
13. Wanden-Berghe Lozano C, Virgili Casas N, Ramos Boluda E, Cuerda Compés C, Moreno Villares JM, Pereira Cunill JL, Gómez Candela C, Burgos Peláez R, Penacho Lázaro MÁ, Pérez de la Cruz A, Álvarez Hernández J, Gonzalo Marín M, Matía Martín P, Martínez Faedo C, Sánchez Martos EÁ, Sanz Paris A, Campos Martín C, Martín Folgueras T, Martín Palmero MÁ, Martín Fontalba MLÁ, Luengo Pérez LM, Zugasti Murillo A, Martínez Ramírez MJ, Carabaña Pérez F, Martínez Costa C, Díaz Guardiola P, Tejera Pérez C, Parés Marimón RM, Irlés Rocamora JA, Garde Orbaiz C, Ponce González MÁ, García Zafra MV, Sánchez Sánchez R, Urgeles Planella JR, Apezetxea Celaya A, Sánchez-Vilar Burdiel O, Joaquín Ortiz C, Suárez Llanos JP, Pintor de la Maza B, Leyes García P, Gil Martínez MC, Mauri Roca S, Carrera Santaliestra MJ. [Home and Ambulatory Artificial Nutrition (NADYA) Group Report -Home parenteral nutrition in Spain, 2016]. Nutr Hosp. 2017 Nov24;34(5):1497-1501. doi: 10.20960/nh.1686. Spanish. PubMed PMID: 29280669.

**Table 1.** Patients on home parenteral nutrition for chronic intestinal failure (CIF) due to non-malignant (benign) or malignant disease, enrolled by countries contributing in the survey.

	<b>Total n.</b>	<b>Benign-CIF n. (%)</b>	<b>Malignant-CIF n. (%)</b>
UK	781	738 (94.5)	43 (5.5)
France	478	441 (92.3)	37 (7.7)
Italy	362	326 (90.1)	36 (9.9)
Poland	283	224 (79.2)	59 (20.8)
Denmark	262	233 (88.9)	29 (11.1)
The Netherlands	257	229 (89.1)	28 (10.9)
Spain	43	40 (93.0)	3 (7.0)
Slovenia	39	31 (79.5)	8 (20.5)
Sweden	25	24 (96.0)	1 (4.0)
Hungary	22	20 (90.9)	2 (9.2)
Belgium	21	21 (100)	0
Germany	10	1 (10)	9 (90)
Bulgaria	5	4 (80)	1 (20)
Croatia	3	3 (100)	0
Lithuania	3	2 (66.7)	1 (33.3)
USA	429	389 (90.7)	40 (9.3)
Israel	90	71 (78.9)	19 (21.1)
Mexico	4	3 (75)	1 (25)
Argentina	44	44 (100)	0
Brasil	7	7 (100)	0
Australia	44	41 (93.2)	3 (6.8)
New Zealand	27	27 (100)	0
<b>Total</b>	<b>3239</b>	<b>2919 (90.1)</b>	<b>320 (9.9)</b>

**Table 2.** Characteristics of the cohorts of patients with chronic intestinal failure (CIF) enrolled in the study: patients without malignant disease (Benign-CIF), n. 2919; patients with a malignant disease (Malignant-CIF), n.320.

	Benign-CIF	Malignant-CIF	P
Gender			0.202
Males	36.8%	39.45	
Females	63.2%	60.6%	
Age, years	54.9±16.0	60.6±13.5	<0.001
BMI, kg/m <sup>2</sup>	22.2±4.4	21.5±4.4	0.002
HPN duration, months	58.1±71.5	17.1±30.9	<0.001
Pathophysiological mechanism of IF			<0.001
SBS-J	38.6%	28.8%	
SBS-JC	19.9%	9.7%	
SBS-JIC	5.9%	2.8%	
Fistulas	7.0%	3.4%	
Dysmotility	17.5%	3.4%	
Mechanical Obstruction	4.4%	45.9%	
Mucosal Disease	6.8%	5.9%	
IVS volume, mL/day	1877.0±1016.6	1967.6±817.8	0.004
IVS energy, kcal/day	1088.0±649.4	1315.9±560.9	<0.001
Clinical classification of CIF (IVS, mL/day)			<0.001
FE1, ≤1000	5.8%	3.1%	
FE2, 1001-2000	2.2%	1.3%	
FE2, 2001-3000	0.5%	0	
FE4, >3000	0.3%	0	
PN1, ≤1000	15.9%	10.3%	
PN2, 1001-2000	40.9%	47.2%	
PN3, 2001-3000	23.1%	29.4%	
PN4, >3000	11.3%	8.8%	

BMI, body mass index; HPN, home parenteral nutrition; SBS-J, short bowel syndrome with end jejunostomy; SBS-JC, short bowel syndrome with jejunocolon anastomosis; SBS-JIC, short bowel syndrome with jejunocolon anastomosis and total colon; IVS, intravenous supplementation; FE, fluid and electrolytes; PN, parenteral nutrition

**Table 3.** Intravenous supplementation (IVS)-admixture type by home parenteral nutrition (HPN)-provider in patients with chronic intestinal failure (P<0.001)

	<b>Total</b> n.	<b>FE</b> n. (%)	<b>PA</b> n. (%)	<b>PAFE</b> n. (%)	<b>CA</b> n. (%)	<b>CAFE</b> n. (%)	<b>P</b>
<b>HCC</b>	2117	224 (10.5)	149 (7.0)	304 (14.3)	906 (42.8)	534 (25.2)	<0.001
<b>LP</b>	1111	88 (7.9)	407 (36.6)	237 (21.3)	318 (28.6)	61 (5.4)	

HCC, home care company

LP, health care system local pharmacy

FE, fluids and electrolytes

PA, commercially premixed ready-to-use parenteral nutrition admixture


PAFE, commercially premixed ready-to-use parenteral nutrition admixture plus extra fluids and electrolytes

CA, parenteral nutrition admixture customized (tailored) to the individual patient requirements

CAFE, parenteral nutrition admixture customized (tailored) to the individual patient requirements plus extra fluids and electrolytes




**Figure 1.** Home parenteral nutrition (HPN)-providers by countries in patients with chronic intestinal failure. **LP**, health care system local pharmacy. **HCC**, home care company. (P<0.001)



**Europe others:** Belgium, Bulgaria, Croatia, Germany, Hungary, Lithuania, Slovenia, Spain, Sweden. **Oceania:** Australia, New Zealand  
**S. America:** Argentina, Brasil

**Figure 2.** Intravenous supplementation (IVS)-admixture type by countries in patients with chronic intestinal failure. **FE**, fluids and electrolytes; **PA**, commercially premixed ready-to-use parenteral nutrition admixture; **PAFE**, commercially premixed ready-to-use parenteral nutrition admixture plus extra fluids and electrolytes; **CA**, parenteral nutrition admixture customized (tailored) to the individual patient requirements; **CAFE**, parenteral nutrition admixture customized (tailored) to the individual patient requirements plus extra fluids and electrolytes (P<0.001)



**Europe others:** Belgium, Bulgaria, Croatia, Germany, Hungary, Lithuania, Slovenia, Spain, Sweden. **Oceania:** Australia, New Zealand  
**S. Am.:** Argentina, Brasil

**Figure 3.** Home parenteral nutrition (HPN)-provider ( $P=0.083$ ) and intravenous supplementation (IVS)-admixture type ( $P<0.001$ ) by nature of the underlying disease in patients with chronic intestinal failure.

**HPN-provider:** LP, health care system local pharmacy. HCC, home care company.

**IVS-admixture:** FE, fluids and electrolytes; PA, commercially premixed ready-to-use parenteral nutrition admixture; PAFE, commercially premixed ready-to-use parenteral nutrition admixture plus extra fluids and electrolytes; CA, parenteral nutrition admixture customized (tailored) to the individual patient requirements; CAFE, parenteral nutrition admixture customized (tailored) to the individual patient requirements plus extra fluids and electrolytes

