



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Intravenous supplementation type and volume are associated with 1-year outcome and major complications in patients with chronic intestinal failure

Pironi, Loris; Steiger, Ezra; Joly, Francisca; Wanten, Geert J A; Chambrier, Cecile; Aimasso, Umberto; Sasdelli, Anna Simona; Szczepanek, Kinga; Jukes, Amelia; Theilla, Miriam; Kunecki, Marek; Daniels, Joanne; Serlie, Mireille J; Cooper, Sheldon C; Poullenot, Florian; Rasmussen, Henrik Højgaard; Compher, Charlene W; Crivelli, Adriana; Hughes, Sarah-Jane; Santarpia, Lidia; Guglielmi, Francesco William; Rotovnik Kozjek, Nada; Ellegard, Lars; Schneider, Stéphane M; Matras, Przemysław; Forbes, Alastair; Wyer, Nicola; Zmarzly, Anna; Taus, Marina; O'Callaghan, Margie; Osland, Emma; Thibault, Ronan; Cuerda, Cristina; Jones, Lynn; Chapman, Brooke; Sahin, Peter; Virgili, Núria M; Lee, Andre Dong Won; Orlandoni, Paolo; Matysiak, Konrad; Di Caro, Simona; Doitchinova-Simeonova, Maryana; Masconale, Luisa; Spaggiari, Corrado; Garde, Carmen; Serralde-Zúñiga, Aurora E; Olveira, Gabriel; Krznaric, Zeljko; Petrina Jáuregui, Estrella; Zugasti Murillo, Ana; Suárez-Llanos, José P; Nardi, Elena; Van Gossum, André; Lal, Simon

Published in:
Gut

DOI (link to publication from Publisher):
[10.1136/gutjnl-2018-318172](https://doi.org/10.1136/gutjnl-2018-318172)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2020

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

1
2 1 **The intravenous supplementation type and volume is associated with one-year outcome and the**
3
4 2 **major complications in patients with chronic intestinal failure**

5
6
7 3
8
9 4 Loris Pironi, Ezra Steiger, Francisca Joly, Geert Wanten, Cecile Chambrier, Umberto Aimasso, Anna Simona
10
11 5 Sasdelli, Kinga Szczepanek, Amelia Jukes, Miriam Theilla, Marek Kunecki, Joanne Daniels, Mireille Serlie,
12
13 6 Sheldon C. Cooper, Florian Poullenot, Henrik H. Rasmussen, Charlene Compher, Adriana Crivelli, Sarah-Jane
14
15 7 Hughes, Lidia Santarpia, Francesco W. Guglielmi, Nada Rotovnik Kozjek, Lars Ellegard, Stéphane M.
16
17 8 Schneider, Przemysław Matras , Alastair Forbes, Nicola Wyer, Anna Zmarzly, Marina Taus, Margie
18
19 9 O'Callaghan, Emma Osland, Ronan Thibault, Cristina Cuerda, Lynn Jones, Brooke Chapman, Peter Sahin,
20
21 10 Nuria M. Virgili, Andre Dong Won Lee, Paolo Orlandoni, Konrad Matysiak, Simona Di Caro, Maryana
22
23 11 Doitchinova-Simeonova, Luisa Masconale, Corrado Spaggiari, Carmen Garde, Aurora E. Serralde-Zúñiga,
24
25 12 Gabriel Oliveira, Zeljko Krznicaric, Estrella Petrina Jáuregui, Ana Zugasti Murillo, José P. Suárez-Llanos, Elena
26
27 13 Nardi, André Van Gossum and Simon Lal.

28
29 14 The Home Artificial Nutrition and Chronic Intestinal Failure special interest group of the European Society
30
31 15 for Clinical Nutrition and Metabolism (ESPEN)

32
33
34
35
36
37
38 17 Corresponding author

39
40 18 Loris Pironi

41
42 19 Center for Chronic Intestinal Failure

43
44 20 Department of Digestive System

45
46 21 St. Orsola Hospital, University of Bologna

47
48 22 Via Massarenti, 9 - 40138 Bologna, Italy

49
50 23 Tel: +39 - 051 - 6363073

51
52 24 Fax: +39 - 051 -6364193

53
54 25 Email: loris.pironi@unibo.it

55
56 26 Word count: 3883

57
58
59
60

Loris Pironi 1, Ezra Steiger 2, Francisca Joly 3, Geert J A Wanten 4, Cecile Chambrier 5, Umberto Aimasso 6, Anna Simona Sasdelli 1, Kinga Szczepanek 7, Amelia Jukes 8, Miriam Theilla 9, Marek Kunecki 10, Joanne Daniels 11, Mireille J Serlie 12, Sheldon C Cooper 13, Florian Poullenet 14, Henrik Højgaard Rasmussen 15, Charlene W Compher 16, Adriana Crivelli 17, Sarah-Jane Hughes 18, Lidia Santarpia 19, Francesco William Guglielmi 20, Nada Rotovnik Kozjek 21, Lars Ellegard 22, Stéphane M Schneider 23, Przemysław Matras 24, Alastair Forbes 25, Nicola Wyr 26, Anna Zmarzly 27, Marina Taus 28, Margie O'Callaghan 29, Emma Osland 30, Ronan Thibault 31, Cristina Cuerda 32, Lynn Jones 33, Brooke Chapman 34, Peter Sahin 35, Núria M Virgili 36, Andre Dong Won Lee 37, Paolo Orlandoni 38, Konrad Matysiak 39, Simona Di Caro 40, Maryana Doitchinova-Simeonova 41, Luisa Masconale 42, Corrado Spaggiari 43, Carmen Garde 44, Aurora E Serralde-Zúñiga 45, Gabriel Olveira 46, Zeljko Krznaric 47, Estrella Petrina Jáuregui 48, Ana Zugasti Murillo 49, José P Suárez-Llanos 50, Elena Nardi 1, André Van Gossum 51, Simon Lal 52

1. *Medical and Surgical Sciences, University of Bologna, Bologna, Italy*
2. *Digestive Disease and Surgery Institute, Cleveland Clinic, Cleveland, Ohio, USA*
3. *Service de Gastroentérologie et d'Assistance nutritive, Hôpital Beaujon, Assistance Publique - Hôpitaux de Paris, University of Paris, Clichy, France*
4. *Department of Gastroenterology and Hepatology, Radboud University Medical Center, Nijmegen, The Netherlands*
5. *Centre Hospitalier Universitaire de Lyon, Lyon, Rhône-Alpes, France*
6. *Azienda Ospedaliero Universitaria Citta della Salute e della Scienza di Torino, Torino, Piemonte, Italy*
7. *Stanley Dudrick's Memorial Hospital, Skawina, Poland*
8. *University Hospital of Wales, Cardiff, Cardiff, UK*
9. *Nursing Department, Steyer School of Health Professions, Sackler School of Medicine, Tel Aviv, Israel*
10. *Clinical Nutrition Department, M Pirogow Hospital, Lodz, Poland*
11. *Nottingham University Hospital NHS Trust, Nottingham, UK*
12. *Endocrinology and Metabolism, Academic Medical Center, Amsterdam, The Netherlands*
13. *University Hospitals Birmingham NHS Foundation Trust, Birmingham, Birmingham, UK*
14. *Hôpital Haut-Lévêque, Service d'hépatogastroentérologie, CHU Bordeaux, Pessac, France*
15. *Center for Nutrition and Bowel Disease, Department of Medical Gastroenterology, Aalborg University Hospital, Aalborg, Denmark*
16. *School of Nursing, University of Pennsylvania, Philadelphia, Pennsylvania, USA*
17. *Fundacion Favaloro Hospital Universitario, Buenos Aires, Federal District, Argentina*
18. *Belfast Health and Social Care Trust, Belfast, Belfast, UK*
19. *Department of Clinical Medicine and Surgery, Università degli Studi di Napoli Federico II, Napoli, Campania, Italy*
20. *Gastroenterology and Artificial Nutrition, Hospital Mons. Dimiccoli, Barletta, Trani, Italy*
21. *Institute of Oncology, Ljubljana, Slovenia*
22. *Sahlgrenska Universitetssjukhuset, Goteborg, Sweden*
23. *Centre Hospitalier Universitaire de Nice, Nice, Provence-Alpes-Côte d'Azur, France*
24. *Uniwersytet Medyczny w Lublinie, Lublin, Lubelskie, Poland*

25. *Faculty of Medicine and Health Sciences, University of East Anglia, Norwich, Norfolk, UK*
26. *University Hospital Coventry, Coventry, Coventry, UK*
27. *J Gromkowski City Hospital, Wroclaw, Poland*
28. *Azienda Ospedaliero Universitaria Ospedali Riuniti di Ancona Umberto I G M Lancisi G Salesi, Ancona, Marche, Italy*
29. *Flinders Medical Centre, Bedford Park, Adelaide, Australia*
30. *Royal Brisbane and Women's Hospital, Herston, Brisbane, Australia*
31. *Centre de référence Maladies Rares Digestives, Unité de Nutrition, CHU Rennes, INRAE, INSERM, Université de Rennes, Nutrition Metabolisms and Cancer institute, NuMeCan, Rennes, Bretagne, France*
32. *Hospital General Universitario Gregorio Marañon, Madrid, Madrid, Spain*
33. *Royal Prince Alfred Hospital, Camperdown, New South Wales, Australia*
34. *Nutrition and Dietetics, Austin Health, Heidelberg, Victoria, Australia*
35. *St Imre Hospital, Budapest, Hungary*
36. *Unitat de Nutrició i Dietètica, Hospital Universitari Bellvitge, L'Hospitalet Llobregat, Barcelona, Spain*
37. *Hospital das Clinicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil*
38. *Clinical Nutrition, IRCCS-INRCA, Ancona, Marche, Italy*
39. *Centre for Intestinal Failure, Uniwersytet Medyczny imienia Karola Marcinkowskiego w Poznaniu, Poznan, Poland*
40. *Gastroenterology, UCLH, London, UK*
41. *Bulgarian Executive Agency of Transplantation, Sofia, Bulgaria*
42. *Unita' Locale Socio-Sanitaria N° 22, Bussolengo, Verona, Italy*
43. *Azienda Unita Sanitaria Locale di Parma, Parma, Emilia-Romagna, Italy*
44. *Hospital Universitario de Donostia, San Sebastian, País Vasco, Spain*
45. *Instituto Nacional de Ciencias Medicas y Nutricion Salvador Zubiran, Tlalpan, DF, Mexico*
46. *Hospital Regional Universitario de Málaga, Universidad de Málaga, Málaga, Spain*
47. *University Hospital Centre Zagreb, Zagreb, Croatia*
48. *Complejo Hospitalario de Navarra, Pamplona, Spain*
49. *Hospital Virgen del Camino, Pamplona, Navarra, Spain*
50. *Hospital Universitario Nuestra Señora de la Candelaria, Santa Cruz de Tenerife, Canarias, Spain*
51. *Department of Gastroenterology, Hepatopancreatology and Digestive Oncology, Erasme Hospital, Université Libre de Bruxelles, Brussels, Belgium*
52. *Gastroenterology and Intestinal Failure Unit, Salford Royal Foundation Trust, University of Manchester, Manchester, UK*

1
2 27 **What is already known on this subject?**
3

- 4 28 • Previous studies have demonstrated that several clinical risk factors are associated with outcome
5
6 29 and the risk of parenteral nutrition/intestinal failure-related major complications in patients on
7
8 30 long-term home parenteral nutrition. However, no objective indicator has yet been identified to
9
10 categorize the severity of chronic intestinal failure.
11 31

12
13 32 **What are the new findings?**
14

- 15 33 • The one-year odds of major complications of home parenteral nutrition/intestinal failure (liver
16
17 34 disease and catheter-related blood stream infection) and weaning from home parenteral nutrition
18
19 are independently associated with the type and volume of the intravenous supplementation
20 35 required.
21
22 36 • The one-year odds of death is non-significantly associated with the type of intravenous
23
24 37 supplementation required; future study is required to determine if the latter significantly impacts
25
26 38 on the longer term risk of death in this patient cohort.
27
28 39
29
30

31 40 **How might it impact on clinical practice in the foreseeable future?**
32

- 33 41 • The type and the volume of the intravenous supplementation could be indicators to categorize the
34
35 42 severity of chronic intestinal failure in clinical and research settings.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 43 **Abstract**

3
4 44 **Background and aim**

5
6 45 No marker to categorize the severity of chronic intestinal failure (CIF) has yet been developed. A one-
7
8 46 year international survey was carried out to investigate whether the European Society for Clinical Nutrition
9
10 47 and Metabolism (ESPEN) clinical classification of CIF, based on the type and the volume of the intravenous
11
12 48 supplementation (IVS) could be an indicator of CIF severity.

13
14
15 49 **Methods**

16
17 50 At baseline, participating home parenteral nutrition (HPN)-centers enrolled all adults with ongoing
18
19 51 CIF due to non-malignant disease; demographic data, body mass index, CIF mechanism, underlying disease,
20
21 52 HPN duration and IVS category were recorded for each patient. The type of IVS was classified as fluid and
22
23 53 electrolyte alone (FE) or parenteral nutrition admixture (PN). The mean daily IVS volume, calculated on
24
25 54 weekly basis, was categorized as: <1, 1-2, 2-3, >3 L/day. The severity of CIF was determined by patient
26
27 55 outcome (still on HPN, weaned from HPN, deceased) and the occurrence of major HPN/CIF-related
28
29 56 complications: intestinal failure associated liver disease (IFALD), catheter-related venous thrombosis (CVC-
30
31 57 VT) and catheter-related bloodstream infection (CRBSI).

32
33
34
35 58 **Results**

36
37 59 Fifty-one HPN-centers included 2194 patients. The analysis showed that both IVS type and volume
38
39 60 were independently associated with the odds of weaning from HPN (significantly higher for PN <1 L/day
40
41 61 than for FE and all the PN >1 L/day), patient's death (lower for FE, $p=0.079$), presence of IFALD-
42
43 62 cholestasis/liver failure and occurrence of CRBSI (significantly higher for PN 2-3 and PN >3 L/day).

44
45
46 63 **Conclusions**

47
48 64 The type and the volume of the IVS required by patients with CIF could be indicators to categorize
49
50 65 the severity of CIF in both clinical practice and research protocols.

66 Introduction

67 Intestinal failure (IF) is defined as the reduction of gut function below the minimum necessary for the
68 absorption of macronutrients and/or water and electrolytes, such that intravenous supplementation (IVS) is
69 required to maintain health and/or growth, [1]. Chronic intestinal failure (CIF) is a long-lasting condition
70 that may be reversible or irreversible. Patients with CIF are metabolically stable and receive IVS at home
71 (home parenteral nutrition, HPN) for months, years or lifelong, [2]. Single or multicenter, mostly
72 retrospective, surveys have described risk factors associated with the patient's outcome, such as survival
73 and reversibility of CIF, and with the risk of HPN/IF-related major complications, [3-5]. However, no simple
74 indicator, such as creatinine for kidney disease and SaO₂ for respiratory disease, has yet been identified to
75 categorize the severity of CIF. Such an indicator would be a useful criterion for both clinical practice and
76 research protocols.

77 The European Society for Clinical Nutrition and Metabolism (ESPEN) devised a clinical classification of
78 CIF, to facilitate communication among professionals through an objective categorization of the patients.
79 This was based on patients' requirements for energy and volume of IVS and originally comprised 16
80 categories, [1]. An international cross-sectional survey was carried out to investigate the applicability of this
81 classification and to evaluate factors associated with the IVS requirements of individual patients, [6]. In
82 adult patients with CIF due to non-malignant disease (benign-CIF), the loss of intestinal function appeared
83 more comprehensively represented by IVS volume requirement than by energy requirement. The results
84 enabled the derivation of a new simplified 8-category classification of CIF, based on two types of IVS, either
85 fluid and electrolyte alone (FE) or parenteral nutrition admixture containing energy (PN), and four
86 categories of volume, [6].

87 In order to determine whether ESPEN clinical classification categories could be used as indicators of
88 the severity of CIF, a prospective, multi-center international study was carried out to investigate their
89 association with the patient's outcome and the major complications related to HPN/IF. The results of one-
90 year follow up are reported.

1
2 92 **Material and methods**

3
4 93 *Study design*

5
6 94 This was an international survey involving the retrospective collection of data prospectively recorded
7
8 95 during a one-year follow-up period. The severity of CIF was based on both patient outcome and major
9
10 96 complications related to HPN/IF. The patient's outcome was categorized as still on HPN, weaned from HPN
11
12 97 or deceased. The HPN/IF-related complications were described as the occurrence of intestinal failure
13
14 98 associated liver disease (IFALD-cholestasis or liver failure), central venous catheter associated vein
15
16 99 thrombosis (CVC-VT) or central venous catheter related bloodstream infection (CRBSI) at one-year follow-
17
18
19
20 100 up, [2].

21
22 101 *Baseline HPN center enrollment and patient inclusion*

23
24 102 The baseline data collection was performed on March 1st, 2015. Details regarding HPN-center
25
26 103 enrollment and the patient inclusion criteria have been published in the previous cross-sectional survey
27
28 104 carried out to evaluate the applicability the clinical classification of CIF, [4]. Sixty-five HPN centers from 22
29
30 105 countries enrolled all adult patients (≥ 18 years old) dependent on HPN for CIF on March 1st, 2015. Patients
31
32 106 with either benign or malignant disease were included. Patients with active malignant disease were termed
33
34 107 as having "cancer-CIF". Patients without malignant disease at time of inclusion in the study were termed as
35
36 108 having "benign-CIF". Invasive intra-abdominal desmoid disease was included in the benign group, because
37
38 109 of the chronic nature of the condition and reflecting the fact that it is an established indication for intestinal
39
40 110 transplantation, [2]. A total of 3239 patients, 9.9% with cancer-CIF and 91.1% with benign-CIF were
41
42 111 included, [4]. For the purpose of the present study, only patients with benign-CIF were investigated.

43
44
45
46 112 *Follow up data collection*

47
48 113 The one-year follow up was carried out on patients enrolled in the 2015 baseline cross-sectional
49
50 114 study. In February 2016, the study coordinator (LP) sent an email to the HPN centers that participated in
51
52 115 the 2015 cross-sectional survey, to invite them to participate in the follow up. The study protocol
53
54 116 (**Supplemental material 1**) and the structured database for the data collection were attached to the
55
56 117 invitation letter. Centers were asked to include relevant data from the patient's medical records between
57
58 118 March 1st, 2015 and March 1st, 2016 and details of the patient's outcome on March 1st, 2016.

1
2 119 Data were collected into a structured questionnaire embedded in an Excel (Microsoft Co., 2013) database
3
4 120 (the ESPEN CIF Action Day database) (**Supplemental material 2**). Centers were invited to contact the study
5
6 121 coordinator for any additional explanation or instruction.
7

8 9 122 *Ethical statement*

10
11 123 The study was approved by the Home Artificial Nutrition and Chronic Intestinal Failure (HAN&CIF)
12
13 124 special interest group of ESPEN. The research was based on anonymized information taken from patient
14
15 125 records at time of data collection. The study was conducted with full regard to confidentiality of the
16
17 126 individual patient. Each patient was labelled with two anonymized identifications: one given by the HPN-
18
19 127 center and one given by the database manager. Ethical committee approval was obtained by the individual
20
21 128 HPN centers according to local regulations. The collected data were used only for the study purpose.
22
23 129 Contributing centers have been anonymized for data analysis and presentation.
24
25

26 27 130 *Statistical analysis*

28
29 131 The term HPN described the provision of IVS, either FE or PN, at the patient's home. Weaning from
30
31 132 equated to stopping IVS [1]. The clinical classification of CIF consisted of 8 categories, based on the type
32
33 133 and volume of IVS, calculated as daily mean of the total volume infused per week: volume per day of
34
35 134 infusion \times number of infusions per week / 7 (mL/day): FE1 or PN1, ≤ 1000 ; FE2 or PN2, 1001–2000; FE3 or
36
37 135 PN3, 2001–3000; FE4 or PN4, >3000 , [6].
38
39

40 136 The pathophysiological mechanisms of IF were classified as short bowel syndrome with end-
41
42 137 jejunostomy (SBS-J), with jejunocolic anastomosis (SBS-JC) or with jejunoleileal anastomosis and total colon
43
44 138 in continuity (SBS-JIC), intestinal dysmotility (Dysmotility), intestinal fistulas (Fistulas), mechanical
45
46 139 obstruction (Obstruction) and extensive small bowel mucosa disease (Mucosal disease), [6].
47
48

49 140 The underlying diseases were grouped as follows: inflammatory bowel disease (IBD), comprising
50
51 141 Crohn's disease and ulcerative colitis; mesenteric ischemia, comprising mesenteric arterious or venous
52
53 142 infarction and non-occlusive ischemia; acute post-surgical complications; chronic intestinal pseudo-
54
55 143 obstruction, idiopathic or secondary to intestinal or systemic diseases (CIPO); short bowel syndrome due to
56
57 144 causes other than mesenteric ischemia, including intra-abdominal adhesions, volvulus, cured cancer,
58
59 145 abdominal trauma, intestinal malformation (Other causes of SBS); radiation enteritis; miscellaneous
60

1
2 146 (collagenous diseases, intra-abdominal desmoids, intestinal polyposis, autoimmune enteropathy,
3
4 147 neurological disease, congenital mucosal disease, celiac disease and other diseases not included in the
5
6 148 above categories).

8
9 149 The reasons for weaning from HPN were categorized as medical and surgical, the latter included non-
10
11 150 transplant procedures and intestinal transplantation. The odds of weaning from HPN were evaluated by
12
13 151 two models of analysis: one including all the weaned patients and one excluding patients weaned because
14
15 152 of non-transplant surgery.

17
18 153 HPN/IF-related complications at March 1st 2016 were categorised as follows: IFALD and CVC-VT
19
20 154 reported as prevalent cases, when they were already present at baseline, and as incident case, when they
21
22 155 developed during the one-year follow up. CRBSI were categorized as incident cases occurring during the
23
24 156 one-year follow up. The incident or prevalent nature of CVC-TV was collected at time of filling out the
25
26 157 database on March 1st 2016, whereas the incident or prevalent nature of IFALD was collected after
27
28 receiving the filled out database, by asking a specific question to the participating centers.

30
31 159 Practice variation by HPN-center was weighted by including in the statistical analysis the number of
32
33 160 patients enrolled in the study by the individual HPN-center. We also estimated a model with center as a
34
35 161 random effect. Although a slight improvement was noted in model fitting (i.e. in terms of percentage of
36
37 162 correctly classified subjects, with subjects correctly classified: 81% vs 80%), the random effect was omitted
38
39 in order to preserve the ease of interpretation of the model.

41
42 164 Data are reported as mean \pm standard deviation (SD), median and range, absolute and relative
43
44 165 frequencies.

46
47 166 Binomial and multinomial logistic regressions were carried out for multivariate analysis. The odds
48
49 167 ratio was used to measure the association between the independent variables and the outcome. A
50
51 168 competing risk regression model based on Fine and Gray's proportional subhazards approach was also
52
53 169 performed in order to model the time to occurrence of competing outcomes. Subhazard ratios were
54
55 presented together with the Cumulative Incidence Functions.

57
58 171 Missing data were excluded from the analysis. Two-tailed p-values less than 0.05 were considered as
59
60 172 statistically significant. All p-values were not corrected for multiple-hypothesis testing.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

173
174
175

The analyses were performed using the IBM SPSS Statistics package for Windows, version 23.0 (BM
Co., Armonk, NY, USA), R software for Windows, version 3.5.1 (<http://cran.r-project.org>) and STATA/IC 16.0
for Windows

Accepted Manuscript

1
2 176 **Results**

3
4 177 *Study population*

5
6 178 Fifty-one of the 65 HPN-centers which contributed in the 2015 database collection, participated in
7
8
9 179 the 2016 follow up; this included 2194 of the 2919 benign-CIF (75.1%) patients enrolled in 2015. Most of
10
11 180 the patients (79.7%) were from European Countries, the remaining were from Israel, US, Mexico, Argentina,
12
13 181 Brazil and Australia. The mean number of patients included in the follow up by center was 43.0 ± 54.1
14
15 182 (median: 19; range: 1-231); number of patients by center: ≤ 19 , n. 26 (51.0%) centers, patients n. 198 (9.0%
16
17
18 183 of total); 20-49, n. 8 (15.7%), patients 253 (11.5%); 50-99, n. 10 (19.6%) centers, patients n.657 (29.9%);
19
20 184 ≥ 100 n. 7 (13.7%) centers, patients n. 1086 (49.5%). Nine centers included < 5 patients, each.

21
22 185 **Table 1** shows the baseline characteristics and the one-year outcome of the cohort of patients with
23
24 186 benign-CIF included in the present study. Two-thirds were females. At baseline, the mean \pm SD, (median;
25
26
27 187 range) patient age, BMI and HPN duration were 51.1 ± 16.2 (56.5;18.0-98.0) years, 22.3 ± 4.4 (21.7; 10.5-59.6)
28
29 188 kg/m² and 58.0 ± 70.2 (33.2,0-474) months, respectively. SBS-J was the most frequent pathophysiological
30
31 189 mechanism of IF (35.9% of cases). The most frequent underlying disease was Crohn's disease (21.1%). The
32
33 190 type of IVS was FE in 7.9% of patients and PN in 92.1%. The IVS volume was significantly lower in the
34
35
36 191 subgroup of patients receiving FE 1055.8 ± 859.6 mL/day (median 857.1, range 107.1–4800.0) than in those
37
38 192 receiving PN 1055.8 ± 859.6 mL/day (median 1785.7, range 81.7-7542.8) ($P < 0.001$).

39
40 193 *One-year outcome*

41
42 194 On March 1st, 2016, 1740 (79.3%) patients were still on HPN, 298 (13.6%) were weaned from HPN
43
44
45 195 and 156 (7.1%) were deceased (**Table 1**). The reason for weaning from HPN was reported in 272 cases:
46
47 196 spontaneous intestinal adaptation in 138 (50.7%), non-transplant surgery in 114 (41.9%) (surgical intestinal
48
49 197 continuity reconstruction in 97 cases), ITx in 14 (5.1%) and intestinal growth factor therapy in 6 (2.2%)
50
51 198 cases. The cause of death was reported in 146 cases: HPN/IF-related in 6 (4.1%) patients (CRBSI 5, IFALD 1),
52
53
54 199 underlying disease-related in 64 (43.8%) (4 due to ITx complications) and other causes (neither HPN/IF nor
55
56 200 underlying disease-related) in 76 (52.1%) cases.

57
58 201 HPN/IF complications were recorded in 1859 of the 2194 (84.7%) patients. The presence of IFALD-
59
60 202 cholestasis/liver failure was reported in 97 patients (4.4%), 66 prevalent and 31 of which were incident

1
2 203 cases: cholestasis 63 (64.9%), impending liver failure 11 (11.3%), overt liver failure 18 (18.6%), not specified
3
4 204 5 (5.1%). A CVC-VT was present in 53 patients (2.9%), 23 prevalent and 30 incident cases. During the follow
5
6 205 up, 273 patients (14.7%) had 344 episodes of CRBSI: one episode in 224 (82.0%); two episodes in 40
7
8 206 (14.7%); three episodes in 5 (1.8%); four episodes in 2 (0.7%); 7 and 10 episodes in 1 (0.4%) patient each
9
10
11 207 one.

12
13 208 A variation of IVS type and volume between baseline and the end of follow up was observed in 317
14
15 209 (14.4%) patients: 22 patients changed the IVS type from FE to PN (12.6% of FE), 26 patients changed from
16
17 210 PN to FE (1.3% of PN) and 269 patients changed PN volume (13.3% of PN) (**Supplemental table 1**).

Table 1. Baseline characteristics and one-year outcome and home parenteral nutrition (HPN)/intestinal failure (IF) – related major complications of adult patients with chronic intestinal failure due to benign disease. (IFALD, intestinal failure associated liver disease: cholestasis or liver failure; CVC-VT, central venous catheter-associated deep vein thrombosis; CRBSI, catheter related bloodstream infections). Data are reported as percentages of cases.

Baseline patient cohort n. 2194	One-year outcome			One-year HPN/IF major complications		
	Still on HPN n. 1740	Weaned from HPN n. 298	Deceased n. 156	IFALD n. 97	CVC-VT n. 53	CRBSI n. 273
Gender (%)						
Male (n. 811)	77.3	15.4	7.3	6.7	3.1	14.8
Female (n. 1383)	80.5	12.5	7.0	4.4	2.7	14.6
Age, years (%)						
≤29 (n.187)	78.6	19.8	1.6	8.1	1.9	24.2
30-49 (n.575)	80.3	16.2	3.5	5.8	3.1	16.4
50-69 (n.990)	80.2	12.5	7.3	4.4	3.0	12.3
≥70 (n.442)	76.2	10.0	13.8	5.1	2.7	13.8
BMI, kg/m², (%)						
≤15 (n.57)	80.7	10.5	8.8	6.3	4.2	14.6
15-18.5 (n.324)	74.4	15.7	9.9	6.3	4.7	14.6
18.5-25 (n.1334)	82.5	10.9	6.6	5.5	2.4	13.3
25-30 (n.363)	75.5	18.7	5.8	3.8	2.5	16.8
≥30 (n.111)	67.6	24.3	8.1	3.0	4.0	25.3
Not reported (n.5)						
Duration of HPN, yrs (%)						
≤1 (n.575)	60	31.7	8.3	5.4	1.3	14.8
1-3 (n.575)	79.1	13.2	7.7	4.9	1.9	16.1
3-10 (n.748)	89.6	4.5	5.9	5.3	4.0	14.7
>10 (n.293)	91.5	1.7	6.8	5.5	4.4	12.2
Not reported (n.3)						

1							
2	Mechanism of IF (%)						
3	SBS-J (n.788)	78.3	14.6	7.1	7.5	2.0	14.6
4	SBS-JC (n.459)	88.2	7.4	4.4	4.0	5.4	12.1
5	SBS-JIC (n.140)	77.9	17.1	5.0	4.4	2.7	18.6
6	Fistulas (n.149)	64.4	23.5	12.1	4.0	0.8	16.8
7	Dysmotility (n.398)	81.7	10.8	7.5	2.9	3.2	15.7
8	Obstruction (n.104)	70.2	18.3	11.5	5.6	1.1	14.4
9	Mucosal disease (n.156)	73.7	17.9	8.3	4.6	2.3	14.6
10							
11							
12							
13	Underlying disease (%)						
14	IBD (n.480)	79.8	15.0	5.2	4.5	1.9	11.8
15	CIPO (n.299)	85.6	10.4	4.0	3.4	2.7	18.9
16	Other causes of SBS (n.178)	80.7	9.6	9.6	5.8	5.1	19.1
17	Miscellaneous (n.218)	76.4	18.0	5.6	7.3	4.5	19.2
18	Mesenteric ischemia (n.395)	79.2	10.6	10.1	6.4	4.5	13.4
19	Radiation enteritis (n.164)	82.3	10.4	7.3	2.3	1.5	9.8
20	Acute surgical complications (n.306)	70.3	20.6	9.2	7.4	1.2	16.3
21	Not reported (n.154)						
22							
23							
24							
25	Clinical classification of CIF						
26	(IVS volume/day of infusion) (%)						
27	FE1 (≤1 L) (n.118)	89.8	9.3	0.8	0.9	0.9	6.6
28	FE2 (1-2 L) (n.40)	77.5	12.5	10.0	2.7	0	10.8
29	FE3 (2-3 L) (n.10)	80.0	20.0	0	0	22.2	22.2
30	FE4 (>3 L) (n.6)	83.3	16.7	0	16.7	0	0
31	PN1 (≤1 L) (n.384)	77.1	16.9	6.0	1.9	4.5	10.7
32	PN2 (1-2 L) (n.944)	78.9	13.3	7.7	3.8	3.1	15.0
33	PN3 (2-3 L) (n.482)	78.4	13.5	8.1	8.6	2.0	16.2
34	PN4 (>3 L) (n.210)	81.4	11.0	7.6	12.4	1.6	22.0
35							
36							
37							
38							
39	Type of IVS (%)						
40	Total FE (n.174)	86.2	10.9	2.9	1.9	1.9	8.2
41	Total PN (n.2020)	78.7	13.8	7.5	12.4	2.9	15.3
42							
43							
44							
45							
46							

1
2
3
4
5
6
7
8
9

215 BMI, body mass index; HPN, home parenteral nutrition; IF, intestinal failure; SBS-J, short bowel syndrome with jejunostomy; SBS-JC, short bowel syndrome with
216 jejuno-colic anastomosis with partial colon; SBS-JIC, short bowel syndrome with jejunio-ileal anastomosis with intact colon in continuity; CIF chronic intestinal
217 failure; IVS, intravenous supplementation; FE, fluid and electrolytes alone; PN, parenteral nutrition-admixture; IBD (inflammatory bowel disease): Crohn's disease
218 n.462, Ulcerative colitis n.18; CIPO (chronic intestinal pseudo-obstruction): primary n.222, secondary n.77; Other causes of SBS: Intra-abdominal adhesions n.72,
219 Volvulus n.46, Cured cancer n.21, Abdominal trauma n.26, Intestinal malformation n.13; Miscellaneous: Collagenous disease n. 40, Intra-abdominal desmoids n.22,
220 Intestinal polyposis n.16, Autoimmune enteropathy n.14, Neurological disease n.11, Congenital mucosal disease n.14, Celiac disease n.8, Other diseases n.93

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

1
2 221 *Factors associated with the patient's one-year outcome and HPN/IF-complications*

3
4 222 Weaning from HPN, death, presence of IFALD-cholestasis/liver failure or CVC-VT at the end of the
5
6 223 follow up, and occurrence of CRBSI during the one-year follow up were considered the dependent
7
8 224 variables. The baseline patient's demographics, IF mechanism, underlying disease, IVS characteristics and
9
10
11 225 number of patients enrolled in the study by individual HPN-center were included as independent variables.

12
13 226 Considering the low number of total patients receiving the FE type as well as the very low number of
14
15 227 those receiving FE3 and FE4, patients on FE were grouped in a unique cohort for the outcome analyses.

16
17
18 228 **Figure 1** shows the cumulative incidence of weaning off and of death according with the type and the
19
20 229 volume of the IVS. When comparing the cumulative incidence function of weaning off and death among
21
22 230 groups, non-statistically significant differences were found ($p=0.329$ and $p=0.148$ for weaning off and death
23
24 231 respectively): incidence of weaning off was greater in PN1 group and lower in PN4 and FE groups; incidence
25
26 232 of death was lower in the FE group .

27
28
29 233 *Odds of one-year outcome*

30
31 234 The associations with the IVS type and volume were analyzed in comparison with the PN1 category.
32
33 235 The odds of weaning from HPN (**Figure 2 and Supplemental table 2**): a) were lower in the FE type category
34
35 236 and in the greatest PN volume categories (PN2, PN3 and PN4); b) were lower in the oldest decades of age,
36
37
38 237 in the longest duration of HPN categories and in the miscellaneous group of underlying diseases; c) were
39
40 238 higher in the underweight, overweight and obese BMI categories; d) showed no association with the
41
42 239 number of patients included in the study by individual HPN-centers.

43
44 240 The results were confirmed when excluding those patients who were weaned because of a non-
45
46 241 transplant surgical procedure (**Supplemental table 3**). Furthermore, significant lower odds of weaning off
47
48
49 242 were observed in patients who had SBS-J or SBS-JC as mechanisms of IF and in those who had an underlying
50
51 243 disease categorized in the miscellaneous group.

52
53 244 The odds of death on HPN (**Figure 2**): a) showed a non-statistically significant decreased risk for the
54
55 245 FE type of IVS with respect to PN1 type ; b) were higher in the oldest age categories and in the lowest BMI
56
57
58 246 categories; c) in comparison with SBS-J mechanism of IF, they were lower in the other SBS types and were
59
60 247 higher in the other mechanisms of IF, excepting the extensive mucosal disease; d) were increased in the

1
2 248 mesenteric ischemia and decreased in the CIPO groups of underlying disease; e) showed a negative
3
4 249 association with the number of patients included in the study by individual HPN-centers.
5

6 250 The competing risk analysis for the risk of death and of weaning from HPN confirmed the results of
7
8
9 251 the multinomial analysis (**Supplemental table 4**).

10
11 252 *Odds of major complications of HPN/IF*

12
13 253 The results are reported in **table 2**.

14
15 254 The odds of the presence of IFALD-cholestasis/liver failure: a) were higher in the greatest PN volume
16
17
18 255 categories in comparison with PN1 and were similar between PN1 and the FE type of IVS; b) were lower in
19
20 256 dysmotility mechanism of IF; c) were higher in the group with acute surgical complications as underlying
21
22 257 disease; d) showed no association with the number of patients included in the study by individual HPN-
23
24 258 centers

25
26 259 The odds of the presence of CVC-VT: a) showed no association with the IVS categories; b) were
27
28
29 260 higher in the in the longest HPN duration categories and in the underweight and obese categories of BMI;
30
31 261 c) showed a negative association with the number of patients included in the study by individual HPN-
32
33 262 centers.

34
35 263 The odds of CRBSI: a) were higher with the increase of the volume of the PN and were similar
36
37
38 264 between PN1 and the FE type of IVS; b) were lower in older patients; c) were higher in the overweight and
39
40 265 obese category of BMI and in the CIPO and miscellaneous categories of underlying disease; d) showed a
41
42 266 negative association with the number of patients included in the study by individual HPN-centers.
43

44 267 The analyses including only the incident cases of IFALD and of CVC-VT showed non-statistically
45
46
47 268 significant odds ratios (**Supplemental table 5**).

48
49 269

50

51

52

53

54

55

56

57

58

59

60

Table 2. Binomial logistic analysis of factors independently associated with the one-year probability of the intestinal failure associated liver disease, cholestasis or liver failure (IFALD), central venous catheter-associated deep vein thrombosis (CVC-VT) and catheter related bloodstream infections (CRBSI) in adult patients on home parenteral nutrition for chronic intestinal failure. Number of analyzed cases due to complete case approach: IFALD-cholestasis/liver failure (presence: 91, absence: 1610), CVC-VT (presence: 49, absence: 1652), CRBSI (presence: 257, absence: 1443).

Independent factors	IFALD-cholestasis/liver failure			CVC-VT			CRBSI		
	OR	95% CI	P	OR	95% CI	P	OR	95% CI	P
IVS type and volume									
PN1 (≤ 1 L)	1			1			1		
PN2 (1-2 L)	1.824	0.738-4.508	0.193	0.784	0.370-1.663	0.527	1.507	0.967-2.346	0.070
PN3 (2-3 L)	3.794	1.522-9.458	0.004	0.513	0.194-1.354	0.178	1.664	1.018-2.721	0.042
PN4 (> 3 L)	4.828	1.792-13.004	0.002	0.437	0.110-1.732	0.239	2.523	1.423-4.475	0.002
FE (total)	0.849	0.202-3.569	0.823	0.384	0.078-1.894	0.240	0.743	0.362-1.525	0.418
Gender									
Male	1			1			1		
Female	0.703	0.449-1.099	0.122	0.798	0.434-1.470	0.470	1.066	0.797-1.427	0.666
Age (years)									
≤ 29	1			1			1		
30-49	0.775	0.355-1.693	0.522	1.875	0.495-7.108	0.355	0.597	0.371-0.961	0.034
50-69	0.694	0.317-1.517	0.360	2.017	0.545-7.461	0.293	0.469	0.291-0.756	0.002
≥ 70	0.822	0.339-1.994	0.665	1.657	0.396-6.939	0.489	0.551	0.319-0.952	0.033
BMI (kg/m²)									
18.5-25.0	1			1			1		
≤ 15.0	1.064	0.233-4.858	0.936	2.705	0.566-12.935	0.212	0.837	0.333-2.106	0.706
15.0-18.5	1.612	0.872-2.983	0.128	2.335	1.093-4.987	0.028	1.061	0.697-1.615	0.782
25.1-30.0	0.611	0.310-1.207	0.156	1.134	0.470-2.738	0.779	1.460	1.012-2.108	0.043
≥ 30.0	0.591	0.172-2.036	0.405	3.124	0.829-11.769	0.092	2.769	1.580-4.851	0.000
Duration of HPN (years)									
≤ 1	1			1			1		

1										
2	1-3	1.025	0.553-1.899	0.938	1.526	0.516-4.515	0.445	1.157	0.787-1.703	0.458
3	3-10	1.070	0.594-1.929	0.821	2.889	1.093-7.636	0.032	1.110	0.762-1.618	0.586
4	>10	0.982	0.470-2.052	0.962	3.477	1.179-10.256	0.024	0.853	0.522-1.393	0.524
5										
6	Mechanism of IF									
7	SBS-J	1			1			1		
8	SBS-JC	0.649	0.338-1.245	0.193	1.926	0.848-4.374	0.117	0.964	0.629-1.479	0.868
9	SBS-JIC	0.629	0.208-1.902	0.411	0.760	0.189-3.060	0.699	1.377	0.754-2.516	0.298
10	Fistulas	0.593	0.219-1.607	0.304	0.580	0.071-4.733	0.611	1.207	0.676-2.154	0.525
11	Dysmotility	0.317	0.115-0.878	0.027	1.431	0.428-4.790	0.561	0.759	0.422-1.365	0.357
12	Obstruction	1.072	0.373-3.083	0.897	0.438	0.050-3.859	0.457	0.800	0.376-1.704	0.563
13	Mucosal disease	0.712	0.260-1.945	0.507	0.868	0.203-3.713	0.849	0.824	0.445-1.526	0.538
14										
15										
16	Underlying disease									
17	IBD	1			1			1		
18	CIPO	1.459	0.481-4.427	0.505	1.244	0.303-5.113	0.762	2.098	1.092-4.03	0.026
19	Other causes of SBS	1.265	0.501-3.192	0.619	2.308	0.742-7.18	0.149	1.490	0.843-2.634	0.170
20	Miscellaneous	2.038	0.909-4.567	0.084	2.308	0.723-7.366	0.158	1.712	1.002-2.924	0.049
21	Mesenteric ischemia	1.468	0.732-2.943	0.280	2.063	0.798-5.333	0.135	1.012	0.626-1.635	0.963
22	Radiation enteritis	0.613	0.170-2.210	0.454	0.824	0.161-4.206	0.816	0.967	0.487-1.918	0.923
23	Acute surgical complications	2.210	1.089-4.482	0.028	0.573	0.142-2.311	0.433	1.143	0.700-1.864	0.593
24										
25										
26	N. of patients included by	0.939	0.765-1.153	0.551	0.580	0.451-0.745	0.000	0.710	0.626-0.806	0.000
27	center									
28										
29										

304 BMI, body mass index; HPN, home parenteral nutrition; IF, intestinal failure; SBS-J, short bowel syndrome with jejunostomy; SBS-JC, short bowel syndrome with jejunocolon
315 anastomosis with partial colon; SBS-JIC, short bowel syndrome with jejunocolon anastomosis with intact colon; IBD, inflammatory bowel disease, CIPO, chronic intestinal
326 pseudo-obstruction; CIF chronic intestinal failure; IVS, intravenous supplementation; FE, fluid and electrolytes alone; PN, parenteral nutrition-admixture.
337
34
35
36
37
38
39
40
41
42
43
44
45
46

Discussion

This is the first study aimed at investigating the association between IVS requirement, CIF outcome and the occurrence of major complications in a very large international cohort of HPN-dependent patients with CIF due to benign underlying disease. The results show that both the type and the volume of the IVS are independently associated with the one-year odds of weaning from HPN and of HPN/IF-associated major complications, as well as with the risk of mortality, albeit the latter observation being based on a non-statistically significant finding. In patients with CIF, the type and the volume of the IVS requirement primarily depends on the degree of the reduction of gut function, [6]. However, other factors may be involved, such as the patient's metabolic condition and vital organ function, the patient's compliance with the prescribed treatment (e.g. drugs and dietary prescriptions) as well as the treatment protocols of the multidisciplinary team caring for him/her, [1,2]. Therefore, while any association between IVS characteristics and the patient's outcome or the occurrence of HPN/IF complications may not be considered causal, they may indicate that the type and the volume of the IVS reflect comprehensive odds of morbidity and mortality for HPN-dependent patients, independently from the factors that may have determined their prescription. This is further strengthened by the observation that none of the other independent factors entered in the multivariate analysis was contemporaneously associated with odds of weaning from HPN, death and occurrence of IFALD and CRBSI. These data support the potential role of the ESPEN clinical classification of CIF, based on the type and the volume of the IVS, as a potential indicator of CIF severity. Further follow up surveys are required to investigate if this could be translated into a long-term marker of CIF.

The one-year odds of death depended on the interaction between the IVS type and volume rather than on either characteristic alone. Indeed, a non-statistically significant decreased risk of death was observed in those receiving the FE type of IVS, but no association was found with the PN-volume alone; since HPN-related deaths were very rare, [3], these results would suggest a less severe clinical condition in patients with CIF requiring only FE supplementation. Future studies will clarify whether the association between the current volume categories of IVS and the risk of death will prove to be statistically significant

1
2 305 in the long-term and/or whether a different categorization of the IVS volume will capture any association
3
4 306 between IVS volume and risk of death, both in the short and in the long-term.
5

6 307 The one-year probability of weaning from HPN was associated with both the type and volume of IVS.
7
8
9 308 The PN1 volume (≤ 1 L/day) showed higher odds of weaning than either the greater PN-volumes or FE-type
10
11 309 IVS. There could be several reasons for a longer maintenance of low volume FE than of low volume PN IVS:
12
13 310 a more difficult intestinal rehabilitation of fluid and electrolytes than of macronutrient absorption due to
14
15 311 concomitant secondary mechanisms of IF causing increased intestinal secretion [1]; the concomitant
16
17 312 presence of a reduced kidney function requiring the maintenance of optimal hydration, [2,7]; physician's
18
19
20 313 and/or patient's perception of a lower risk of IVS-associated complications with FE than with PN; patient's
21
22 314 better acceptance of FE than of PN, because of shorter duration of FE infusion compared to PN, [2]; the
23
24 315 lower cost of FE. All of these factors would make weaning from FE slower/less likely than weaning from PN.
25

26 316 The risk of IFALD and of the occurrence of CRBSIs were also associated with both the type and the
27
28
29 317 volume of the IVS, whereas no association was observed with the presence of CVC-VT. The odds of IFALD
30
31 318 and of CRBSI were greater in patients receiving the highest volumes of PN in comparison with the lowest
32
33 319 PN-volumes and the FE-type of IVS. Furthermore, the odds of these complications were higher in the
34
35 320 greater PN volume categories. These data are in keeping with previous studies, [2, 8, 9]. The pathogenesis
36
37
38 321 of IFALD is multifactorial, including factors related to the IVS, underlying gastrointestinal disease and
39
40 322 systemic factors, especially episodes of sepsis [2, 10]. Intravenous supplementation overfeeding and a high
41
42 323 amount of lipid emulsion are recognized causes of IFALD [2, 10]. Similarly, CRBSI occurrence has also
43
44 324 previously been reported to occur more frequently in those dependent on an increased number of days of
45
46
47 325 IVS, [8]; this may relate to more frequent handling of the central venous catheter increasing infection risk
48
49 326 or the association between macronutrients, vitamins and trace metals affecting microbial growth in the PN
50
51 327 admixture, [11, 12].
52

53 328 Most of the other independent factors found to be associated with patient's outcome and HPN/IF
54
55 329 complications were in keeping with data from previous studies, [2, 3, 8, 10]. As expected, non-transplant
56
57
58 330 surgery was the cause of weaning off HPN in a large percentage of patients, [13]. Notably, data on the
59
60 331 causes of death on long-term HPN are consistent with previous observations, [3-5, 13-15], even though the

1
2 332 percentage of HPN-related deaths (4%) was lower than that reported in longer retrospective surveys (10-
3
4 333 14%), [3-5,13-15]. This could be due to the short duration of the present follow up, as it is known that the
5
6 334 rate of the HPN-related death increases with the duration of the treatment, [4]. The 344 episodes of CRBSI
7
8 335 registered in the 1859 patients accounted for a rate of CRBSI of 0.18 per catheter-year, or 0.50 per 1000
9
10
11 336 catheter-days, a rate that is in the range reported in the literature, [2]. The 30 incident cases of CVC-VT
12
13 337 observed at one-year follow up, accounted for an incidence rate of 0.016 per catheter-year, that is also in
14
15 338 the lower range of the literature (0.02-0.09 cases per catheter-year), [2,16]. The same incidence rate can be
16
17
18 339 accounted for the 31 incident cases of IFALD-cholestasis/liver failure. This data is of some relevance as no
19
20 340 prospective study has yet been carried out on this HPN/IF complication, [17].
21

22 341 The weakness of the study is mainly represented by the retrospective collection of data prospectively
23
24 342 recorded in the previous 12 months, which would imply a risk of some underreporting, and by not using a
25
26 343 co-morbidity index to assess the patient's general condition. However, CIF can develop as a complication of
27
28
29 344 a number of gastrointestinal or systemic underlying diseases having different pathogenesis and outcomes,
30
31 345 [1]. Therefore, the underlying disease that we collected and included in the statistical analysis, could be
32
33 346 considered as a surrogate co-morbidity index, as supported by the finding of an increased risk of death in
34
35 347 patients with CIF due to mesenteric ischemia. The strength of the study is clearly reflected by its
36
37
38 348 international multicenter structure and by the study population, which is the largest cohort of patients with
39
40 349 CIF ever enrolled in a single survey. These characteristics should avoid the potential bias associated with
41
42 350 the analysis of individual center cohorts, which could be influenced by local practice and expertise and
43
44 351 mitigate the impact of the above possible weakness on statistical analyses. Considering that CIF is a rare
45
46 352 disease, [1] the observation of lower odds of death and of HPN/IF complications in HPN-center which
47
48
49 353 included the larger number of patients in the study would support the importance of creating networks
50
51 354 facilitating the referral of CIF patients to few but appropriately organized and experienced centers, [18].
52
53 355 Finally, the agreement between our results and the risk factors, other than IVS, reported by previous
54
55
56 356 studies would support the overall reliability of our findings.
57

58 357 In conclusion, the type of the IVS, either FE or PN, and the volume of the PN-admixture, as
59
60 358 categorized by the ESPEN clinical classification of CIF, were found to be independently associated with the

1
2 359 one-year risk of death, of weaning from HPN and of major complications of HPN/IF. These results support
3
4 360 the ESPEN categorization of the IVS as potential marker of the severity of CIF.
5

6 361
7

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Manuscript

References

1. Pironi L, Arends J, Baxter J, et al.; Home Artificial Nutrition & Chronic Intestinal Failure and the Acute Intestinal Failure Special Interest Groups of ESPEN. ESPEN endorsed recommendations. Definition and classification of intestinal failure in adults. *Clin Nutr* 2015;34:171-80 doi: 10.1016/j.clnu.2014.08.017. Epub 2014 Sep 21
2. Pironi L, Arends J, Bozzetti F, et al.; Home Artificial Nutrition & Chronic Intestinal Failure Special Interest Group of ESPEN. ESPEN guidelines on chronic intestinal failure in adults. *Clin Nutr* 2016;35:247-307 doi: 10.1016/j.clnu.2016.01.020. Epub 2016 Feb 8.
3. Pironi L, Goulet O, Buchman A, et al.; Home Artificial Nutrition and Chronic Intestinal Failure Working Group of ESPEN. Outcome on home parenteral nutrition for benign intestinal failure: a review of the literature and benchmarking with the European prospective survey of ESPEN. *Clin Nutr* 2012;31:831-45 doi: 10.1016/j.clnu.2012.05.004. Epub 2012 Jun 2.
4. Pironi L, Joly F, Forbes A, et al.; Home Artificial Nutrition & Chronic Intestinal Failure Working Group of the European Society for Clinical Nutrition and Metabolism (ESPEN). Long-term follow-up of patients on home parenteral nutrition in Europe: implications for intestinal transplantation. *Gut* 2011;60:17-25 doi: 10.1136/gut.2010.223255. Epub 2010 Nov 10.
5. Joly F, Baxter J, Staun M, et al.; ESPEN HAN CIF group. Five-year survival and causes of death in patients on home parenteral nutrition for severe chronic and benign intestinal failure. *Clin Nutr* 2018;37:1415-1422 doi: 10.1016/j.clnu.2017.06.016. Epub 2017 Jun 19.
6. Pironi L, Konrad D, Brandt C, et al. Clinical classification of adult patients with chronic intestinal failure due to benign disease: An international multicenter cross-sectional survey. *Clin Nutr* 2018;37:728-738 doi: 10.1016/j.clnu.2017.04.013. Epub 2017 Apr 19.
7. Agostini F, Sasdelli AS, Guidetti M, et al. Outcome of kidney function in adults on long-term home parenteral nutrition for chronic intestinal failure. *Nutrition* 2018, doi: <https://doi.org/10.1016/j.nut.2018.10.005>

- 1
2 387 8. Dreesen M, Foulon V, Spriet I, et al. Epidemiology of catheter-related infections in adult patients
3
4 388 receiving home parenteral nutrition: a systematic review. *Clin Nutr* 2013;32:16-26 doi:
5
6 389 10.1016/j.clnu.2012.08.004. Epub 2012 Aug 21.
7
8
9 390 9. Sasdelli AS, Agostini F, Pazzeschi C, et al. Assessment of Intestinal Failure Associated Liver Disease
10
11 391 according to different diagnostic criteria. *Clin Nutr* Published online first: 8 May 2018. doi:
12
13 392 10.1016/j.clnu.2018.04.019.
14
15 393 10. Lal S, Pironi L, Wanten G, et al.; Home Artificial Nutrition & Chronic Intestinal Failure Special Interest
16
17 Group of the European Society for Clinical Nutrition and Metabolism (ESPEN). Clinical approach to
18 394 the management of Intestinal Failure Associated Liver Disease (IFALD) in adults: A position paper
19
20 395 from the Home Artificial Nutrition and Chronic Intestinal Failure Special Interest Group of ESPEN. *Clin*
21
22 396 *Nutr* 2018;37:1794-1797 doi: 10.1016/j.clnu.2018.07.006. Epub 2018 Jul 9.
23
24 397
25
26 398 11. Austin PD, Hand KS, Elia M. Systematic review and meta-analyses of the effect of lipid emulsion on
27
28 microbial growth in parenteral nutrition. *J Hosp Infect* 2016;94:307-319 doi:
29 399 10.1016/j.jhin.2016.08.026. Epub 2016 Sep 7.
30
31 400
32
33 401 12. Buchman AL, Opilla M, Kwasny M, et al. Risk factors for the development of catheter-related
34
35 bloodstream infections in patients receiving home parenteral nutrition. *JPEN J Parenter Enteral Nutr*
36 402 2014;38:744-9 doi: 10.1177/0148607113491783. Epub 2013 Jun 6.
37
38 403
39
40 404 13. Amiot A, Messing B, Corcos O, et al. Determinants of home parenteral nutrition dependence and
41
42 405 survival of 268 patients with non-malignant short bowel syndrome. *Clin Nutr* 2013;32:368-74 doi:
43
44 406 10.1016/j.clnu.2012.08.007. Epub 2012 Aug 23.
45
46
47 407 14. Dibb M, Soop M, Teubner A, et al. Survival and nutritional dependence on home parenteral nutrition:
48
49 408 Three decades of experience from a single referral centre. *Clin Nutr* 2017;36:570-576 doi:
50
51 409 10.1016/j.clnu.2016.01.028. Epub 2016 Feb 22.
52
53
54 410 15. Brandt CF, Tribler S, Hvistendahl M, et al. Home Parenteral Nutrition in Adult Patients With Chronic
55
56 411 Intestinal Failure: Catheter-Related Complications Over 4 Decades at the Main Danish Tertiary
57
58 412 Referral Center. *JPEN J Parenter Enteral Nutr* 2018;42:95-103 doi: 10.1177/0148607116678766. Epub
59
60 413 2017 Dec 11.

- 1
2 414 16. Pironi L, Corcos O, Forbes A, et al.; ESPEN Acute and Chronic Intestinal Failure Special Interest
3
4 415 Groups. Intestinal failure in adults: Recommendations from the ESPEN expert groups. *Clin Nutr*
5
6 416 2018;37:1798-1809 doi: 10.1016/j.clnu.2018.07.036. Epub 2018 Aug 18..
7
8
9 417 17. Bond A, Huijbers A, Pironi L, Schneider SM, Wanten G, Lal S. Review article: diagnosis and
10
11 418 management of intestinal failure-associated liver disease in adults. *Aliment Pharmacol Ther*. 2019
12
13 419 Sep;50(6):640-653. doi: 10.1111/apt.15432. Epub 2019 Jul 25. Review. PubMed PMID: 31342540.
14
15 420 18. <https://www.engage.England.nhs.uk/consultation/severe-intestinal-failure-services-for-adults/>
16
17
18 421
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1

2 422 **Figure headings and footnotes**

3

4 423

5

6

7 424 **Figure 1.** Cumulative one-year incidence of weaning from home parenteral nutrition (HPN) and of death
8 425 according to the intravenous supplementation (IVS) type and volume. IVS type: FE, fluid and electrolyte
9 426 alone; PN, parenteral nutrition admixture containing energy. IVS volume (L/day): PN1, ≤ 1 ; PN2, 1-2; PN3, 2-
10 427 3; PN4, >3 . FE1, FE2, FE3 and FE4 were grouped in a unique cohort (FE). The patient risk set is reported in
11 428 **supplemental table 2.**

12 429

13 430

14

15 431

16

17 432

18 **Figure 2.** Forest plot of the analysis of factors associated with the one-year probability of weaning from
19 433 home parenteral nutrition (HPN) or of death and of home parenteral nutrition /intestinal failure (HPN/IF) in
20 434 adult patients with chronic intestinal. Number of analyzed cases due to complete case approach: 2035 (Still
21 435 in HPN: 1610, Weaned from HPN: 278, Deceased: 147).

22 436

23 437

24 438

25 439

26 439 Footnote figure 2

27 440 BMI, body mass index; HPN, home parenteral nutrition; IF, intestinal failure; SBS-J, short bowel syndrome
28 441 with jejunostomy; SBS-JC, short bowel syndrome with jejunocolon anastomosis with partial colon; SBS-JIC,
29 442 short bowel syndrome with jejunocolon anastomosis with intact colon; IBD, inflammatory bowel disease,
30 443 CIPO, chronic intestinal pseudo-obstruction; CIF chronic intestinal failure; IVS, intravenous
31 444 supplementation; FE, fluid and electrolytes alone; PN, parenteral nutrition-admixture.

32 445

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

1
2 446 **Acknowledgements**
3

4 447 **Contributing coordinators and centers by country**
5

6 448
7 449 **Argentina**

8 450 Adriana N. Crivelli, Hector Solar Muñiz; Hospital Universitario Fundacion Favaloro, Buenos Aires

9 451 **Australia**

10 452 Brooke R. Chapman; Austin Health, Melbourne

11 453 Lynn Jones, Peter Lim; Royal Prince Alfred Hospital, Sydney

12 454 Margie O'Callaghan; Flinders Medical Centre, Adelaide

13 455 Emma Osland, Ruth Hodgson, Siobhan Wallin, Kay Lasenby; Royal Brisbane and Women's Hospital,
14 456 Herston

15 457 **Belgium**

16 458 Andre Van Gossum: Hôpital Erasme, Brussels

17 459 **Brazil:**

18 460 Andre Dong Won Lee; Hospital das Clinicas da Faculdade de Medicina da Universidade de São
19 461 Paulo, São Paulo

20 462 **Bulgaria**

21 463 Maryana Doitchinova-Simeonova; Bulgarian Executive Agency of Transplantation, Sofia

22 464 **Croatia**

23 465 Zeljko Krznaric; University Hospital Centre Zagreb, Zagreb

24 466 **Denmark**

25 467 Henrik Højgaard Rasmussen; Center for Nutrition and Bowel Disease, Aalborg University Hospital,
26 468 Aalborg

27 469 **France**

28 470 Cecile Chambrier; Hospices Civils de Lyon, Centre Hospitalier Lyon Sud, Lyon

29 471 Francisca Joly, Vanessa Boehm, Julie Bataille, Lore Billiauws; Beaujon Hospital, Clichy

30 472 Florian Poullenot; CHU de Bordeaux, Hôpital Haut-Lévêque, Pessac

31 473 Stéphane M. Schneider; CHU Archet, Nice

32 474 Ronan Thibault; Nutrition unit, CHU Rennes, Nutrition Metabolisms and Cancer institute,

33 475 NuMeCan, INRA, INSERM, Université Rennes, Rennes

34 476 **Hungary**

35 477 Peter Sahin, Gábor Udvarhelyi; St. Imre Hospital, Budapest

36 478 **Israel**

37 479 Miriam Theilla; Rabin Medical Center, Petach Tikva

38 480 **Italy**

39 481 Anna Simona Sasdelli, Loris Pironi; S. Orsola University Hospital, Bologna

40 482 Umberto Aimasso, Merlo F. Dario; Città della Salute e della Scienza, Torino

41 483 Luisa Masconale, Valentino Bertasi; ULSS 22 Ospedale Orlandi, Bussoleto

42 484 Francesco W. Guglielmi, Nunzia Regano; San Nicola Pellegrino Hospital, Trani

43 485 Paolo Orlandoni; IRCCS- INRCA, Ancona

44 486 Santarpia Lidia, Maria C. Pagano, Lucia Alfonsi; Federico II University, Italy

45 487 Corrado Spaggiari; AUSL di Parma, Parma

46 488 Marina Taus, Debora Busni; Ospedali Riuniti, Ancona

47 489 **México**

48 490 Aurora E. Serralde-Zúñiga; Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán,
49 491 México City

50 492 **Poland**

51 493 Marek Kunecki; M. Pirogow Hospital, Lodz

- 1
2 494 Przemysław Matras; Medical University of Lublin, Lublin
3 495 Konrad Matysiak; H.Święcicki University Hospital, Poznań
4 496 Kinga Szczepanek; Stanley Dudrick's Memorial Hospital, Skawina
5 497 Anna Zmarzły; J. Gromkowski City Hospital, Wrocław
6 498 **Slovenia**
7 499 Nada Rotovnik Kozjek; Institute of Oncology, Ljubljana
8 500 **Spain**
9 501 Cristina Cuerda; Hospital General Universitario Gregorio Marañón, Madrid
10 502 Carmen Garde; Hospital Universitario Donostia, San Sebastian
11 503 Nuria M. Virgili; Hospital Universitari de Bellvitge, Barcelona
12 504 Gabriel Olveira; IBIMA, Hospital Regional Universitario de Málaga, Universidad de Málaga, Málaga
13 505 M^a Estrella Petrina Jáuregui; Complejo Hospitalario de Navarra, Pamplona
14 506 José P. Suárez-Llanos; Hospital Universitario Nuestra Señora de Candelaria, Santa Cruz de Tenerife
15 507 Ana Zugasti Murillo; Hospital Virgen del Camino, Pamplona
16 508 **Sweden**
17 509 Lars Ellegard; Sahlgrenska University Hospital, Gothenburg
18 510 **The Netherlands**
19 511 Mireille Serlie, Cora Jonker; Academic Medical Center, Amsterdam
20 512 Geert Wanten; Radboud University Medical Center, Nijmegen
21 513 **United Kingdom**
22 514 Sheldon C. Cooper; University Hospitals Birmingham NHS Foundation Trust, Birmingham
23 515 Joanne Daniels; Nottingham University Hospital NHS Trust, Nottingham
24 516 Simona Di Caro, Niamh Keane, Pinal Patel; University College Hospital, London
25 517 Alastair Forbes; Norfolk and Norwich University Hospital, Norwich
26 518 Sarah-Jane Hughes; Regional Intestinal Failure Service, Belfast Health and Social Care Trust,
27 519 Northern Ireland
28 520 Amelia Jukes, Rachel Lloyd; University Hospital of Wales, Cardiff
29 521 Simon Lal, Arun Abraham, Gerda Garside, Michael Taylor; Salford Royal NHS Foundation Trust,
30 522 Salford
31 523 Nicola Wyer, Reena Parmar, Nicola Burch; University Hospital, Coventry
32 524 **United States of America**
33 525 Charlene Compher; Hospital of the University of Pennsylvania, Philadelphia, PA
34 526 Denise Jezerski, Ezra Steiger; Cleveland Clinic Foundation, Cleveland, OH
35
36 527
37
38
39
40
41
42
43
44
45
46 528 **Statistical analysis performed by:** Elena Nardi, PhD statistician, Giorgia Brillanti, statistician, and
47
48 529 Dr. Fabrizio Ghigliano, statistician; Department of Medical and Surgical Sciences, University of
49
50
51 530 Bologna, Italy
52
53 531 **Database manager:** Dr. Marianna Mastroberto, MD, PhD and Dr. Fabrizio Ghigliano, statistician;
54
55
56 532 Department of Medical and Surgical Sciences, University of Bologna, Italy
57
58 533
59
60 534 **Funding source**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

535 The project of the ESPEN database for Chronic Intestinal Failure was promoted by the ESPEN
536 Executive Committee in 2013, was approved by the ESPEN Council and was supported by an ESPEN
537 grant.


538 **Statement of authorship**

539 LP devised the study protocol, collected the data, analyzed the results and drafted the manuscript.

540 The Home Artificial Nutrition & Chronic Intestinal Failure Special Interest Group of ESPEN
541 discussed and approved the protocol study, discussed the results and reviewed the manuscript
542 before submission. All the co-Authors participated in the acquisition of data, revised the final
543 analysis, approved the final version of the manuscript and were accountable for all aspects of the
544 work.

546 **Conflict of interest statements**

547 None declared






Figure 2

1
2 1 **The intravenous supplementation type and volume is associated with one-year outcome and the**
3
4 2 **major complications in patients with chronic intestinal failure**
5

6
7 3
8
9 4 Loris Pironi, Ezra Steiger, Francisca Joly, Geert Wanten, Cecile Chambrier, Umberto Aimasso, Anna Simona
10
11 5 Sasdelli, Kinga Szczepanek, Amelia Jukes, Miriam Theilla, Marek Kunecki, Joanne Daniels, Mireille Serlie,
12
13 6 Sheldon C. Cooper, Florian Poullenot, Henrik H. Rasmussen, Charlene Compher, Adriana Crivelli, Sarah-Jane
14
15 7 Hughes, Lidia Santarpia, Francesco W. Guglielmi, Nada Rotovnik Kozjek, Lars Ellegard, Stéphane M.
16
17 8 Schneider, Przemysław Matras , Alastair Forbes, Nicola Wyer, Anna Zmarzly, Marina Taus, Margie
18
19 9 O'Callaghan, Emma Osland, Ronan Thibault, Cristina Cuerda, Lynn Jones, Brooke Chapman, Peter Sahin,
20
21 10 Nuria M. Virgili, Andre Dong Won Lee, Paolo Orlandoni, Konrad Matysiak, Simona Di Caro, Maryana
22
23 11 Doitchinova-Simeonova, Luisa Masconale, Corrado Spaggiari, Carmen Garde, Aurora E. Serralde-Zúñiga,
24
25 12 Gabriel Oliveira, Zeljko Krznicaric, Estrella Petrina Jáuregui, Ana Zugasti Murillo, José P. Suárez-Llanos, Elena
26
27 13 Nardi, André Van Gossum and Simon Lal.
28

29
30 14 The Home Artificial Nutrition and Chronic Intestinal Failure special interest group of the European Society
31
32 15 for Clinical Nutrition and Metabolism (ESPEN)
33
34
35

36 16
37
38 17 Corresponding author

39
40 18 Loris Pironi

41
42 19 Center for Chronic Intestinal Failure

43
44 20 Department of Digestive System

45
46 21 St. Orsola Hospital, University of Bologna

47
48 22 Via Massarenti, 9 - 40138 Bologna, Italy

49
50 23 Tel: +39 - 051 - 6363073

51
52 24 Fax: +39 - 051 -6364193

53
54 25 Email: loris.pironi@unibo.it

55
56 26 Word count: 37163883
57
58
59
60