



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

ADHD and communication difficulties

A meta-analysis based on parental CCC reports

De Lopez, Kristine M. Jensen; Møbjerg Sørensen, Lise; Jensen, Christina Mohr

Published in:
ChildLanguage Symposium 2015

Publication date:
2015

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

De Lopez, K. M. J., Møbjerg Sørensen, L., & Jensen, C. M. (2015). ADHD and communication difficulties: A meta-analysis based on parental CCC reports. In *ChildLanguage Symposium 2015* (pp. 98). University of Warwick.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



CeDAPS
AALBORG UNIVERSITY

ADHD and communication difficulties: a meta-analysis based on parental CCC reports

Kristine Jensen de López¹, Lise Møbjerg Sørensen¹ & Christina Mohr Jensen^{1,2}

¹Clinic for Developmental Communication Disorder,

Center for Developmental & Applied Psychological Science (CeDAPS)

²Research Unit for Child and Adolescent Psychiatry, Aalborg University Hospital, Denmark



CeDAPS
AALBORG UNIVERSITY

Child Language Symposium, July 20th - 21st, 2015, Warwick University, UK

Child Language Symposium, July 20th - 21st, 2015, Warwick University, UK

Introduction

Several scholars have suggested that SLI and ADHD show overlapping symptoms.^[1,4,7,11,13,14] Children with ADHD are often neither assessed for language problems nor treated for such, potentially increasing the risk for negative outcomes^[3] and being a barrier for the effectiveness of classical non-pharmacologically therapies such as parent-training programs.

The few studies comparing expressive language use by children with SLI, ADHD and typically developing children (TD) show that children with ADHD produce more “mazes” – disfluencies, false starts ect. compared to the children with SLI and TD children, and are more challenged in sentence recall compared to TD children.^[12] Tense/agreement difficulties in combination with sentence recall difficulties have showed to distinguish SLI children from children with ADHD.^[11]

Overall aim:

The aim of this study is to systematically review studies comparing communicative abilities of children with ADHD to their typically developing peers and carry out a meta-analysis in order to gain better knowledge of the boundaries between these two groups.

Methods and Materials

Two databases; PsycINFO & PubMed searched from November 6th, 2014 to November 11th, 2014 using combinations of the specific keywords (ADHD, language, speech, communication, pragmatic, language disorder, SLI, Children’s communication checklist and CCC). Reference lists of retrieved articles were also examined.

The most used tool for assessing communication abilities of children with ADHD was the Children’s Communication Checklist, version 1 and 2. Our meta-analysis therefore focused on results that included this tool and reported by parents.

Both checklists have 70-item that cover children’s communication behaviors across respectively 9 and 10 subscales. The checklists are noted as an objective measure with acceptable levels of reliability ($r = .80$)^[2,4,10] and validity ($J = 0.85$).^[4,10]

The meta-analysis was performed using a random-effect, standardized mean differences (SMD), 95% confidence intervals (95% CI) and interpretation of effect size as small effect ≥ 0.2 ; medium effect ≥ 0.5 ; large effect ≥ 0.8 .

See Table 1. for specific criteria used in the study

Inclusion criteria	Exclusion criteria
Participants $\geq 5 - 17 \geq$ years	Non peer-reviewed studies
At least one clinical diagnosed ADHD group compared with a group of typical developed children	Dyslexia, stuttering, speech disorder, aphasia
Use CCC / CCC-2	IQ < 85
Primary studies	No report of means and standard deviations

Table 1. Criteria used in the CCC / CCC2 literature search

Results

- Six unique studies
- A total of 171 children with a clinical diagnosis of ADHD
- Age 5 - 17 years (mean age 9.8 years)
- All studies included a TD control group

Description of included studies

- 3 studies used 1st version of CCC^[2,5,6]
- 3 studies used the 2nd edition^[4,7,15]


The results of the synthesized studies indicate that children diagnosed with ADHD have significantly more language problems compared to peers on all 10 subscales of CCC with a large overall effect size ($d = .86$, $z = 5.48$, $p < .001$).

The largest effects sizes were found on the scales Inappropriate Initiation ($d = 1.82$), Use of Context ($d = 1.86$) and Social Relations ($d = 1.92$). These scales are all related to the pragmatic aspect of language thereby indicating that children with ADHD have pragmatic language problems.

Scale	SMD	95% confidence interval		Z-value	p-value
		Lower limit	Upper limit		
A: Speech output	0.88	0.55	1.20	5.29	< .001
B: Syntax	0.82	0.61	1.04	7.88	< .001
C: Semantic	1.26	0.91	1.62	6.94	< .001
D: Coherence	1.60	1.15	2.04	7.09	< .001
E: Inappropriate initiation	1.82	1.39	2.25	8.31	< .001
F: Stereotyped language use	1.35	0.98	1.73	7.02	< .001
G: Use of context	1.86	1.38	2.33	7.68	< .001
H: Nonverbal communication	1.21	0.65	1.77	4.25	< .001
I: Social relations	1.92	1.48	2.35	8.68	< .001
J: Interests CCC	0.70	0.32	1.08	3.62	< .001
J: Interests CCC-2	1.55	0.91	2.18	4.78	< .001

Table 2. Summary of indices obtained in the meta-analysis

Flowchart



Discussion

ADHD & SLI two of a kind?

Our results show that children with ADHD are reported as having significantly more language problems than TD children, and that they particularly show problems initiating speech and using context correctly. These disabilities are characterized as pragmatic disabilities and the findings are consistent with previous research. The relatively high difficulties reported for Social Relations could be due to these pragmatic difficulties. Together these challenges could be explained related to the executive functioning problems characterizing ADHD as proposed by Leonard.^[8]

According to the guidelines for clinical interpretation of the CCC2 children with SLI are expected to show profiles reflecting least challenges on scales E to H, which include Inappropriate initiation and Use of context; the two scales that were most challenging for children with ADHD. Children with a pragmatic language disorder on the other hand are also characterized with these same type of problems.

Whether CCC2 could be used as an additional clinical tool for identifying pragmatic deficits among children with ADHD, as well as identifying children holding a double deficit (ADHD + SLI) warrants further investigation.

Conclusions

Our meta-analysis shows that children with ADHD possess challenges on all ten domains of the CCC compared to TD peers, while adequate pragmatic skills might be the most challenging area. However children with ADHD are exclusively offered treatment for behavioral problems alone, whereas pragmatic treatment should be an additional option.

Contact

kristine@hum.aau.dk, Clinic for Developmental Communication Disorders, Aalborg University, Denmark,

References

1. Baker, L., & Cantwell, D. (1982). Developmental, social and behavioral characteristics of speech and language disordered children. *Child Psychiatry and Human Development*, 12, 195-206
2. Bishop, D.V., Baird, G. (2001) Parent and teacher report of pragmatic aspects of communication: use of the Children’s communication checklist in a clinical setting. *Developmental Medicine & Child Neurology*, 43(12), 809-818
3. Clegg, J., Hollis, C., Mawhood, L. & Rutter, M. (2005). Developmental language disorders – follow-up in later adult life. *Cognitive, language and psychosocial outcomes. Journal of Child Psychology and Psychiatry*, 46 (2), 128-149
4. Geurts, H.M., Embrechts, M. (2008) Language profiles in ASD, SLI, and ADHD. *Journal of Autism and Developmental Disorders*, 10(1), 1931-1943
5. Geurts, H.M., Verté, S., Oosterlaan, J., Roeyers, H., Hartman, C.A., Mulder, E.J., Bervoets-Corriu, J.A., Sergeant, J.A. (2004) Can the Children’s Communication Checklist differentiate between children with autism, children with ADHD, and normal controls? *Journal of Child Psychology and Psychiatry*, 45(8), 1437-1453
6. Geurts, H.M., Verté, S., Oosterlaan, J., Roeyers, H., & Sergeant, J.A. (2004). How specific are executive functioning deficits in attention deficit hyperactivity disorders and autism? *Journal of Child Psychology and Psychiatry*, 45(4-5), 836-854
7. Holland, W.A., Binger, E., Holland, J., Heimann, M. (2012) Exploring language profiles for children with ADHD and children with asperger syndrome. *Journal of Attention Disorders*, 16(1), 34-43
8. Leonard, L. B. (2014). *Children with specific language impairment*. 2nd edition. MIT Press
9. Love, A., & Thompson, M. (1988). Language disorders and attention disorders in young children referred for psychiatric services: Analysis of prevalence and a conceptual synthesis. *American Journal of Orthopsychiatry*, 58, 52-64
10. Norbury, C. F., Nash, M., Baird, G., & Bishop, D. (2004). Using a parental checklist to identify diagnostic groups in children with communication impairment: a validation of the Children’s Communication Checklist-2. *International Journal of Language and Communication Disorders*, 39(3), 343-364
11. Redmond, S. M., Thompson, H.L., Goldstein, S. (2011). Psycholinguistic profiling differentiates specific language impairment from typical development and from attention-deficit/hyperactivity disorder. *Journal of Speech, Language and Hearing Research*, 54(1), 99-117
12. Redmond, S. M. (2004). Conversational profiles of children with ADHD, SLI and typical development. *Clinical Linguistics & Phonetics*, 18(10) 107-125
13. Timler, G. R. (2014). Use of the Children’s Communication Checklist-2 for classification of language impairment risk in young school-age children with attention-deficit/hyperactivity disorder. *American Journal of Speech-Language Pathology*, 23(1), 79-83
14. Tiroch, E., & Cohen, A. (1998). Language deficit with attention-deficit disorder: A prevalent comorbidity. *Journal of Child Neurology*, 13, 493-497
15. Vainanen, R., Loukusa, S., Molanen, I., Vilhena, A. (2014) Language and pragmatic profile in children with ADHD measured by Children’s Communication Checklist 2nd edition. *Logopedics Phoniatrics Vocology* 39, 179-187