



AALBORG UNIVERSITY
DENMARK

Aalborg Universitet

Long-term follow-up after acute achilles tendon rupture — Does treatment strategy influence functional outcomes?

Nilsson, Laurine; Thorlund, Jonas Bloch; Kjær, Inge Lunding; Kazlauskas, Andrius; Christensen, Marianne

Published in:
Foot

DOI (link to publication from Publisher):
[10.1016/j.foot.2020.101769](https://doi.org/10.1016/j.foot.2020.101769)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2021

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Nilsson, L., Thorlund, J. B., Kjær, I. L., Kazlauskas, A., & Christensen, M. (2021). Long-term follow-up after acute achilles tendon rupture — Does treatment strategy influence functional outcomes? *Foot*, 47, Article 101769. <https://doi.org/10.1016/j.foot.2020.101769>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

1 Title

2 Long-Term Follow-Up after Acute Achilles Tendon Rupture - Does Treatment
3 Strategy Influence Functional Outcomes?
4

5 Abstract

6 **Background:** Patients struggle to fully recover after an Achilles tendon rupture.

7 Although several studies has investigated surgical and non-surgical treatment, the
8 best treatment is still uncertain. The aim of this study was to investigate long-term
9 patient-reported outcomes and objective measures 4 years after acute Achilles
10 tendon rupture and compare whether outcomes differed between patients treated on
11 basis of the previous regimen preferring surgical treatment and the new regimen
12 preferring functional rehabilitation.

13 **Methods:** Achilles tendon Total Rupture Score (ATRS), number of re-ruptures and
14 the objective measures; Achilles tendon resting angle, calf circumference, heel-rise
15 height, and muscle endurance were measured at a 4-year follow-up. Patients were
16 recruited from Aalborg University Hospital.

17 **Results:** Seventy-six patients were included (29% female). The mean ATRS was
18 71.4 (95% CI: 65.8 to 77.1) at 4 years follow-up. No difference in ATRS was
19 observed between Previous regimen and New regimen at any timepoint (time x
20 group interaction, $p=0.8509$). The injured side was still significantly impaired
21 compared with the non-injured side in terms of all objective measures. Impairments
22 in objective measures were not dependent on the preferred treatment strategy.

23 **Conclusions:** Patient reported impairments and objective functional deficits persist
24 4 years after an acute Achilles tendon rupture. No differences in patient reported
25 outcome or objective measures at the 4 years follow-up was observed between the

1 old treatment regimen preferring surgery compared with the new treatment regimen
2 preferring functional rehabilitation.

3

4 **Keywords**

5 Achilles, functional rehabilitation, rupture, surgery, tendon

6

7 **Abbreviations**

8 ATRA: Achilles tendon resting angle

9 ATRS: Acute Achilles tendon rupture score

10 DADB: Danish Achilles tendon Database

11 HRH: Heel-rise height

12 ME: Muscle Endurance

13

14 **Funding**

15 This research did not receive any specific grant from funding agencies in the public,
16 commercial, or not-for-profit sectors.

17

18 **1. Introduction**

19 *1.1 Background*

20 Acute Achilles tendon rupture is a common injury with an incidence of 8 to 37 per

21 100,000 person years.¹⁻⁵ Over the last decades, the incidence has increased,^{1,3,5}

22 presumably due to a larger elderly population and their activity level.⁶ Previous

23 studies found functional deficits 1-2 years after Achilles tendon rupture,^{7,8} but long-

24 term follow-up studies are scarce.

1

2 Despite several randomized controlled trials and meta-analyses investigating
3 surgical versus non-surgical treatment of an acute Achilles tendon rupture, the
4 optimal treatment is still uncertain.⁹⁻¹¹ The risk of re-rupture is generally considered
5 lower with surgical treatment compared to non-surgical treatment with
6 immobilization.^{12,13} The advantage of non-surgical treatment is the absence of
7 complications due to anesthesia and surgical complications such as risk of infection,
8 nerve injury, scar tissue, and other surgery-related complications.^{14,15} Historically,
9 since surgery was introduced, it has been the preferred treatment, but during the last
10 10 years, the number of non-surgically treated patients has increased.^{1,3,5}

11 The change in treatment appears to have been inspired by publications of
12 randomized controlled trials,^{16,17} followed by a meta-analysis showing no difference
13 between surgical and non-surgical treatment regarding re-rupture rate if a functional
14 rehabilitation protocol with early mobilization was followed.¹⁸ These results indicate
15 that rehabilitation is important for the outcome and could be more important than if
16 the rupture is treated surgically or not.

17

18 Previously, the preferred treatment strategy for acute Achilles tendon ruptures in
19 Denmark country was surgical treatment. However, later research evidence has
20 suggested that many patients with acute Achilles tendon ruptures may do well with
21 conservative treatment (i.e. functional rehabilitation). Thus, the preferred (or
22 recommended) treatment strategy was changed. As part of quality assurance and
23 research, it is important to investigate if such decisions (i.e. to change the preferred
24 treatment strategy) manifest in better or at least similar outcomes compared with the
25 previous strategy. In this study we included patients from a single hospital clinical

1 database, where the preferred treatment strategy was changed from regimen with
2 surgery as the preferred, to a regimen with functional rehabilitation (non-surgical) as
3 the preferred treatment to investigate if this impacted the long term outcomes.

4

5 *1.2 Aim*

6 The aim of this study was to investigate long-term patient-reported outcomes and
7 objective measures 4 years after acute Achilles tendon rupture and compare
8 whether outcomes differed between patients treated on basis of the previous
9 regimen preferring surgical treatment and the new regimen preferring functional
10 rehabilitation.

11

12 **2. Material and Method**

13 *2.1 Materials*

14 Patients included in this long-term follow-up were treated at Aalborg University
15 Hospital and recruited from the Danish Achilles tendon Database (DADB)¹⁹
16 Long-term follow-up was conducted from 03/2018 to 04/2018. Inclusion criteria were
17 first-time Achilles tendon rupture between 04/2012 and 03/2015, minimum 18 years
18 old at time of rupture, and patients registered in the DADB in Aalborg University
19 Hospital. Exclusion criteria were bilateral rupture, delayed treatment onset (> 14
20 days), and if patients were not able to participate in testing. Patients were invited
21 through secure e-mail.

22

1 *2.2 Treatment procedure*

2 The regimens were as described below, but each surgeon could deviate from the
3 regimen if they considered it relevant in the specific case due contraindications,
4 patient preferences or other clinical reasons.

5 *2.2.1 Previous regimen – preferring surgery as first choice treatment*

6 Before November 2013, the primary treatment for Achilles tendon rupture was
7 surgical treatment using open end-to-end sutures. Surgery was performed with the
8 patient in prone position, usually under local anesthesia, otherwise under general or
9 spinal anesthesia. Postoperatively, the patients were placed in a ROM walker for 6
10 weeks. The ROM walker was locked in 30 degrees plantarflexion for two weeks,
11 followed by 30- to 15-degree range of motion in the third week and 30- to 0-degree
12 range of motion in the last 3 weeks. Severe comorbidities such as diabetes or
13 vascular disease with risk of impaired healing were contraindications to surgical
14 treatment. Patients treated non-surgically were placed in a ROM walker for 8 weeks.
15 During the first 2 weeks, the ROM walker was locked at 30 degrees of plantarflexion,
16 and in the third and fourth weeks, it was locked at 15 degrees plantarflexion. In the
17 following weeks, range of motion in the ROM walker was allowed, with 30- to 15-
18 degree range of motion in the fifth week and 30-0 degrees in the last 3 weeks.
19 Protected weight bearing was allowed when the foot position reached neutral
20 position for both surgical and non-surgical treatment. All patients were referred to
21 physiotherapy for one instruction in home exercises upon removal of the orthosis.

22

23 *2.2.2 New regimen – preferring functional rehabilitation as first choice treatment*

24 In the new treatment regimen functional rehabilitation was preferred treatment. In
25 this study, functional rehabilitation is defined as an organized treatment with early

1 mobilization for patients treated non-surgically. After November 2013, the primary
2 treatment was changed to non-surgical treatment with functional rehabilitation. After
3 diagnosis, the leg was placed in a walker boot with wedges for 8 weeks. For the first
4 2 weeks, the patients were immobilized with the foot in plantar flexion with three
5 wedges and were non-weight bearing. One wedge was removed every second week
6 starting from the second week. From the third week, weight bearing and range of
7 motion exercises were allowed. Five times per day, the patient removed the walker
8 boot and performed 25 repetitions of dorsi-plantarflexion and inversion-eversion with
9 restricted dorsiflexion to neutral (0 degrees). In the sixth week, exercises with an
10 elastic band (25 repetitions 5 times per day) were added. After 8 weeks, the walker
11 boot was removed, and the patients attended 8 weeks of rehabilitation in group
12 sessions led by a physiotherapist. Surgical treatment were indicated for Achilles
13 avulsion from the calcaneus or open lacerations. Patients treated surgically after
14 November 2013 followed the same rehabilitation program as patients treated with
15 functional rehabilitation.

16

17 *2.3 Outcome*

18 The main outcome were patient-reported Achilles tendon Total Rupture Score
19 (ATRS),²⁰ and the number of re-ruptures. The ATRS is determined based on 10
20 questions about pain and function. The score ranges from 0 to 100, where 100
21 equals no limitations. For objective measurements, Achilles tendon resting angle
22 (ATRA), calf circumference, heel-rise height (HRH), and muscle endurance (ME)
23 were used. ATRA measures the angle of the foot with the knee in 90-degree flexion.
24 The foot will be relative more in dorsiflexion if the Achilles tendon is elongated. ATRA
25 is used as a clinical measure for Achilles tendon length. It was performed as

1 described in the reliability study by Carmont et al.²¹ Calf circumference was
2 measured 15 cm distal from the anteromedial knee joint line.²¹ HRH is the maximum
3 height at one repetition single-leg heel rise on a 10-degree incline. ME was
4 measured using MuscleLab (Ergotest Technology, Oslo, Norway).²² MuscleLab has
5 previously been used for monitoring patients with Achilles tendon ruptures.⁷

6

7 *2.4 Four-year Follow-up*

8 At the 4-year follow-up, patients were all welcomed by an orthopedic surgeon (ILK)
9 for a consultation and information about the testing procedures. Another orthopedic
10 surgeon (AK) examined all objective measures, and a physiotherapist (LN) guided all
11 the patients to complete the ME test.

12

13 *2.5 Statistics*

14 Descriptive data is presented as mean with standard deviation, median with
15 interquartile range or numbers and percentages as appropriate. Differences between
16 groups in continuous descriptive data were compared using an unpaired t-test (for
17 ATRS also a Mann-Whitney test was performed). Categorical demographics were
18 compared using a chi-squared test. Differences in objective measures between the
19 affected side and the contralateral side was tested using paired t-tests. The Limb
20 Symmetry Index (LSI) was used when comparing objective measures between
21 groups and was calculated as follows: $(\text{Injured Side} \div \text{Non-injured side}) \times 100$.
22 Difference between groups in LSI was assessed using unpaired t-tests.

23 To assess the change in ATRS over time (1 year, 2 years, and 4 years after rupture),
24 a mixed linear effect model was used with time and group as fixed effects and
25 person (i.e. ID) as random effect. Due to the low number of re-ruptures, statistical

1 analyses for this outcome were not conducted. We used STATA, version 14.0 for all
2 analyses.

3

4 *2.6 Ethics*

5 All patients received written and oral information before giving written consent. The
6 Danish National Ethics Committee on Health Research was consulted, but approval
7 was not necessary for this type of study.

8

9 **3. Results**

10 One hundred and ninety-two patients were invited to participate. Of them, 116 did
11 not want to participate or did not show up at the follow-up as invited, 76 were
12 included in the study. Details can be seen in Fig. 1. Characteristics of included
13 patients are presented in Table 1. Patients who refused to participate or did not show
14 up did not differ from included patients in age, sex, or type of treatment (Table 2).

15

16 Figure 1

17

18 Table 1

19

20 Table 2

21

22 At the 4-year follow-up, the mean ATRS for all patients was 71.4 (95% CI: 66.4 to
23 77.1). Patients following Previous regimen had mean ATRS of 69.1, 65.9, and 73.9
24 at 1-, 2-, and 4-year follow-ups (Fig. 2). For patients following New regimen, the
25 mean scores were 61.5, 66.1, and 69.4 (Fig. 2). No difference was observed

1 between groups at any time. At 4 years follow-up the difference between treatment
2 strategies were 4.5 points (95% CI: -6.8 to 15.8)(Table 1). Change in ATRS over
3 time did not differ between groups (time x group interaction, $p=0.8509$) (Fig. 2).
4 Number of re-ruptures did not differ between groups. The Previous regimen group
5 had one re-rupture, and the New regimen group had two.

6

7 Figure 2

8

9 At the 4-year follow-up, there was a significant difference between the injured and
10 non-injured side for all objective measures. The mean difference was 8.1 degrees
11 (95% CI: 6.4 to 10.6) for ATRA, 2.9 cm (95% CI: 2.2 to 3.6) for HRH, and 1.6 cm
12 (95% CI: 1.3 to 1.9) for calf circumference (Table 3). For ME, the mean differences
13 were 2.7 cm (95% CI: 2.2 to 3.3) for average height, 5.0 (95% CI: 3.0 to 7.1) for
14 number of repetitions, and 807 newtons (95% CI: 643 to 972) for the total work
15 (Table 3). Regarding the objective measures and ME results for the Previous
16 regimen and New regimen groups, there were no differences (Table 4).

17

18 Table 3

19

20 Table 4

21

22 4. Discussion

23 The results of the present study demonstrate that patients struggle to fully recover
24 after an Achilles tendon rupture. For all objective measures, the injured side were
25 impaired compared to the non-injured side 4 years after an acute Achilles tendon

1 rupture. No difference in any outcome was observed between patients treated
2 according to the old regimen with surgery as first choice treatment compared with
3 the new regimen with functional rehabilitation as first choice treatment.

4
5 The results of this study support the results from previous studies evaluating 2- and
6 7-year follow-ups and underline that regaining full function following an Achilles
7 tendon rupture is difficult and only accomplished by few patients.^{23,24} Previous
8 studies comparing surgical and conservative treatment found no differences in ATRS
9 for different treatment strategies but observed a difference in favor of surgical
10 treatment in HRH and calf circumference.¹⁵

11
12 Previous research suggests that surgical treatment has a lower re-rupture rate than
13 non-surgical treatment, although the risk difference is low (1.6%).¹⁵ This small
14 difference found in previous studies¹⁵ indicates that clinical decision-making should
15 not be based exclusively on re-rupture. Surgical treatment has a higher risk of other
16 complications (3.3%), primarily due to a higher risk of infection. Functional outcome
17 have been reported in several studies,^{7,8,25} but to our knowledge, only two studies
18 have measured ATRS more than 4 years after rupture.^{24,26} In the present study,
19 patients had a mean ATRS of 71 at the 4-year follow-up. This is about 20 points
20 lower than the mean score reported by Brorsson et al.²⁴ at a 7-year follow-up. The
21 exact reason for this difference is unknown but may reflect differences in patient
22 populations. For example, Brorsson et al.²⁴ excluded patients older than 65 years
23 and patients with diabetes, neurovascular diseases, and other diseases; such
24 patients are likely to have poorer outcomes and were not excluded in the present
25 study. Another reason for the difference could be that the potential for improvement

1 continues even after 4 years. Brorsson et al. found no difference in any outcomes
2 between 2- and 7-year follow-ups, which indicates that improvement does not
3 continue after 2 years.²⁴

4

5 Similar to previous studies, we found no difference between treatment strategies
6 preferring surgical treatment and functional rehabilitation, respectively, measured by
7 ATRS, re-rupture rate, and objective measures at long-term follow-up (i.e., 4 years).

8 The data in re-rupture rate in this study support previous reports on low re-rupture
9 rates for both treatments¹⁵. This strengthens the existing evidence that the re-rupture
10 rate is low for both treatments.¹⁵

11

12 As severe comorbidity was considered a contraindication to surgery not all patients
13 underwent surgery in the period where surgery where surgery was the preferred
14 treatment. Therefore, it is not possible to compare surgery vs. non-surgery,
15 independent from regime, as it would result in significant bias. In the Previous
16 regimen group, only 63% were treated surgically. The reason for not performing
17 surgery despite this being the preferred strategy was mainly contraindications such
18 as comorbidities or low functional status, but could also be due to increased attention
19 to the topic in medical literature and at national and international conferences.

20 Implementation of non-surgical treatment at several hospitals close by might also
21 have made the surgeon more inclined to treat non-surgical instead of following the
22 guidelines. In the local guideline for New regimen it was described that patients with
23 open ruptures should be treated surgically. This might be the reason for 3 non-
24 included patients treated surgical as presented in the drop-out analysis (table 2) .

25 Comparing outcomes based on having surgery or not would introduce the risk of

1 confounding by indication. It is expected that the patients treated non-surgically in
2 Previous regimen are patients with comorbidities or other factors due to which
3 surgery would be a risk. Based on this, an analysis comparing treatments would
4 presumably result in incorrect high functional mean scores for patients treated
5 surgically.

6

7 There is a substantial number of studies examining whether patients should be
8 treated surgically or non-surgically. As this study supports, patients do not fully
9 recover at 4 years post injury. This raises the question of the importance of looking
10 at other parts of the treatment. There might not be a treatment that fits all patients,
11 and maybe the choice between surgery or non-surgery should be decided based on
12 patient preferences or objective measures. New research is examining whether
13 patients will have a better functional outcome if the choice between surgery or not is
14 based on the gap between the tendon ends measured immediately after rupture.²⁷

15

16 The functional deficits that patients experience in the lower leg affects the possibility
17 of returning to previous sport.^{19,28-30} Returning to physically demanding work is
18 substantially delayed.¹³ Professional athletes have to meet the high demands of
19 returning to high-level sports, and they also struggle to return to play, but some do
20 succeed.^{29,30} Anticipating that consistent adherence to rehabilitation and ongoing
21 training is part of the reason for athletes succeeding to return to highly demanding
22 sports, it will be relevant in future research to investigate the contents and the
23 potential of functional rehabilitation for both recreational and professional athletes.

24

25 **Limitations**

1 This study is a cohort study which have limitations compared to randomized
2 controlled trials and the quality of data depend on the data available from the
3 database.

4 The participation rate for the study was relatively low (40%), which could affect study
5 generalizability. The invitation was sent through a secure online mailing system,
6 which might have resulted in some patients not seeing the invitation. For ethical
7 reasons, it was not possible to call the patients who did not respond to the written
8 invitation.

9 Low participation rate could result in attrition bias. To examine if included patients
10 differed from those who did not show up at the 4-year follow-up, a dropout analysis
11 was performed. We found no difference between the included and non-included
12 patients in terms of age, sex, or previous ATRS (i.e., 1 year) (Table 2) suggesting
13 that we did not have a systematic loss to follow-up. For this study only data from the
14 DADB was available. As we only have data if the patients attended the follow up at
15 different timepoints, the missing data for ATRS at one and two years follow-up is
16 relative high. We could not provide information on re-ruptures from patients not
17 participating, as additional review of patient records require consent from the
18 patients. For data collected from a database, the quality of data is dependent on the
19 database, and due to changes in objective measures over time, comparing objective
20 measures over time was not possible. The treatment regimen differed in both time of
21 immobilization, type of cast and number of appointments with physiotherapist as the
22 two regimens where different, not only on whether they had surgery or not. As data
23 from time of rupture is from the database, we neither have information about specific
24 treatments for each patient or whether they supplemented the rehabilitation with

1 other treatments. Collecting this information at 4 years follow-up will have a high risk
2 of recall bias.

3 The difference between treatment strategies in ATRS score was 4.5 points (CI: -6.8
4 to 15.8). It is lower than the 7 points which is usually considered to the minimal
5 clinical important difference for the ATRS score³¹. As 95% CI expands over the 7
6 points we can not refute that there could potentially be a clinically important
7 difference between groups.

8

9 **Perspective**

10 This study contributes to previous findings that there is no difference between
11 treatment strategies for acute Achilles tendon ruptures. In future studies, it would be
12 relevant to add an economic analysis to expand the parameters based on which
13 treatment decisions are made. This analysis should include the cost of surgery and
14 rehabilitation.

15

16 **5. Conclusions**

17 In conclusion, we found that patients at an average age of 53 years still had patient
18 reported impairments and objective functional deficits 4 years after an acute Achilles
19 tendon rupture compared with their uninjured side. No differences in patient reported
20 outcome or objective measures at the 4 years follow-up were observed between two
21 consecutive cohorts with either a regimen with surgery as the preferred treatment
22 option compared with a regimen where functional rehabilitation (non-surgical
23 treatment) was the preferred strategy.

24

1 *Brief Summary*

2 What is already known

- 3 • Acute Achilles tendon rupture is a common injury and the incidence has
4 increased the last decades.
- 5 • Optimal treatment for acute Achilles tendon rupture is still uncertain.
- 6 • Patients have functional deficits 1-2 years after acute Achilles tendon rupture, but
7 long-term follow-up studies are scarce .

8

9 What this study adds

- 10 • Functional deficits persists 4 years after Achilles tendon rupture.
- 11 • Difference between injured and non-injured side in objective measures is
12 observed 4 years after an Acute Achilles tendon rupture.
- 13 • No improvements in functional outcomes was observed between 2 and 4 years
14 after an Acute Achilles tendon rupture.

15

16 **6. Acknowledgements**

17 The authors would like to thank Jette Kristiansen for assistance with correspondence
18 to all eligible patients.

1 7. References

- 2 1. Ganestam A, Kallemose T, Troelsen A, Barfod KW. Increasing incidence of
3 acute Achilles tendon rupture and a noticeable decline in surgical treatment
4 from 1994 to 2013. A nationwide registry study of 33,160 patients. *Knee
5 Surgery, Sport Traumatol Arthrosc.* 2016;24(12):3730–7.
- 6 2. Suchak AA, Bostick G, Reid D, Blitz S, Jomha N. The incidence of Achilles
7 tendon ruptures in Edmonton, Canada. *Foot Ankle Int.* 2005;26(11):932–6.
- 8 3. Sheth U, Wasserstein D, Jenkinson R, Moineddin R, Kreder H, Jaglal SB. The
9 epidemiology and trends in management of acute Achilles tendon ruptures in
10 Ontario, Canada a population-based study of 27 607 patients. *Bone Jt J.*
11 2017;99–B(1):78–86.
- 12 4. Lantto I, Heikkinen J, Flinkkilä T, Ohtonen P, Leppilahti J. Epidemiology of
13 Achilles tendon ruptures: Increasing incidence over a 33-year period. *Scand J
14 Med Sci Sport.* 2015;25(1):e133–8.
- 15 5. Huttunen TT, Kannus P, Rolf C, Felländer-Tsai L, Mattila VM. Acute achilles
16 tendon ruptures: Incidence of injury and surgery in Sweden between 2001 and
17 2012. *Am J Sports Med.* 2014;42(10):2419–23.
- 18 6. Egger AC, Berkowitz MJ. Achilles tendon injuries. *Curr Rev Musculoskelet
19 Med. Current Reviews in Musculoskeletal Medicine;* 2017;10(1):72–80.
- 20 7. Nilsson-Helander K, Grävare Silbernagel K, Thomeé R, Faxén E, Olsson N,
21 Eriksson BI, et al. Acute achilles tendon rupture: A randomized, controlled
22 study comparing surgical and nonsurgical treatments using validated outcome
23 measures. *Am J Sports Med.* 2010;38(11):2186–93.
- 24 8. Olsson, N., Nilsson-Helander, K., Karlsson, J., Eriksson, B. I., Thomée, R.,
25 Faxén, E., & Silbernagel, K. G. (2011). Major functional deficits persist 2 years

- 1 after acute Achilles tendon rupture. *Knee Surgery, Sports Traumatology,*
2 *Arthroscopy*, 19(8), 1385–1393. <https://doi.org/10.1007/s00167-011-1511-3>.
- 3 9. Deng S, Sun Z, Zhang C, Chen G, Li J. Surgical Treatment Versus
4 Conservative Management for Acute Achilles Tendon Rupture: A Systematic
5 Review and Meta-Analysis of Randomized Controlled Trials. *J Foot Ankle Surg*
6 [Internet]. Elsevier Ltd; 2017;56(6):1236–43. Available from:
7 <https://doi.org/10.1053/j.jfas.2017.05.036>
- 8 10. Wu Y, Lin L, Li H, Zhao Y, Liu L, Jia Z, et al. Is surgical intervention more
9 effective than non-surgical treatment for acute Achilles tendon rupture? A
10 systematic review of overlapping meta-analyses. *Int J Surg* [Internet]. Elsevier
11 Ltd; 2016;36:305–11. Available from:
12 <http://dx.doi.org/10.1016/j.ijisu.2016.11.014>
- 13 11. Erickson BJ, Mascarenhas R, Saltzman BM, Walton D, Lee S, Cole BJ, et al.
14 Operative vs. non-operative treatment of Achilles tendon ruptures: A
15 systematic review of overlapping meta-analyses. *Orthop J Sport Med*.
16 2015;3(7):1–11.
- 17 12. Bergkvist D, Åström I, Josefsson PO, Dahlberg LE. Acute achilles tendon
18 rupture: A questionnaire follow-up of 487 patients. *J Bone Jt Surg - Ser A*.
19 2012;94(13):1229–33.
- 20 13. Moller M, Movin T, Granhed H, Lind K, Faxen E, Karlsson J. Acute rupture of
21 tendon Achillis. A prospective randomised study of comparison between
22 surgical and non-surgical treatment. *J Bone Jt Surg* [Internet]. 2001;83(6):843–
23 8. Available from:
24 <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&do>
25 [pt=Citation&list_uids=11521926](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11521926)

- 1 14. Zhang H, Tang H, He Q, Wei Q, Tong D, Wang C, et al. Surgical Versus
2 Conservative Intervention for Acute Achilles Tendon Rupture. *Med (United*
3 *States)*. 2015;94(45):e1951.
- 4 15. Ochen Y, Beks RB, Van Heijl M, Hietbrink F, Leenen LPH, Van Der Velde D,
5 et al. Operative treatment versus nonoperative treatment of Achilles tendon
6 ruptures: Systematic review and meta-analysis. *BMJ*. 2019;364.
- 7 16. Twaddle BC, Poon P. Early motion for Achilles tendon ruptures: Is surgery
8 important? A randomized, prospective study. *Am J Sports Med*.
9 2007;35(12):2033–8.
- 10 17. Willits K, Amendola A, Bryant D, Mohtadi NG, Giffin JR, Fowler P, et al.
11 Operative versus Nonoperative Treatment of Acute Achilles Tendon Ruptures.
12 *J Bone Jt Surgery-American Vol [Internet]*. 2010;92(17):2767–75. Available
13 from: <https://insights.ovid.com/crossref?an=00004623-201012010-00001>
- 14 18. Soroceanu A, Sidhwa F, Aarabi S, Kaufman A, Glazebrook M. Surgical Versus
15 Nonsurgical Treatment. *J Bone Joint Surg Am*. 2012;94(23):2136–43.
- 16 19. Hansen M.S., Christensen M., Budolfsen T., Østergaard T.F., Kallermose T.,
17 Troelsen A., et. al.: Achilles tendon total rupture score at 3 months can predict
18 patients' ability to return to sport 1 year after injury. *Knee Surg Sport Traumatol*
19 *Arthrosc* 2016; 24: pp. 1365-1371.
- 20 20. Nilsson-Helander K, Thomeé R, Grävare-Silbernagel K, Thomeé P, Faxén E,
21 Eriksson BI, et al. The Achilles tendon Total Rupture Score (ATRS):
22 Development and validation. *Am J Sports Med*. 2007;35(3):421–6.
- 23 21. Carmont MR, Silbernagel KG, Mathy A, Mulji Y, Karlsson J, Maffulli N.
24 Reliability of Achilles Tendon Resting Angle and Calf Circumference
25 measurement techniques. *Foot Ankle Surg*. 2013;19(4):245–9.

- 1 22. Silbernagel KG, Gustavsson A, Thomeé R, Karlsson J. Evaluation of lower leg
2 function in patients with Achilles tendinopathy. *Knee Surgery, Sport Traumatol*
3 *Arthrosc.* 2006;14(11):1207–17.
- 4 23. Olsson N, Nilsson-Helander K, Karlsson J, Eriksson BI, Thomée R, Faxén E,
5 et al. Major functional deficits persist 2 years after acute Achilles tendon
6 rupture. *Knee Surgery, Sport Traumatol Arthrosc.* 2011;19(8):1385–93.
- 7 24. Brorsson A, Grävare Silbernagel K, Olsson N, Nilsson Helander K. Calf Muscle
8 Performance Deficits Remain 7 Years After an Achilles Tendon Rupture. *Am J*
9 *Sports Med.* 2018;46(2):470–7.
- 10 25. Jackson G, Sinclair VF, McLaughlin C, Barrie J. Outcomes of Functional
11 Weight-bearing Rehabilitation of Achilles Tendon Ruptures. *Orthopedics.*
12 2014;36(8):e1053–9.
- 13 26. Lim CS, Lees D, Gwynne-Jones DP. Functional Outcome of Acute Achilles
14 Tendon Rupture With and Without Operative Treatment Using Identical
15 Functional Bracing Protocol. *Foot Ankle Int.* 2017;38(12):1331–6. 27.
- 16 27. Amlang MH, Rammelt S, Peaden A, Zwipp H, Friedrich A, Bunk A.
17 Ultrasonographic Classification of Achilles Tendon Ruptures as a Rationale for
18 Individual Treatment Selection. *ISRN Orthop.* 2011;2011:1–10.
- 19 28. Carmont MR, Grävare Silbernagel K, Brorsson A, Olsson N, Maffulli N,
20 Karlsson J. The Achilles tendon resting angle as an indirect measure of
21 Achilles tendon length following rupture, repair, and rehabilitation. *Asia-Pacific*
22 *J Sport Med Arthrosc Rehabil Technol.* 2015;2(2):49–55.
- 23 29. Trofa DP, Miller JC, Jang ES, Woode DR, Greisberg JK, Vosseller JT.
24 Professional Athletes' Return to Play and Performance after Operative Repair
25 of an Achilles Tendon Rupture. *Am J Sports Med [Internet].* 2017;45(12):2864–

- 1 71. Available from: <https://doi.org/10.1177/0363546517713001>
- 2 30. Trofa DP, Noback PC, Caldwell JME, Miller JC, Greisberg JK, Ahmad CS, et
3 al. Professional Soccer Players' Return to Play and Performance After
4 Operative Repair of Achilles Tendon Rupture. *Orthop J Sport Med.*
5 2018;6(11):1–7.
- 6 31. Ganestam A, Barfod K, Klit J, Troelsen A. Validity and reliability of the Achilles
7 tendon total rupture score. *J Foot Ankle Surg [Internet].* 2013;52(6):736–9.
8 Available from: <http://dx.doi.org/10.1053/j.jfas.2013.07.004>

9

10

11 **Tables**12 *Table 1*13 **Table 1** Patient characteristics at the 4-year follow-up (n=76)

	All patients (n=76)	Previous regimen (n=35)	New regimen (n=41)	Difference	P-value
Age (years)	53.7 ± 12.5 (25.8-81.3)	53.7 ± 12.6 (25.7-81.3)	53.7 ± 12.6 (27.4-76.7)	0.0	0.994
Height (cm)	177 ± 7 (154-196)	178 ± 8 (160-196)	177 ± 7 (154-192)	1.4	0.387
Weight (kg)	82 ± 14 (52-134)	82 ± 14 (52-110)	83 ± 14 (63-134)	-0.6	0.859
BMI (kg/m ²)	26.2 ± 4.3 (18.7-45.8)	25.8 ± 3.7 (18.7-32.8)	26.5 ± 4.8 (21.4-45.8)	-0.8	0.435

Sex					
Female	22 (29%)	8 (23%)	14 (34%)	-11%	0.279
Male	54 (71%)	27 (77%)	27 (66%)		
Time since rupture (years)					
	4.3 ± 0.8 (3.0-5.9)	5.0 ± 0.5 (4.3-5.9)	3.7 ± 0.4 (3.1-4.2)	1.4	<0.001
Treatment					
Surgical	22 (29%)	22 (63%)	0 (0%)	63%	<0.001
Non-surgical	54 (71%)	13 (37%)	41 (100%)		
Injured side					
Right	31 (41%)	16 (46%)	15 (37%)	9%	0.420
Left	45 (59%)	19 (54%)	26 (63%)		
Dominant side					
Yes	31 (50%)	11 (52%)	20 (49%)	3%	0.788
No	31 (50%) ^b	10 (48%) ^c	21 (51%)		
Pre-ATRS score ^a					
Mean	95 ± 13 ^d	98 ± 4 ^e	93 ± 17	5.2	0.081
Median	100 (25-100)	100 (84-100)	100 (25-100)	0	0.359
4 years ATRS score					
Mean	71.4 ± 27	73.9 ± 22	69.4 ± 27	4.5	0.432
Median	79.5 (9-100)	79 (20-100)	80 (9-100)	0.5	0.628

Continuous variables are presented with mean, standard deviation and min-max values.

ATRS are presented with with mean, standard deviation and median and min-max values

Dichotomous variables are presented as numbers and percentage.

^aThe pre-ATRS score was assessed at inclusion and not at 4 years follow-up. ^bn=62, ^cn=21, ^dn=75, ^en=34.

1 *Table 2*

2 **Table 2** Treatment, ATRS-scores at 1 and 2 years follow-up and health data for
 3 included and non-included patients (n=192)

	Included (n=76)	Non-included (n=116)	P-value
<i>Surgical vs. non-surgical</i>			
Treatment Previous regimen			
Surgical	22(63%)	27 (44%)	0.068
Non-surgical	13 (37%)	36 (56%)	
Treatment New regimen			
Surgical	0 (0%)	3 (6%)	0.122
Non-surgical	41 (100%)	50 (94%)	
<i>ATRS and health data</i>			
Age	53.7 ± 12.5 ^a (25.8-81.3)	50.9 ± 13.5 ^b (26.3-87.3)	0.149
Sex			
Female	22 (29%)	23 (21%)	0.234
Male	54 (71%)	85 (79%)	
Pre-ATRS			
Mean	95 ± 13 ^c	92 ± 17 ^d	0.197
Median	100 (25-100)	100 (25-100)	0.693
ATRS at 1-year follow-up			
Mean	65 ± 23 ^e	58 ± 27 ^f	0.125
Median	78.5 (12-100)	64 (5-99)	0.159
ATRS at 2-year follow-up			
Mean	66 ± 25 ^g	68 ± 27 ^h	0.681
Median	67 (8-100)	70.5 (2-100)	0.498
Diabetes			
Yes	2 (3%)	6 (5%)	0.389

No	74 (97%)	110 (95%)	
Smoker			
Yes	9 (12%)	26 (22%)	0.064
No	67 (88%)	90 (78%)	

Continuous variables are presented with mean, standard deviation and min-max values.

ATRS are presented with with mean, standard deviation, median and min-max values

^an=76, ^bn=116, ^cn=75, ^dn=110, ^en=63, ^fn=73, ^gn=59, ^hn=60.

1 *Table 3*2 **Table 3** Objective measures at the 4-year follow-up (n=76)

	Injured side (n=76)	Non-injured side (n=76)	Difference	P-value
ATRA (degrees)	54.9 (53.2 to 56.5)	46.7 (45.0 to 48.4)	-8.1 ^r (6.4 to 9.8)	<0.001
Heel-rise height (cm)	8.8 (7.8 to 9.7)	11.6 (10.9 to 12.4)	2.9 (2.2 to 3.6)	<0.001
Calf circumference (cm)	36.1 (35.3 to 36.9)	37.7 (36.9 to 38.5)	1.6 (1.3 to 1.9)	<0.001
Muscle endurance				
Avg. height (cm)	7.9 (7.1 to 8.7)	10.7 (10.1 to 11.3)	2.8 (2.2 to 3.3)	<0.001
Repetitions (n)	21.5 (18.6 to 24.5)	26.5 (23.6 to 29.4)	5.0 (3.0 to 7.1)	<0.001
Total work (joule)	1412 (1202 to 1623)	2219 (1996 to 2443)	807 (643 to 972)	<0.001

Values are reported as mean with 95% confidence intervals.

Difference is reported as mean difference between injured and non-injured side with 95% confidence intervals.

^rDifference do not correspond with difference in injured side and non-injured side because of rounding.

1 **Table 4**2 **Table 4** Difference in objective measures between groups at 4 years follow-up (Limb

3 Symmetry Index) (n=76)

	Previous regimen (n=35)	New regimen (n=41)	Difference	P-value
ATRA (degrees)	117.7% (111.4 to 124.0)	120.6% (115.0 to 126.3)	-2.9% (-11.2 to 5.4)	0.488
Heel-rise height (cm)	78.9% (69.8 to 88.1)	69.3% (59.8 to 78.8)	9.7% ^r (-3.4 to 22.8)	0.146
Calf circumference (cm)	96.0% (94.9 to 97.1)	95.6% (94.4 to 96.7)	0.5% ^r (-1.1 to 2.0)	0.555
Muscle endurance				
Avg. height (cm)	76.7% (68.8 to 84.6)	69.2% (60.2 to 78.1)	7.5% (-4.4 to 19.5)	0.219
Repetitions (n)	87.1% (74.8 to 99.3)	72.2% (61.8 to 82.6)	14.8% ^r (-0.8 to 30.5)	0.063
Total work (joule)	69.4% (56.8 to 82.1)	54.9% (45.3 to 64.5)	14.5% (-0.9 to 29.9)	0.064

Values are reported as percentage difference between injured and non-injured side (Limb Symmetry Index) with 95% confidence intervals.

Values above 100 indicate highest mean score for injured side and values below 100 indicate highest score for non-injured side.

^rDifference do not correspond with difference in Previous regimen and New regimen values because of rounding.

1 **Figures**

2 *Figure 1*

3 **Figure 1** Flow-chart of recruiting (n=222)

4

5 *Figure 2*

6 **Figure 2** Mean ATRS-scores with 95% confidence intervals at different time

7 points.(n=76)

8 Legend: Mean scores with 95% confidence intervals. Time x group interaction,

9 $p=0.8509$