



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Long-term cognitive and pulmonary functions following a lower versus a higher oxygenation target in the HOT-ICU and HOT-COVID trials: A protocol update

Crescioli, Elena; Riis, Jens Østergaard; Weinreich, Ulla Møller; Jensen, Jens Ulrik Staehr; Poulsen, Lone Musaeus; Brøchner, Anne Craveiro; Lange, Theis; Perner, Anders; Klitgaard, Thomas Lass; Schjørring, Olav Lilleholt; Rasmussen, Bodil Steen

Published in:
Acta Anaesthesiologica Scandinavica

DOI (link to publication from Publisher):
[10.1111/aas.14379](https://doi.org/10.1111/aas.14379)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2024

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Crescioli, E., Riis, J. Ø., Weinreich, U. M., Jensen, J. U. S., Poulsen, L. M., Brøchner, A. C., Lange, T., Perner, A., Klitgaard, T. L., Schjørring, O. L., & Rasmussen, B. S. (2024). Long-term cognitive and pulmonary functions following a lower versus a higher oxygenation target in the HOT-ICU and HOT-COVID trials: A protocol update. *Acta Anaesthesiologica Scandinavica*, 68(4), 575-578. <https://doi.org/10.1111/aas.14379>




General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

RESEARCH ARTICLE

Long-term cognitive and pulmonary functions following a lower versus a higher oxygenation target in the HOT-ICU and HOT-COVID trials: A protocol update

Elena Crescioli^{1,2}  | Jens Østergaard Riis³ | Ulla Møller Weinreich^{2,4} |
Jens Ulrik Stæhr Jensen⁵ | Lone Musaeus Poulsen⁶ | Anne Craveiro Brøchner⁷ |
Theis Lange⁸ | Anders Perner⁹ | Thomas Lass Klitgaard¹ |
Olav Lilleholt Schjørring^{1,2}  | Bodil Steen Rasmussen^{1,2} 

¹Department of Anaesthesia and Intensive Care, Aalborg University Hospital, Aalborg, Denmark

²Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

³Department of Neurology and Neurosurgery, Aalborg University Hospital, Aalborg, Denmark

⁴Department of Respiratory Medicine, Aalborg University Hospital, Aalborg, Denmark

⁵Section of Respiratory Medicine, Department of Medicine, Herlev and Gentofte Hospital, University of Copenhagen, Hellerup, Denmark

⁶Department of Anaesthesiology, Zealand University Hospital, Køge, Denmark

⁷Department of Anaesthesia and Intensive Care, Kolding Hospital, University of Southern Denmark, Kolding, Denmark

⁸Section of Biostatistics, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

⁹Department of Intensive Care, Rigshospitalet, Copenhagen, Denmark

Correspondence

Elena Crescioli, Department of Anaesthesia and Intensive Care, Aalborg University Hospital, Hobrovej 18-22, 9100 Aalborg, Denmark.

Email: e.crescioli@rn.dk

Funding information

Novo Nordisk Foundation, Grant/Award Number: NNF18OC0034614; Danish Ministry of Higher Education and Research, Grant/Award Number: 0238-00004B

Abstract

Background: The Handling Oxygenation Targets in the Intensive Care Unit (HOT-ICU) trial was a multicentre, randomised, parallel-group trial of a lower oxygenation target (arterial partial pressure of oxygen [PaO₂] = 8 kPa) versus a higher oxygenation target (PaO₂ = 12 kPa) in adult ICU patients with acute hypoxaemic respiratory failure; the Handling Oxygenation Targets in coronavirus disease 2019 (HOT-COVID) tested the same oxygenation targets in patients with confirmed COVID-19. In this study, we aim to evaluate the long-term effects of these oxygenation targets on cognitive and pulmonary function. We hypothesise that a lower oxygenation target throughout the ICU stay may result in cognitive impairment, whereas a higher oxygenation target may result in impaired pulmonary function.

Methods: This is the updated protocol and statistical analysis plan of two pre-planned secondary outcomes, the long-term cognitive function, and long-term pulmonary function, in the HOT-ICU and HOT-COVID trials. Patients enrolled in both trials at selected Danish sites and surviving to 1 year after randomisation are eligible to participate. A Repeatable Battery for the Assessment of Neuropsychological Status score and a full-body plethysmography, including diffusion capacity for carbon monoxide, will be obtained. The last patient is expected to be included in the spring of 2024.

Conclusion: This study will provide important information on the long-term effects of a lower versus a higher oxygenation target on long-term cognitive and pulmonary functions in adult ICU patients with acute hypoxaemic respiratory failure.

KEYWORDS

clinical trial, critical care outcomes, ICU, oxygen inhalation therapy

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *Acta Anaesthesiologica Scandinavica* published by John Wiley & Sons Ltd on behalf of Acta Anaesthesiologica Scandinavica Foundation.

1 | INTRODUCTION

Cognitive impairment and pulmonary dysfunction have been described as longstanding complications in intensive care unit (ICU) survivors.¹ A low arterial partial pressure of oxygen (PaO₂) has been associated with poor cognitive performance,² while concerns about higher oxygenation targets being possibly disadvantageous in terms of pulmonary function have been raised.³ However, the evidence on the long-term effects of oxygen therapy in adult ICU survivors is still very uncertain.⁴

The Handling Oxygenation Targets in the Intensive Care Unit (HOT-ICU) trial investigated a lower versus a higher oxygenation target in adult patients acutely admitted to ICU with hypoxaemic respiratory failure,⁵ and the Handling Oxygenation Targets in coronavirus disease 2019 (HOT-COVID) trial tested similar interventions in hypoxaemic patients with a confirmed COVID-19 infection.⁶ In the present study, we aim to evaluate the effects of the two oxygenation targets on cognitive and pulmonary functions in survivors 1 year after randomisation in the HOT-ICU or HOT-COVID trials. We hypothesise that the lower oxygenation target results in long-term cognitive impairment, whereas the higher oxygenation target results in impaired long-term pulmonary function.

2 | METHODS

2.1 | Study design

This is the updated protocol and statistical analysis plan for a sub-study of the HOT-ICU and HOT-COVID trials evaluating the two pre-planned secondary long-term outcomes; cognitive function and pulmonary function,⁵⁻⁷ 1 year after randomisation in survivors included at selected Danish sites. This is an amendment to the previously published protocol for these outcomes in the HOT-ICU trial.⁸ In the following, we will only consider the elements that have been changed from the primary version of the protocol.⁸

This protocol amendment was written in accordance with the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) 2013 statement.⁹ An updated SPIRIT 2013 checklist is presented in Appendix S1.

2.2 | The Danish cohort of the HOT-ICU and HOT-COVID trials

The HOT-ICU and HOT-COVID trials were both investigator-initiated, pragmatic, multi-centre, randomised, parallel-group trials of a lower versus a higher oxygenation target in adult patients acutely admitted to the ICU with severe hypoxaemic respiratory failure, with outcome-assessor blinding of the present long-term cognitive and pulmonary outcomes. Inclusion criteria defining severe hypoxaemic respiratory failure were provision of at least 10 L of oxygen per minute in an open system in both trials, or a fraction of inspired oxygen (FiO₂) of at least

0.50 in a closed system in the HOT-ICU trial, but with no limits for FiO₂ in closed systems in the HOT-COVID trial.^{5,6} Additionally, the HOT-COVID trial required a confirmed severe acute respiratory syndrome coronavirus 2 infection.⁶ Full lists and definitions of inclusion and exclusion criteria can be found elsewhere.⁵⁻⁷ Patients were randomised 1:1 within 12 h of ICU admission to either a PaO₂ target of 8 kPa (lower oxygenation target) or 12 kPa (higher oxygenation target) during ICU admission up to a maximum of 90 days after randomisation, including ICU readmissions. The HOT-ICU trial was performed in 35 ICUs across 7 countries, and the HOT-COVID trial was performed in 11 ICUs across 3 countries. The Danish trial cohorts consist of 2332 patients in the HOT-ICU trial, and of 593 patients in the HOT-COVID trial, respectively.

2.3 | Approvals and registrations

The HOT-COVID trial was approved as an amendment to the HOT-ICU trial. Thus, approvals for both trials are identical as follows: from the Danish Health and Medicines Agency (AAUH-ICU-01), the Committee on Health Research Ethics in the North Denmark Region (N-20170015), the Danish Data Protection Agency (2017-055), all required authorities in participating countries, and the European clinical trials database (EudraCT number 2017-000632-34). Both trials were prospectively registered at [Clinicaltrials.gov](https://clinicaltrials.gov) (Identifier: NCT03174002 for the HOT-ICU trial; NCT04425031 for the HOT-COVID trial).^{6,7} Additional details on both trials are available elsewhere.⁵⁻⁷

2.4 | Study population and 1-year cognitive and pulmonary follow-up

Patients enrolled in the HOT-ICU and HOT-COVID trials at selected Danish sites and surviving until 1 year after randomisation are eligible to participate. The inclusion criteria are the following: (1) included in the HOT-ICU or HOT-COVID trials; (2) able to speak and understand the Danish language; (3) informed consent to participate in the long-term evaluations of cognitive and lung functions. Patients will be excluded from the study if they meet any of the following criteria: (1) more than 20 months since inclusion; (2) consent not obtainable according to national regulations; (3) body weight above 150 kg (only an exclusion criterion for the lung function test).

2.5 | Outcome measures

The two pre-planned long-term outcomes covered in this study are:

1. A global cognitive score obtained from the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS),¹⁰ which is a comprehensive test battery for the neuropsychological status consisting of 12 subtests designed to produce a global cognitive score and index scores for five different cognitive

domains (immediate memory; delayed memory; attention; language; and visuospatial/constructional abilities).

2. A full-body plethysmography with measurement of pulmonary diffusion capacity for carbon monoxide as the inactive tracer gas mixed with oxygen and nitrogen. The value obtained corresponds to the diffusing capacity for carbon monoxide (DLCO).¹¹

Exploratory outcomes include the five cognitive domains of the RBANS, and the following measurements of dynamic and static lung function: (a) forced expiratory volume in the first second (FEV₁), (b) forced vital capacity (FVC), (c) the ratio between FEV₁ and FVC (FEV₁/FVC), (d) total lung capacity (TLC), (e) inspiratory capacity (IC), (f) ratio between IC and TLC (IC/TLC), (g) residual volume, and (h) intrathoracic gas volume.

2.6 | Statistics

We will conduct all analyses according to the intention-to-treat principle among 1-year survivors unless specified otherwise.¹² The intention-to-treat population includes all randomised Danish patients at selected sites surviving to 1 year after randomisation, except where follow-up data cannot be obtained due to loss to follow-up, for example, due to withdrawal of consent according to national regulations.^{13–15} Individual patient-level data from the two trials will be appended and analysed in a one-step approach using a linear mixed model, accounting for clustering of patients within the trials and sites. Specifically, we will include a random effect for trial and sites. We will not adjust for other stratification variables as these were not similar in the two trials. We will use a robust estimator of clustered variance–covariance¹⁶; preferably a VCE(robust)¹⁷ option in Stata. However, if more than 20 iterations are required to obtain convergence, a VCE(cluster)¹⁷ will be used instead to avoid overfitting. Additionally, we will perform supplemental analyses of all outcomes in each trial separately using a linear regression adjusted for each trial's stratification variables (site, presence or absence of chronic obstructive pulmonary disease, and presence or absence of active haematological malignancy in the HOT-ICU trial; site in the HOT-COVID trial) and a robust estimator of clustered variance–covariance, as described above. All results will be presented as mean differences with 95% confidence interval and a *p*-value below 0.05 will be considered statistically significant. All statistical analyses will be conducted using Stata version 17, STATA Nordic.

We will report the RBANS scores (comprising the global cognitive score and the five index scores) in each intervention group, stratified by the age-adjusted mean and standard deviation in cognitive test performance (above normal: >115 points; normal: between 85 and 115 points; mild impairment: between 70 and 84 points; moderate impairment: between 55 and 69 points; severe impairment: <54 points)¹⁰; DLCO will be presented as the percentage of the predicted values in each intervention group categorised by severity (normal:> 80%; mildly impaired: from 60% to 80%; moderately impaired: from 40% to 59%; severely impaired:<40%).¹⁸ Finally, to quantify the proportions of patients with obstructive or restrictive

patterns in the exploratory pulmonary function outcomes, we will report proportions of patients with an FEV₁/FVC ratio below 70% as indicative of obstructive pulmonary disease,¹⁹ and an FVC below 80% or a TLC below 90% as indicative of restrictive pulmonary disease.^{20,21}

2.7 | Evaluation of sample representativeness

In order to evaluate if our sample is representative of the entire trial population alive at 1 year, we will compare the baseline variables and non-mortality outcomes at 90 days (i.e., serious adverse events in the ICU, days alive without life-support in the ICU, days alive and out of hospital), and 1 year (i.e., EuroQol visual analogue scale score)²² between the subpopulation included in this study and the remaining combined trial cohort alive at 1 year from randomisation.^{6,7} We will use a chi-squared test for categorical data and a parametric or non-parametric test for continuous data, as appropriate.

3 | DISCUSSION

This 1-year follow-up sub-study of the HOT-ICU and HOT-COVID trials will provide new important information on cognitive and pulmonary long-term outcomes following lower versus higher oxygenation targets in acutely ill patients with hypoxaemic respiratory failure admitted to the ICU. Thus far, only one randomised clinical trial (RCT) investigating two oxygenation strategies has performed cognitive follow-up at 180 days post-randomisation,²³ whereas long-term pulmonary function has never been studied in RCTs exploring lower versus higher oxygenation strategies, except in a very small COPD population.²⁴ Since the COVID-19 pandemic has significantly affected the 1-year follow-up inclusion rate of the HOT-ICU trial, we decided to analyse and present the 1-year pulmonary and cognitive outcomes for both trials in combination.

4 | STATUS

The 1-year follow-up of the HOT-ICU trial was completed in October 2021 with a total of 216 patients included in this sub-study. The 1-year follow-up of the HOT-COVID trial is currently ongoing and the last patient is expected to be assessed during spring 2024.

AUTHOR CONTRIBUTIONS

EC and BSR wrote the first draft of the article. All authors made substantial contributions to the article and provided important scientific input and read plus approved the final version of the article.

ACKNOWLEDGEMENTS

We thank all participating patients and relatives, and the participating clinical research staff at Aalborg University Hospital, Kolding Hospital, Zealand University Hospital Køge, and Herlev & Gentofte Hospital.

FUNDING INFORMATION

This study is funded by the Novo Nordisk Foundation (NNF18OC0034614) and the Danish Ministry of Higher Education and Research (0238-00004B), who had no influence on study design, scientific content of the protocol, trial conduct, data collection or analysis, interpretation of the results, or on the final publication.

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study will be available from the corresponding author upon reasonable request.

ORCID

Elena Crescioli  <https://orcid.org/0000-0002-8267-7634>

Olav Lilleholt Schjørring  <https://orcid.org/0000-0002-7749-6003>

Bodil Steen Rasmussen  <https://orcid.org/0000-0003-2190-145X>

REFERENCES

- Desai SV, Law TJ, Needham DM. Long-term complications of critical care. *Crit Care Med*. 2011;39(2):371-379. doi:10.1097/CCM.0b013e3181fd66e5
- Mikkelsen ME, Christie JD, Lanke PN, et al. The adult respiratory distress syndrome cognitive outcomes study: long-term neuropsychological function in survivors of acute lung injury. *Am J Respir Crit Care Med*. 2012;185(12):1307-1315. doi:10.1164/rccm.201111-2025OC
- Mikkelsen ME, Anderson B, Christie JD, Hopkins RO, Lanke PN. Can we optimize long-term outcomes in acute respiratory distress syndrome by targeting normoxemia? *Ann Am Thorac Soc*. 2014;11(4):613-618. doi:10.1513/AnnalsATS.201401-001PS
- Crescioli E, Krejberg KU, Klitgaard TL, et al. Long-term effects of lower versus higher oxygenation levels in adult ICU patients—a systematic review. *Acta Anaesthesiol Scand*. 2022;66(8):910-922. doi:10.1111/aas.14107
- Schjørring OL, Klitgaard TL, Perner A, et al. Lower or higher oxygenation targets for acute hypoxemic respiratory failure. *N Engl J Med*. 2021;384(14):1301-1311. doi:10.1056/NEJMoa2032510
- Mølgaard Nielsen F, Lass Klitgaard T, Crescioli E, et al. Handling oxygenation targets in ICU patients with COVID-19-protocol and statistical analysis plan in the HOT-COVID trial. *Acta Anaesthesiol Scand*. 2021;65(10):1497-1504. doi:10.1111/aas.13956
- Schjørring OL, Perner A, Wetterslev J, et al. Handling Oxygenation Targets in the Intensive Care Unit (HOT-ICU)-protocol for a randomised clinical trial comparing a lower vs a higher oxygenation target in adults with acute hypoxaemic respiratory failure. *Acta Anaesthesiol Scand*. 2019;63(7):956-965. doi:10.1111/aas.13356
- Crescioli E, Riis J, Weinreich UM, et al. Long-term cognitive and pulmonary functions following a lower versus a higher oxygenation target in the HOT-ICU trial: protocol and statistical analysis plan. *Acta Anaesthesiol Scand*. 2022;66(2):282-287. doi:10.1111/aas.13995
- Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med*. 2013;158(3):200-207. doi:10.7326/0003-4819-158-3-201302050-00583
- Randolph C, Tierney MC, Mohr E, Chase TN. The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): preliminary clinical validity. *J Clin Exp Neuropsychol*. 1998;20(3):310-319. doi:10.1076/jcen.20.3.310.823
- DR Borland C, B Hughes JM. Lung diffusing capacities (D(L)) for nitric oxide (NO) and carbon monoxide (CO): the evolving story. *Compr Physiol*. 2019;10(1):73-97. doi:10.1002/cphy.c190001
- DeMets DL, Cook T. Challenges of non-intention-to-treat analyses. *Jama*. 2019;321(2):145-146. doi:10.1001/jama.2018.19192
- International conference on harmonisation; guidance on statistical principles for clinical trials; availability – FDA. Notice. *Fed Regist*. 1998;63(179):49583-49598. International conference on harmonisation; guidance on statistical principles for clinical trials; availability—FDA. Notice.
- Gabriel SE, Normand SL. Getting the methods right—the foundation of patient-centered outcomes research. *N Engl J Med*. 2012;367(9):787-790. doi:10.1056/NEJMp1207437
- Schmiemann G. The preliminary draft of the methodology report by the Patient-Centered Outcomes Research Institute. *Z Evid Fortbild Qual Gesundheitswes*. 2012;106(7):496-499. Der vorläufige Methodenbericht des Patient-Centered Outcomes Research Institute. doi:10.1016/j.zefq.2012.07.002
- Cameron AC, Douglas LM. A Practitioner's guide to cluster-robust inference. *Journal of Human Resources*. 2015;50(2):317-372. doi:10.3368/jhr.50.2.317
- vce options. Variance estimators. Accessed October 31, 2023. https://www.stata.com/manuals/xtvce_options.pdf
- Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J*. 2005;26(5):948-968. doi:10.1183/09031936.05.00035205
- Vogelmeier CF, Criner GJ, Martinez FJ, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report. GOLD executive summary. *Am J Respir Crit Care Med*. 2017;195(5):557-582. doi:10.1164/rccm.201701-0218PP
- Aggarwal AN, Aggarwal R. The new ATS/ERS guidelines for assessing the spirometric severity of restrictive lung disease differ from previous standards. *Respirology*. 2007;12(5):759-762. doi:10.1111/j.1440-1843.2007.01117.x
- Hariri LP, Smith ML, Mino-Kenudson M, et al. Pulmonary pathology perspective on the 2018 American Thoracic Society, European Respiratory Society, Japanese Respiratory Society, and Latin American Thoracic Society idiopathic pulmonary fibrosis clinical practice guidelines. *Ann Am Thorac Soc*. 2020;17(5):550-554. doi:10.1513/AnnalsATS.201910-801PS
- Herdman M, Gudex C, Lloyd A, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res*. 2011;20(10):1727-1736. doi:10.1007/s11136-011-9903-x
- Mackle D, Bellomo R, Bailey M, et al. Conservative oxygen therapy during mechanical ventilation in the ICU. *N Engl J Med*. 2020;382(11):989-998. doi:10.1056/NEJMoa1903297
- Gomersall CD, Joynt GM, Freebairn RC, Lai CK, Oh TE. Oxygen therapy for hypercapnic patients with chronic obstructive pulmonary disease and acute respiratory failure: a randomized, controlled pilot study. *Crit Care Med*. 2002;30(1):113-116. doi:10.1097/00003246-200201000-00018

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Crescioli E, Riis JØ, Weinreich UM, et al. Long-term cognitive and pulmonary functions following a lower versus a higher oxygenation target in the HOT-ICU and HOT-COVID trials: A protocol update. *Acta Anaesthesiol Scand*. 2024;68(4):575-578. doi:10.1111/aas.14379