



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Biochemical Changes in Adult Male Gamers During Prolonged Gaming: Pilot Study

Krarp, Kasper Bygum; Riis, Johannes; Mørk, Morten; Nguyen, Hien Thi Thu; Søkilde Pedersen, Inge; Risom Kristensen, Søren; Handberg, Aase; Krarp, Henrik Bygum

Published in:
Interactive journal of medical research

DOI (link to publication from Publisher):
[10.2196/46570](https://doi.org/10.2196/46570)

Creative Commons License
CC BY 4.0

Publication date:
2024

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Krarp, K. B., Riis, J., Mørk, M., Nguyen, H. T. T., Søkilde Pedersen, I., Risom Kristensen, S., Handberg, A., & Krarp, H. B. (2024). Biochemical Changes in Adult Male Gamers During Prolonged Gaming: Pilot Study. *Interactive journal of medical research*, 13, Article e46570. <https://doi.org/10.2196/46570>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Original Paper

Biochemical Changes in Adult Male Gamers During Prolonged Gaming: Pilot Study

Kasper Bygum Krarup¹, BSc, MD; Johannes Riis¹, MD; Morten Mørk², MD, PhD; Hien Thi Thu Nguyen³, MSc, PhD; Inge Søkilde Pedersen^{3,4}, MSc, PhD; Søren Risom Kristensen^{2,4}, MD, PhD; Aase Handberg^{2,4}, MD, PhD; Henrik Bygum Krarup^{3,4}, MD, PhD

¹Department of Geriatrics, Aalborg University Hospital, Aalborg, Denmark

²Department of Clinical Biochemistry, Aalborg University Hospital, Aalborg, Denmark

³Department of Molecular Diagnostics, Aalborg University Hospital, Aalborg, Denmark

⁴Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

Corresponding Author:

Kasper Bygum Krarup, BSc, MD

Department of Geriatrics

Aalborg University Hospital

Reberbansgade 15

Aalborg, 9000

Denmark

Phone: 45 61781289

Email: kasper.krarup@rn.dk

Abstract

Background: Gaming has become an integrated part of life for children and adults worldwide. Previous studies on the impact of gaming on biochemical parameters have primarily addressed the acute effects of gaming. The literature is limited, and the study designs are very diverse. The parameters that have been investigated most thoroughly are blood glucose and cortisol.

Objective: This exploratory study is the first to investigate the effects of long gaming sessions on the biochemical parameters of healthy male adults. The extensive testing allowed us to observe short-term changes (within 6 hours), long-term changes during the duration of the gaming sessions, and follow-up after 1 week to determine whether any changes were longer lasting.

Methods: In total, 9 experienced gamers completed 2 back-to-back 18-hour gaming sessions interspersed with a 6-hour rest period. All participants adhered to a structured sleep pattern due to daytime employment or attending university. Blood, saliva, and urine samples were collected from the participants every 6 hours. Linear mixed-effect models were used to analyze the repeated-measures data accumulated during the study. A total of 51 biochemical parameters were investigated.

Results: In total, 12 of the 51 biochemical parameters significantly changed during the study: alkaline phosphatase, aspartate aminotransferase, bilirubin, chloride, creatinine, glucose, hemoglobin, immature reticulocyte fraction, lactate, methemoglobin, sodium, and thrombocytes. All changes were within the normal range. The mean glucose level of the participants was 4.39 (SD 0.07) mmol/L at baseline, which increased significantly by 0.24 (SD 0.07) mmol/L per 6 hours during the first period and by 0.38 (SD 0.07) mmol/L per 6 hours in the second period ($P < .001$). The glucose levels during the second session increased even though the participants had little energy intake. Cortisol levels did not change significantly, although the cortisol pattern deviated from the typical circadian rhythm. During both gaming sessions, we observed increasing cortisol levels from 6 AM until noon. The participants were relatively dehydrated at the start of the study. The patients were asked to fast before the first blood sampling. Within the first 6 hours of the study, the participants rehydrated, followed by relative dehydration during the remainder of the study. This pattern was identified using the following parameters: albumin, creatinine, hemoglobin, erythrocytes, potassium, and platelets.

Conclusions: This study is the first of its kind, and many of the analyses in the study yielded novel results. The study was designed to emulate the behavior of gamers during the weekend and other long gaming sessions. At this point, we are not able to determine the difference between the effects of gaming and behavior during gaming. Regardless, the results of this study suggest that healthy gamers can partake in long gaming sessions, with ample amounts of unhealthy foods and little rest, without acute impacts on health.

KEYWORDS

long gaming sessions; local area network party; biochemistry; cortisol; glucose; gaming; biochemical; blood sample; hematology; hematological; games; gamers; hemoglobin; adults; males; men; blood

Introduction

Video games have become a favorite pastime among children and adolescents. In the United States, 99% and 94% of boys and girls, respectively, play video games [1]. Video games are part of a larger category of sedentary activities linked to health issues such as physical inactivity, overeating, obesity, and diabetes [2-6]. Collegiate-level gamers have a comparable BMI to their nongaming peers but are less active with a higher body fat percentage, lower lean body mass, and lower bone mineral content [7]. This is particularly troublesome because studies report that children spend as much as 7-11 hours daily engaged in screen-based activities after school [8]. In addition, gaming has been linked to overeating and overconsumption of soft drinks as well as adverse health behaviors [9-12]. In a recent review, the authors found that while gaming may increase energy expenditure above baseline levels, gaming does not constitute physical activity. Energy expenditure may increase, but the activity level is comparable to that of standing or walking [13].

Previously published data from this study showed that the participants ingested an excessive number of calories from both food and drink [14]. During the study, the participants ingested an average of 6160 kcal from food and 1844 kcal from liquid sources. Additionally, the participants consumed 1354 mg of caffeine on average during the same period [14].

Previous studies of the impact of gaming on biochemical parameters have primarily addressed the acute effects of gaming. The literature is limited, and the study designs are very diverse. The parameters that have been investigated most thoroughly are blood glucose and cortisol. For blood glucose levels, gaming does not appear to increase glucose levels within the first 20 minutes of a gaming session [15,16]. Chaput et al [15] reported that gaming increased blood glucose levels after 40 minutes of a 60-minute gaming session.

Cortisol has been used as a marker of both physiological and mental stress during gaming [17-19]. Gaming may affect cortisol levels, but the nature of this relationship has yet to be elucidated. Oxford et al [17] reported that cortisol levels increase acutely when gamers compete against friends. The authors suggested that this was typical of male-male competition behavior [17].

While biochemical markers such as blood glucose and cortisol levels have been investigated during the last 2 decades, many biochemical markers pertaining to acute changes in the health of the human body have yet to be investigated. In this study, we aimed to investigate a broad array of biochemical markers to assess homeostasis, lipid metabolism, internal organ function, hematological balance, acid-base balance, and blood gases during long gaming sessions. This is the first time that most of the included parameters (except for glucose and cortisol) were investigated in gamers and during long gaming sessions. As

such, we did not know what to expect over the course of the study.

According to a recent literature review, knowledge of the effects of long video gaming sessions is minimal [13]. This exploratory study is the first of its kind in gaming. This is true for both the length of the gaming sessions, the physical setup, and the extensive testing.

A limitation of this study was the relatively small number of participants. This study was designed to realistically emulate the gaming behavior of young adults in a controlled setting in a hospital dining hall and an adjourning meeting room. The study was conducted as a local area network party for practical reasons. Four laboratory technicians worked at all times to sample, prepare, and analyze blood samples. A doctor and 2 investigators were also present throughout the gaming sessions. We tried to overcome the artificial situation of the event by discussing the setup with the participants before the study to create the most real-life-like experience (video clip [20]). The participants were asked to consume food and drink according to their wishes and what they would habitually consume [14]. The extensive testing in the study allowed us to observe short-term changes (within 6 hours), long-term changes during the duration of the gaming sessions, and follow-up after 1 week to determine whether any changes were longer lasting. The aim of this study was to investigate the effects of long gaming sessions on the biochemical parameters of healthy male adults.

Methods

Participants and Intervention

We have previously presented how the study was conducted and the physiological response to long gaming sessions [14]. In this paper, the results of an extensive collection of biochemical data are presented. We have adhered to the CONSORT (Consolidated Standards of Reporting Trials) statement regarding pilot and feasibility trials [21]. A CONSORT checklist can be found in [Multimedia Appendix 1](#).

In brief, 9 healthy male participants older than 18 years of age with vast gaming experience were enrolled. The mean age of the participants was 25.8 (SD 2.6) years, and the mean BMI was 24.8 (SD 2.9) years. All participants were either full-time students or employees [14]. According to the protocol (EudraCT 2019-004091-19), the plan was to enroll 6-9 gamers, with at least 6 gamers engaging in gaming for 48 hours. Participants were enrolled after they had called for participants from local e-sports clubs through e-sports instructors, online message boards, and word of mouth. The participants were recruited into 2 teams of 4 and 5 members.

The intervention consisted of two 18-hour gaming sessions interspersed with a 6-hour break. During the break, the participants had approximately 4 hours of sleep. After the last

gaming period, all participants underwent both physiological and biochemical tests.

Throughout the intervention, the participants had ad libitum access to food and drink. Before the study, participants provided lists of their preferred snacks and drinks. An assortment of chips, candies, cookies, buns, cold cuts, cereal, and fruit was available throughout the study. Additionally, an evening meal was provided at 6 PM during both gaming sessions: pizza on the first evening and hamburgers on the second evening. Participants had access to tap water, soda, energy drinks, coffee, milk, and chocolate milk.

The participants were instructed to avoid strenuous physical labor, cardiovascular exercise, alcohol, and junk food for 7 days before the study. Additionally, on the day of the study, the participants were instructed to stop food intake at noon and only drink water in case of thirst.

During the study, the participants were not restricted in any way regarding gaming. Specifically, all types of games across all genres and platforms were allowed. One participant brought his PlayStation, and as a new soccer game had just been released, participants played matches in teams or head-to-head. At other times, the participants played alone in certain games or played on the web with and against players who were not participants in the study. Especially during the last 18 hours of the study, the participants played first-person shooter games against each other on teams.

Participant Involvement

After the protocol was approved by the local ethics committee in February 2018, in total, 2 meetings were held with potential participants and local e-sports instructors, discussing the content of the protocol. In particular, food monitoring and blood sampling methods were changed in accordance with the wishes of the participants.

The participants suggested a “food diary” to monitor their caloric intake. This suggestion was incorporated into the protocol. First, it was suggested by the research team that a venous catheter be inserted for blood sampling throughout the study. Instead, the participants opted for multiple venipunctures. The changes suggested at the meetings were approved by the local ethics committee in an amendment before the study started.

Ethical Considerations

The North Denmark Region Committee approved the study protocol on Health Research Ethics (N-20180011; EudraCT 2019-004091). Each participant provided informed consent in writing twice, 2 weeks prior to the study and again on the day of the study. The study used a standardized consent form that stated that participation was voluntary and that participants could withdraw from the study at any time without reason or consequence. Primary consent was obtained after approval for the secondary analysis. The data were deidentified. Except for travel expenses, participants were not compensated for their participation.

Blood Sampling and Processing

When the participants arrived at the laboratory, baseline samples were collected, including a venous blood sample, a urine sample, and a saliva swab. Blood, urine, and saliva were collected from the participants every 6 hours. After the blood was drawn by venous puncture, it was taken to the laboratory. Biochemical analyses were performed immediately using an ABL800 FLEX Blood Gas Analyzer (Radiometer); other samples were centrifuged (at 3000 rpm/1875×g for 10 minutes), and the plasma was frozen at -80°C for later analysis. Blood samples were analyzed using a Cobas 8000 Modular Analyzer (Roche Applied Science) and a Sysmex XN-9000 Hematology Analyzer (Sysmex Europe, GmbH). In total, 18 parameters were analyzed on the Cobas 8000 (1 parameter is only presented in [Multimedia Appendices 2 and 3](#)), 22 parameters were analyzed on the Sysmex XN-9000 (16 parameters are only presented in [Multimedia Appendices 2 and 3](#)), and 18 parameters were analyzed using the ABL800 (11 parameters are only presented in [Multimedia Appendices 2 and 3](#)). An overview of the analyses by apparatus can be found in [Multimedia Appendix 2](#). The complete summary of the results is provided in [Multimedia Appendix 3](#). A total of 8 parameters were analyzed twice between the 3 machines (calcium [Ca], glucose, lactate, potassium [K], sodium [Na], bilirubin, creatinine, and hemoglobin [Hb]). All the tests were performed in accordance with the manufacturer’s instructions in a nationally accredited biochemistry laboratory (Department of Clinical Biochemistry, Aalborg University Hospital).

The following 30 parameters were measured and are presented in the Results section: glucose (mmol/L), lactate (mmol/L), cortisol (nmol/L), Ca (mmol/L), albumin-corrected calcium (mmol/L), K (mmol/L), Na (mmol/L), and chloride (Cl, mmol/L) were measured to assess homeostasis. Low-density lipoprotein (LDL) cholesterol (mmol/L), high-density lipoprotein cholesterol (mmol/L), total cholesterol (mmol/L), and triglyceride (TG, mmol/L) levels were measured to assess lipid metabolism. Alanine aminotransferase (ALT, U/L), albumin (g/L), alkaline phosphatase (ALP, U/L), bilirubin ($\mu\text{mol/L}$), creatinine ($\mu\text{mol/L}$), C-reactive protein (CRP, mg/L), and ferritin ($\mu\text{g/L}$) were measured to assess organ-specific markers. sO_2 (%), pCO_2 (kPa), pO_2 (kPa), pH, and standard bicarbonate concentration (mmol/L) were measured to assess acid-base balance and blood gases. Erythrocytes ($10^{12}/\text{L}$), erythrocyte volume fraction (EVF), Hb (mmol/L), mean cell Hb (10^{-15} , SE 0.0018 mol), mean cell volume ($10^{-15}/\text{L}$, SE 0.088/L), and platelets ($10^9/\text{L}$) were measured to assess the hematological markers.

Statistical Analyses

Linear mixed-effects models were used to analyze participants’ absolute changes in biochemical parameters throughout the study. This modeling method is a standard extension of linear regression models that controls for the random effects introduced by having paired data and is often used when analyzing studies with repeated measurements. We performed separate univariate analyses for each change in each biochemical parameter; time was the fixed effect, and participants were the random effect.

Additionally, we included an interaction term between time and period. This gave separate results for the first and second gaming periods.

We used a likelihood ratio test to test the statistical significance of the linear mixed effects models compared to a basic model, which included only random effects and thereby assumed no development over time. In total, 51 markers were analyzed (58 markers in total and 7 duplicates), and Bonferroni correction was applied by multiplying α (originally .05) by the number of tests. Statistical significance was set at $\alpha=.05$ before Bonferroni correction [22]. The assumptions of linear mixed-effects models were tested and fulfilled in all the analyses. Linearity and homoscedasticity were assessed by inspecting residual plots. The normality of the residuals was assessed by inspecting histograms and quantile-quantile plots. Given that this was an exploratory study, outliers were not removed. Due to the sample

size and number of tests, we did not perform post hoc tests comparing the data at individual time points.

Graphs illustrating changes over time were made for all variables, consisting of average values among participants for each sampling point, together with SEs presented as error bars from the mean. In addition, selected graphs were included to illustrate developments over time for specific parameters (Figures 1-4) that were not necessarily reflected in the linear mixed-effects models.

One participant's LDL cholesterol level decreased 3-fold below the minimum measuring range. The missing values were substituted by using the previous lowest value of the participant [23]. All the statistical analyses were performed using Microsoft Excel 2013 (Microsoft Corp) and RStudio (version 1.1.383; Posit, PBC). Linear mixed-effects models were generated using the *lme4* package [24].

Figure 1. Dehydration. (A) Albumin, (B) hemoglobin, (C) erythrocyte volume fraction (EVF), and (D) platelets.




Figure 2. Glucose development over time.




Figure 3. Cortisol development over time.





Figure 4. Triglyceride development over time.

Results

Overview

The linear mixed-effects models produced results with a mean baseline value at the start of the study and coefficients of change for the first and second gaming periods. Several parameters, such as albumin, Hb, EVF, and the platelet count, exhibited a pattern of high initial values, followed by a decrease after 6 hours and a continuous increase during the rest of the gaming period (Figure 1). The complete list of results of all analyses is available in [Multimedia Appendix 3](#).

Homeostasis

The mean glucose level of the participants was 4.39 (SD 0.07) mmol/L at baseline, which increased significantly by 0.24 (SD 0.07) mmol/L per 6 hours during the first period and by 0.38 (SD 0.07) mmol/L per 6 hours in the second period (Figure 2). The mean Na level was 141.9 (SE 0.12) mmol/L at baseline and decreased by 0.31 (SD 0.1) mmol/L per 6 hours during the second period. There were no changes during the first period. The mean Cl level was 105.9 (SE 0.14) mmol/L at baseline and decreased significantly by 0.54 (SD 0.14) mmol/L per 6 hours during the second period, and there were no significant changes during the first period. Lactate, Ca, K, and albumin-corrected calcium levels did not significantly change during the study. Cortisol levels also did not change significantly. However, cortisol levels appeared to be affected by long gaming sessions and a lack of rest during the study. A deviation from this pattern is usually associated with daily fluctuations in cortisol levels (Figure 3). During a normal cortisol rhythm, cortisol levels are

at their lowest level at midnight before increasing to their peak value at 6 AM, followed by a steady decline toward midnight.

Lipid Metabolism

Total cholesterol, LDL, high-density lipoprotein, and TG did not significantly change during the study. However, the TG levels during the study showed a distinct pattern: TG levels increased after evening meals but returned to baseline after 6 hours (Figure 4).

Organ-Specific Markers

The mean ALP level was 77.2 (SE 0.63) U/L at baseline and increased during the second period by 3.8 (SD 0.63) U/L per 6 hours. There was no significant change during the first period. The mean bilirubin level was 9.9 (SE 0.22) $\mu\text{mol/L}$ at baseline, decreased by 1.15 (SD 0.22) $\mu\text{mol/L}$ per 6 hours during the first period and decreased by 1.05 (SD 0.22) $\mu\text{mol/L}$ per 6 hours during the second period. The mean creatinine level was 93.6 (0.56) $\mu\text{mol/L}$ at baseline, decreased by 2.9 (SD 0.56) $\mu\text{mol/L}$ every 6 hours during the first period and decreased by 1.15 (SD 0.56) $\mu\text{mol/L}$ every 6 hours during the second period. Albumin, CRP, ferritin, and ALT did not significantly change during the gaming sessions.

Acid-Base Balance and Blood Gases

sO_2 , pCO_2 , pO_2 , pH, and standard bicarbonate concentration did not significantly change during the study.

Hematology

The mean Hb level was 9.18 (SE 0.025) mmol/L at baseline, and it decreased by 0.055 (SD 0.025) mmol/L every 6 hours during the first period. However, during the second period, Hb

increased by 0.027 (SD 0.025) mmol/L per 6 hours. The mean platelet count was $258 \times 10^9/L$ (SE $1.1 \times 10^9/L$) at baseline and increased by $6.9 \times 10^9/L$ (SD $1.1 \times 10^9/L$) per 6 hours during the second period. During the first session, there were no significant changes. Erythrocytes, EVF, mean cell Hb, and mean cell volume did not significantly change during the study.

Discussion

Principal Findings

This study is the first of its kind regarding the design and number of biochemical analyses in the gaming population. The fact that the study stretches over 42 hours of continuous measurements makes it unique compared to the current literature on gaming science. The study applies to recreational gamers who play with various levels of seriousness but lack a singular focus on the competition associated with e-sports. Overall, the effect of gaming on standard biochemical parameters in healthy male adults is limited. Significant changes were found in 12 of the 51 parameters. Most of the results of biochemical tests are novel findings in a gamer population [13]. It is not surprising that most parameters were unaffected by the intervention, but the high number of examined parameters added to our understanding of the effects of gaming. The participants had a large intake of calories throughout the gaming sessions (especially during the first gaming sessions) [14]. We found small changes in the levels of several biochemical and hematological biomarkers, but all the levels were within the normal range. Overall, the results of this exploratory study suggest that, from a biochemical and hematological standpoint, the health of male adults is not altered in the short term by long gaming sessions.

Dehydration

The development of several parameters, including albumin, ALP, creatinine, Ca, ferritin, Hb, erythrocytes, EVF, K, and platelets, over time suggested that the participants were relatively dehydrated at the start of the study and rehydrated within the first 6 hours, followed by relative dehydration during the remainder of the study (Figure 1). Dehydration, despite a 3-L fluid intake per 18-hour gaming session, was most likely aided by the participants' caffeine intake, as a large intake of caffeine during nonstrenuous activities can cause increased diuresis [25].

Homeostasis

Blood glucose levels increased consistently during each of the gaming sessions (Figure 2). This development agrees with the findings of Chaput et al [15], but the changes in this study occurred over a much longer period. The changes in blood glucose levels cannot be attributed to dehydration, as this pathway is under tight hormonal control [26]. The glucose levels during the second session increased even though the participants had little energy intake. This could be due to the unexpectedly high levels of cortisol present at the same time (Figure 3), triggering the release of glucose from body stores. The glucose levels never exceeded the normal range and returned to baseline levels 1 week after the second long gaming session. Short- and long-term changes in blood glucose levels in gamers who

regularly participate in long gaming sessions need further investigation.

The development of cortisol during the study differed from what would be expected (Figure 3). Cortisol is typically at its lowest level at midnight and increases sharply at 6 AM to reach a maximum between 6 and 10 AM, after which a sharp decline is expected [27]. The continued increase at noon during both sessions is surprising. Typically, the cortisol concentration decreases throughout the day and evening, reaching its nadir at midnight. The participants slept between noon (after the first session) and 6 PM. The 6 PM cortisol value could have been influenced by the waking cortisol response, which is related to the circadian rhythm [28]. Multiple factors potentially contribute to the disruption of the regular cortisol rhythm. Going into the study, the participants all adhered to a structured sleep pattern due to daytime employment or attending university classes. During both gaming sessions, we observed a rise in cortisol levels from 6 AM until noon. Gaming could be the cause of this change, as the level of alertness (or "stress") in gaming potentially requires the continuous activity of the hypothalamic-pituitary-adrenal axis [28,29]. An increase in cortisol based on light intensity has been described, and light from monitors could be a factor in the changes observed [30]. Nightshift workers who are habitually awake during the night exhibit markedly lower morning cortisol levels [31]. The sharp decrease in cortisol levels at midnight (both nights) indicated that the regular diurnal slope was not completely changed by the weekend during extreme gaming.

Lipid Metabolism

Gaming and sedentary behavior have been associated with obesity and increased cholesterol and TG levels [4], as they displace other nonsedentary activities [32]. Based on standard biochemistry, cholesterol and TG were unaffected by long gaming sessions and unrestricted food intake. However, TG levels sharply increased after mealtimes, which were normalized 6-12 hours after each meal. This increase was much greater than usual [33,34]. The sharp increase in TG was most likely the result of the high fat content of the ingested foods [14].

Organ-Specific Markers

Overall, this study does not indicate that long gaming sessions impact the kidneys or livers of gamers. Creatinine levels decreased slightly during the study, possibly because of reduced physical activity. A decrease in creatinine at this level is not known to influence health.

Bilirubin and ALP levels decreased significantly during the study. This decrease, while significant, is not associated with any known pathophysiology.

Additionally, the large amounts of caffeine ingested may have lowered the bilirubin levels [35]. In experimental models, caffeine has been shown to have antifibrogenic, anti-inflammatory, and antioxidant effects that potentially exert liver protection [36-39].

We expected an increase in ALT based on food consumption during the study [40]. Likewise, we suspected that the inflammatory parameters would change due to the large intake

of unhealthy food and drinks. As CRP, ferritin, and ALT are sensitive to changes in behavior (strenuous exercise, alcohol consumption, and excessive food intake), their lack of change during the study indicates a minor or no impact of gaming behavior.

Acid-Base Balance and Blood Gases

Our results showed that gaming did not affect the pH balance or its regulatory systems. Furthermore, the pO₂ in the venous blood did not significantly change throughout the trial. This finding agrees with the literature suggesting that sedentary gaming is not physically demanding [13].

Hematology

Hematological parameters have not previously been examined in gamers during long gaming sessions. The Hb concentration decreased significantly during the first 18 hours of gaming and increased during the second 18-hour session. These changes are in accordance with the changes in hydration status outlined earlier. Thrombocyte counts increased throughout the study, especially in the second 18-hour session. Despite the significant changes, the parameters were always within the normal range. Overall, this exploratory study does not suggest a need for further investigation into the association between hematological parameters and gaming.

Strengths and Limitations

The participants were all male individuals in their 20s, and there were only 9 participants. Based on these results, it is not possible

to determine whether gaming has a different or even harmful effect on children, adolescents, or female individuals. The study was conducted in a hospital and not in the familiar environments of the participants. This may have caused participants to alter their behavior in ways we cannot determine. The easy availability of food and snacks may not relate entirely to how long gaming sessions are conducted at home or at local area network parties. We cannot rule out the possibility that overeating occurred, but the available food and drinks were only present at the behest of the participants. During the first 6-12 hours of the study, overeating was perhaps a factor, but during the second session, the participants had a markedly lower intake of calories from both food and drinks [14]. On the basis of this study, we are not able to determine whether the excessive intake of calories was due to convenience or to sustain a high level of performance during strenuous gaming.

Conclusions

This study is the first of its kind, and many of the analyses in the study yielded novel results. The study was designed to emulate the behavior of gamers during the weekend and other long gaming sessions. At this point, we are not able to determine the difference between the effects of gaming and behavior during gaming. Regardless, the results of this study suggest that healthy gamers can partake in long gaming sessions, with ample amounts of unhealthy foods and little rest, without acute impacts on health.

Acknowledgments

The authors thank the participants for their input and participation in this study. This study would not have been possible without the support of Aalborg University Hospital, which generously provided facilities and technical support. Additionally, the authors thank the laboratory staff at the Department of Molecular Diagnostics and the Department of Clinical Biochemistry for their swift and tireless work during the study. This study was funded by grants from “Brødrene Hartmanns Fond” and “Tømmerhandler Johannes Fogs Fond.” The funding sources had no involvement in the study design, data collection, or data interpretation.

Data Availability

Data from this study are not available in a public repository. It can be requested from the corresponding author. The study garnered national media attention in Denmark when it was conducted, and several participants were interviewed through radio and television. The data included physical characteristics, employment status, and other personal information. Due to the low number of participants, it would potentially be possible to identify individual participants' data.

Conflicts of Interest

None declared.

Multimedia Appendix 1

CONSORT checklist.

[\[PDF File \(Adobe PDF File\), 94 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Biochemical markers by analysis instrument.

[\[DOCX File , 16 KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Overview of biochemical markers.

[DOCX File , 55 KB-Multimedia Appendix 3]

References

1. Lenhart J, Kahne J, Middaugh E, Macgill AR, Evans C, Vitak J. Teens, video games, and civics. Pew Research Center. 2008. URL: <https://www.pewresearch.org/internet/2008/09/16/teens-video-games-and-civics/> [accessed 2024-04-19]
2. Turel O, Romashkin A, Morrison KM. Health outcomes of information system use lifestyles among adolescents: videogame addiction, sleep curtailment and cardio-metabolic deficiencies. *PLoS One*. 2016;11(5):e0154764. [FREE Full text] [doi: [10.1371/journal.pone.0154764](https://doi.org/10.1371/journal.pone.0154764)] [Medline: [27149512](https://pubmed.ncbi.nlm.nih.gov/27149512/)]
3. Turel O, Romashkin A, Morrison KM. A model linking video gaming, sleep quality, sweet drinks consumption and obesity among children and youth. *Clin Obes*. 2017;7(4):191-198. [doi: [10.1111/cob.12191](https://doi.org/10.1111/cob.12191)] [Medline: [28320073](https://pubmed.ncbi.nlm.nih.gov/28320073/)]
4. Vella CA, Taylor K, Nelson MC. Associations of leisure screen time with cardiometabolic biomarkers in college-aged adults. *J Behav Med*. 2020;43(6):1014-1025. [FREE Full text] [doi: [10.1007/s10865-020-00161-2](https://doi.org/10.1007/s10865-020-00161-2)] [Medline: [32451650](https://pubmed.ncbi.nlm.nih.gov/32451650/)]
5. Vicente-Rodríguez G, Rey-López JP, Martín-Matillas M, Moreno LA, Wärnberg J, Redondo C, et al. Television watching, videogames, and excess of body fat in Spanish adolescents: the AVENA study. *Nutrition*. 2008;24(7-8):654-662. [doi: [10.1016/j.nut.2008.03.011](https://doi.org/10.1016/j.nut.2008.03.011)] [Medline: [18472399](https://pubmed.ncbi.nlm.nih.gov/18472399/)]
6. Vuong B, Rogers R, Corriveau N, Foti A, Pew A, Wei WC, et al. Passive screen time associated with unhealthy dietary consumption and physiological characteristics: a closer look at childhood behaviors. *J Am Coll Cardiol*. 2014;63(12):A1292. [doi: [10.1016/s0735-1097\(14\)61292-4](https://doi.org/10.1016/s0735-1097(14)61292-4)]
7. DiFrancisco-Donoghue J, Werner WG, Douris PC, Zwibel H. Esports players, got muscle? Competitive video game players' physical activity, body fat, bone mineral content, and muscle mass in comparison to matched controls. *J Sport Health Sci*. 2022;11(6):725-730. [FREE Full text] [doi: [10.1016/j.jshs.2020.07.006](https://doi.org/10.1016/j.jshs.2020.07.006)] [Medline: [32711155](https://pubmed.ncbi.nlm.nih.gov/32711155/)]
8. Strasburger VC, Jordan AB, Donnerstein E. Children, adolescents, and the media: health effects. *Pediatr Clin North Am*. 2012;59(3):533-587, vii. [doi: [10.1016/j.pcl.2012.03.025](https://doi.org/10.1016/j.pcl.2012.03.025)] [Medline: [22643165](https://pubmed.ncbi.nlm.nih.gov/22643165/)]
9. Henderson M, Benedetti A, Barnett TA, Mathieu ME, Deladoëy J, Gray-Donald K. Influence of adiposity, physical activity, fitness, and screen time on insulin dynamics over 2 years in children. *JAMA Pediatr*. 2016;170(3):227-235. [FREE Full text] [doi: [10.1001/jamapediatrics.2015.3909](https://doi.org/10.1001/jamapediatrics.2015.3909)] [Medline: [26857733](https://pubmed.ncbi.nlm.nih.gov/26857733/)]
10. Siervo M, Gan J, Fewtrell MS, Cortina-Borja M, Wells JCK. Acute effects of video-game playing versus television viewing on stress markers and food intake in overweight and obese young men: a randomised controlled trial. *Appetite*. 2018;120:100-108. [doi: [10.1016/j.appet.2017.08.018](https://doi.org/10.1016/j.appet.2017.08.018)] [Medline: [28843974](https://pubmed.ncbi.nlm.nih.gov/28843974/)]
11. Mario S, Hannah C, Jonathan WCK, Jose L. Frequent video-game playing in young males is associated with central adiposity and high-sugar, low-fibre dietary consumption. *Eat Weight Disord*. 2014;19(4):515-520. [doi: [10.1007/s40519-014-0128-1](https://doi.org/10.1007/s40519-014-0128-1)] [Medline: [24821488](https://pubmed.ncbi.nlm.nih.gov/24821488/)]
12. Gheller BJB, Totosy de Zepetnek JO, Welch JM, Rossiter MD, Luhovyy B, Brett NR, et al. Effect of video game playing and a glucose preload on subjective appetite, subjective emotions, and food intake in overweight and obese boys. *Appl Physiol Nutr Metab*. 2019;44(3):248-254. [doi: [10.1139/apnm-2018-0281](https://doi.org/10.1139/apnm-2018-0281)] [Medline: [30096245](https://pubmed.ncbi.nlm.nih.gov/30096245/)]
13. Krarup KB, Krarup HB. The physiological and biochemical effects of gaming: a review. *Environ Res*. 2020;184:109344. [doi: [10.1016/j.envres.2020.109344](https://doi.org/10.1016/j.envres.2020.109344)] [Medline: [32199319](https://pubmed.ncbi.nlm.nih.gov/32199319/)]
14. Krarup KB, Rantanen JM, Mørk M, Pedersen IS, Christensen JH, Kristensen SR, et al. The physiological and cardiologic effects of long video gaming sessions in adult males. *Sports Med Int Open*. 2022;6(1):E39-E46. [FREE Full text] [doi: [10.1055/a-1858-8436](https://doi.org/10.1055/a-1858-8436)] [Medline: [35874049](https://pubmed.ncbi.nlm.nih.gov/35874049/)]
15. Chaput JP, Visby T, Nyby S, Klingenberg L, Gregersen NT, Tremblay A, et al. Video game playing increases food intake in adolescents: a randomized crossover study. *Am J Clin Nutr*. 2011;93(6):1196-1203. [FREE Full text] [doi: [10.3945/ajcn.110.008680](https://doi.org/10.3945/ajcn.110.008680)] [Medline: [21490141](https://pubmed.ncbi.nlm.nih.gov/21490141/)]
16. Wang X, Perry AC. Metabolic and physiologic responses to video game play in 7- to 10-year-old boys. *Arch Pediatr Adolesc Med*. 2006;160(4):411-415. [FREE Full text] [doi: [10.1001/archpedi.160.4.411](https://doi.org/10.1001/archpedi.160.4.411)] [Medline: [16585487](https://pubmed.ncbi.nlm.nih.gov/16585487/)]
17. Oxford J, Ponzi D, Geary DC. Hormonal responses differ when playing violent video games against an ingroup and outgroup. *Evol Hum Behav*. 2010;31(3):201-209. [doi: [10.1016/j.evolhumbehav.2009.07.002](https://doi.org/10.1016/j.evolhumbehav.2009.07.002)]
18. Kindermann H, Javor A, Reuter M. Playing counter-strike versus running: the impact of leisure time activities and cortisol on intermediate-term memory in male students. *Cogn Syst Res*. 2016;40:1-7. [doi: [10.1016/j.cogsys.2016.01.002](https://doi.org/10.1016/j.cogsys.2016.01.002)]
19. Ivarsson M, Anderson M, Akerstedt T, Lindblad F. Playing a violent television game does not affect saliva cortisol. *Acta Paediatr*. 2009;98(6):1052-1053. [doi: [10.1111/j.1651-2227.2009.01271.x](https://doi.org/10.1111/j.1651-2227.2009.01271.x)] [Medline: [19397536](https://pubmed.ncbi.nlm.nih.gov/19397536/)]
20. Dieckmann L. TV2 Nord, 2018. TV2 Nord. Sep 29, 2018. URL: <https://vimeo.com/906653219> [accessed 2024-04-17]
21. Eldridge SM, Chan CL, Campbell MJ, Bond CM, Hopewell S, Thabane L, et al. CONSORT 2010 statement: extension to randomised pilot and feasibility trials. *BMJ*. 2016;355:i5239. [FREE Full text] [doi: [10.1136/bmj.i5239](https://doi.org/10.1136/bmj.i5239)] [Medline: [27777223](https://pubmed.ncbi.nlm.nih.gov/27777223/)]
22. Jafari M, Ansari-Pour N. Why, when and how to adjust your P values? *Cell J*. 2019;20(4):604-607. [FREE Full text] [doi: [10.22074/cellj.2019.5992](https://doi.org/10.22074/cellj.2019.5992)] [Medline: [30124010](https://pubmed.ncbi.nlm.nih.gov/30124010/)]
23. Giskeødegård GF, Lydersen S. Measurements below the detection limit. *Tidsskr Nor Laegeforen*. 2022;142(13):62352. [FREE Full text] [doi: [10.4045/tidsskr.22.0439](https://doi.org/10.4045/tidsskr.22.0439)] [Medline: [36164797](https://pubmed.ncbi.nlm.nih.gov/36164797/)]

24. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Soft.* 2015;67(1):1-48. [doi: [10.18637/jss.v067.i01](https://doi.org/10.18637/jss.v067.i01)]
25. Seal AD, Bardis CN, Gavrieli A, Grigorakis P, Adams JD, Arnaoutis G, et al. Coffee with high but not low caffeine content augments fluid and electrolyte excretion at rest. *Front Nutr.* 2017;4:40. [FREE Full text] [doi: [10.3389/fnut.2017.00040](https://doi.org/10.3389/fnut.2017.00040)] [Medline: [28868290](https://pubmed.ncbi.nlm.nih.gov/28868290/)]
26. Röder PV, Wu B, Liu Y, Han W. Pancreatic regulation of glucose homeostasis. *Exp Mol Med.* 2016;48(3):e219. [FREE Full text] [doi: [10.1038/emm.2016.6](https://doi.org/10.1038/emm.2016.6)] [Medline: [26964835](https://pubmed.ncbi.nlm.nih.gov/26964835/)]
27. Ryan R, Booth S, Spathis A, Mollart S, Clow A. Use of salivary diurnal cortisol as an outcome measure in randomised controlled trials: a systematic review. *Ann Behav Med.* 2016;50(2):210-236. [FREE Full text] [doi: [10.1007/s12160-015-9753-9](https://doi.org/10.1007/s12160-015-9753-9)] [Medline: [27007274](https://pubmed.ncbi.nlm.nih.gov/27007274/)]
28. Kudielka BM, Gierens A, Hellhammer DH, Wüst S, Schlotz W. Salivary cortisol in ambulatory assessment—some dos, some don'ts, and some open questions. *Psychosom Med.* 2012;74(4):418-431. [doi: [10.1097/PSY.0b013e31825434c7](https://doi.org/10.1097/PSY.0b013e31825434c7)] [Medline: [22582339](https://pubmed.ncbi.nlm.nih.gov/22582339/)]
29. Hellhammer DH, Wüst S, Kudielka BM. Salivary cortisol as a biomarker in stress research. *Psychoneuroendocrinol.* 2009;34(2):163-171. [doi: [10.1016/j.psyneuen.2008.10.026](https://doi.org/10.1016/j.psyneuen.2008.10.026)] [Medline: [19095358](https://pubmed.ncbi.nlm.nih.gov/19095358/)]
30. Petrowski K, Buehrer S, Niedling M, Schmalbach B. The effects of light exposure on the cortisol stress response in human males. *Stress.* 2021;24(1):29-35. [FREE Full text] [doi: [10.1080/10253890.2020.1741543](https://doi.org/10.1080/10253890.2020.1741543)] [Medline: [32160826](https://pubmed.ncbi.nlm.nih.gov/32160826/)]
31. Mirick DK, Bhatti P, Chen C, Nordt F, Stanczyk FZ, Davis S. Night shift work and levels of 6-sulfatoxymelatonin and cortisol in men. *Cancer Epidemiol Biomarkers Prev.* 2013;22(6):1079-1087. [FREE Full text] [doi: [10.1158/1055-9965.EPI-12-1377](https://doi.org/10.1158/1055-9965.EPI-12-1377)] [Medline: [23563887](https://pubmed.ncbi.nlm.nih.gov/23563887/)]
32. Turel O. Videogames and guns in adolescents: tests of a bipartite theory. *Comput Human Behav.* 2020;109:106355. [FREE Full text] [doi: [10.1016/j.chb.2020.106355](https://doi.org/10.1016/j.chb.2020.106355)] [Medline: [32372846](https://pubmed.ncbi.nlm.nih.gov/32372846/)]
33. Tiihonen K, Rautonen N, Alhoniemi E, Ahotupa M, Stowell J, Vasankari T. Postprandial triglyceride response in normolipidemic, hyperlipidemic and obese subjects—the influence of polydextrose, a non-digestible carbohydrate. *Nutr J.* 2015;14:23. [FREE Full text] [doi: [10.1186/s12937-015-0009-0](https://doi.org/10.1186/s12937-015-0009-0)] [Medline: [25889643](https://pubmed.ncbi.nlm.nih.gov/25889643/)]
34. Mørk M, Nielsen MH, Bæk R, Jørgensen MM, Pedersen S, Kristensen SR. Postprandial increase in blood plasma levels of tissue factor-bearing (and other) microvesicles measured by flow cytometry: fact or artifact? *TH Open.* 2018;2(2):e147-e157. [FREE Full text] [doi: [10.1055/s-0038-1642021](https://doi.org/10.1055/s-0038-1642021)] [Medline: [31249938](https://pubmed.ncbi.nlm.nih.gov/31249938/)]
35. Modi AA, Feld JJ, Park Y, Kleiner DE, Everhart JE, Liang TJ, et al. Increased caffeine consumption is associated with reduced hepatic fibrosis. *Hepatology.* 2010;51(1):201-209. [FREE Full text] [doi: [10.1002/hep.23279](https://doi.org/10.1002/hep.23279)] [Medline: [20034049](https://pubmed.ncbi.nlm.nih.gov/20034049/)]
36. Amer MG, Mazen NF, Mohamed AM. Caffeine intake decreases oxidative stress and inflammatory biomarkers in experimental liver diseases induced by thioacetamide: biochemical and histological study. *Int J Immunopathol Pharmacol.* 2017;30(1):13-24. [FREE Full text] [doi: [10.1177/0394632017694898](https://doi.org/10.1177/0394632017694898)] [Medline: [28281876](https://pubmed.ncbi.nlm.nih.gov/28281876/)]
37. Helal MG, Ayoub SE, Elkashefand WF, Ibrahim TM. Caffeine affects HFD-induced hepatic steatosis by multifactorial intervention. *Hum Exp Toxicol.* 2018;37(9):983-990. [FREE Full text] [doi: [10.1177/0960327117747026](https://doi.org/10.1177/0960327117747026)] [Medline: [29249184](https://pubmed.ncbi.nlm.nih.gov/29249184/)]
38. Marventano S, Salomone F, Godos J, Pluchinotta F, Del Rio D, Mistretta A, et al. Coffee and tea consumption in relation with non-alcoholic fatty liver and metabolic syndrome: a systematic review and meta-analysis of observational studies. *Clin Nutr.* 2016;35(6):1269-1281. [doi: [10.1016/j.clnu.2016.03.012](https://doi.org/10.1016/j.clnu.2016.03.012)] [Medline: [27060021](https://pubmed.ncbi.nlm.nih.gov/27060021/)]
39. Molloy JW, Calcagno CJ, Williams CD, Jones FJ, Torres DM, Harrison SA. Association of coffee and caffeine consumption with fatty liver disease, nonalcoholic steatohepatitis, and degree of hepatic fibrosis. *Hepatology.* 2012;55(2):429-436. [FREE Full text] [doi: [10.1002/hep.24731](https://doi.org/10.1002/hep.24731)] [Medline: [21987293](https://pubmed.ncbi.nlm.nih.gov/21987293/)]
40. Kechagias S, Ernerson A, Dahlqvist O, Lundberg P, Lindström T, Nystrom FH, et al. Fast Food Study Group. Fast-food-based hyper-alimentation can induce rapid and profound elevation of serum alanine aminotransferase in healthy subjects. *Gut.* 2008;57(5):649-654. [FREE Full text] [doi: [10.1136/gut.2007.131797](https://doi.org/10.1136/gut.2007.131797)] [Medline: [18276725](https://pubmed.ncbi.nlm.nih.gov/18276725/)]

Abbreviations

- ALP:** alkaline phosphatase
- ALT:** alanine aminotransferase
- Ca:** calcium
- Cl:** chloride
- CONSORT:** Consolidated Standards of Reporting Trials
- CRP:** C-reactive protein
- EVF:** erythrocyte volume fraction
- Hb:** hemoglobin
- K:** potassium
- LDL:** low-density lipoprotein
- Na:** sodium

TG: triglyceride

Edited by T de Azevedo Cardoso; submitted 16.02.23; peer-reviewed by X Yao, M Birk; comments to author 15.11.23; revised version received 30.01.24; accepted 02.04.24; published 08.07.24

Please cite as:

Krarup KB, Riis J, Mørk M, Nguyen HTT, Søkilde Pedersen I, Risom Kristensen S, Handberg A, Krarup HB

Biochemical Changes in Adult Male Gamers During Prolonged Gaming: Pilot Study

Interact J Med Res 2024;13:e46570

URL: <https://www.i-jmr.org/2024/1/e46570>

doi: [10.2196/46570](https://doi.org/10.2196/46570)

PMID: [38976326](https://pubmed.ncbi.nlm.nih.gov/38976326/)

©Kasper Bygum Krarup, Johannes Riis, Morten Mørk, Hien Thi Thu Nguyen, Inge Søkilde Pedersen, Søren Risom Kristensen, Aase Handberg, Henrik Bygum Krarup. Originally published in the Interactive Journal of Medical Research (<https://www.i-jmr.org/>), 08.07.2024. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Interactive Journal of Medical Research, is properly cited. The complete bibliographic information, a link to the original publication on <https://www.i-jmr.org/>, as well as this copyright and license information must be included.